International Journal of Zoology and Applied Biosciences Volume 3, Issue 2, pp: 174-177, 2018 https://doi.org/10.5281/zenodo.1314040



Research Article

HISTOPATHOLOGICAL ALTERATIONS IN THE INTESTINE OF FISH CHANNA PUNCTATUS EXPOSED TO SUBLETHAL CONCENTRATIONS OF ATRAZINE

S. Jain, Mansoor majeed lone* and Shama Rukshar

Department of Zoology, Govt. M. H. College of Home Science and Science for Women, Napier Town, Jabalpur-482002. Madhya Pradesh, India

Article History: Received 8th April 2018; Accepted 20th April 2018; Published 22nd April 2018

ABSTRACT

Atrazine commonly used as a herbicide, contaminates the aquatic ecosystem as a toxic pollutant from agricultural and domestic washouts. The aim of the present study was to investigate the toxic effect of Atrazine on intestine tissues in the fish *Channa punctatus* exposed to sublethal concentration of 5 mg/L, 10 mg/L and 15 mg/L for 15 and 30 days respectively. The histological observations of the intestine showed deleterious effect of Atrazine exposure time, the intestine of *C. punctatus* showed many strong changes viz, rupturing of longitudinal and circular muscle layers, reduction of villi, Necrosis at the tips of villi and highly shrinkage of sub mucosa at the tips of the villi were observed, Distortion of goblet cells, Blood vessels and Loss of structural integrity of mucosal folds were also seen in the intestine of *C. punctatus* at higher exposure of Atrazine. This clearly demonstrates that the sublethal concentrations of Atrazine have a deleterious effect on the intestine of *C. punctatus*. So, the herbicide Atrazine can be considered as a potent toxic pollutant capable of destroying the balance of the aquatic ecosystem.

Keywords: Channa Punctatus, Atrazine, Intestine, Pollutant, Herbicide.

INTRODUCTION

Large-scale application of pesticides to crops and plants may contribute to the presence of toxic substances in the environment. These toxicants through winds, floods, rains and surface run off reach unrestricted areas like ponds, springs and rivers thus producing an adverse impact on the aquatic biota including fishes (Rahmathullah et al., 2003). Atrazine (2-chloro-4-ethylamino-6-isopropylamino-1, 3, 5triazine) is one of the most widely used triazine herbicides in the world. It is used to control broad- leaf weeds in corn or crops, including green vegetables, weeds in maize, sorghum, asparagus, citrus, sugar cane, bananas, coffee, oil palms and grass land/forestry (Cui et al., 2002). The major uses being maize and sorghum. It is also used in combination with many other herbicides (Alvarez & Fuiman, 2005; Roberts et al., 1998; Steinberg, 1997; Zhou et al., 2008).

After spraying on crops, it can enter watercourses, because of its high mobility through soil (Waring & Moore,

2004). Hussain *et al.* (1996) pointed out that Atrazine reaches aquatic environments due to proximities of the agricultural country sides to the water places, or directly due to the careless application in such environments. After reaching the environment and ultimately affect non-target organisms such as fish (Bhatnagar & Srivastava, 1992).

In the present study, the freshwater teleost fish, *Channa punctatus* locally called karra is an important and popular fish species in the water bodies located in Jabalpur regions of Madhya Pradesh (India) is selected. This region has been traditionally well developed agricultural area mostly cultivating irrigated wet crops. In view of the incidence of herbs and pests, a variety of herbicides and other chemicals are used on crops for controlling the agriculture pests. The surface runoff from these areas during monsoon flow into the ponds and affect the aquatic fauna mostly fish.

The purpose of present study was to access the Histopathological changes in the intestine of *C. punctatus*

exposed to atrazine which is commonly used to control the herbs of several agricultural fields in India.

MATERIAL AND METHODS

Live specimens of test organism C. punctatus were purchased from Hanumantal fish market (Jabalpur) M.P. Average length and weight of the experimental fish was 10 to 14 cm in length and 50 to 80 gms in weight. The experiments were conducted in the laboratory using glass aquarium ($40 \times 27 \times 26$ cm). The fishes were dipped in tubs and were disinfected with 0.01% of KMnO₄ solution and washed thoroughly to prevent dermal infection and the fishes were maintained in aquariums with dechlorinated water which was continuously aerated. These fishes were acclimatized in the laboratory for two weeks prior to the experimentation.

For sub-acute toxicity, the various sub lethal concentrations of Atrazine viz. 5mg/L, 10 mg/L and 15 mg/L for 15 and 30 days respectively were given. Sets of twelve fishes were transferred to four different aquaria. Out of four, one served as control (contained only dechlorinated water) and rest three contained different concentrations of Atrazine. Six fishes were taken out from each aquarium after Fifteen days and the rest of the fishes were taken out after 30 days.

The fish were sacrificed and intestine was removed for histological examination. The tissues were washed in saline water and preserved in plastic vials with Bouin's fluid for at least 18 hours. The intestine was then dehydrated, embedded in paraffin wax. The embedded blocks were sectioned (6 μ) by a microtome machine (Buffalo. N. Y. 14215. U. S. A.) Fixed and prepared slides were kept for one night. Then the sections were stained with haematoxylin and eosin and were then mounted for later examination. Changes induced by treatment in the intestinal tissues were photographed and analyzed by light microscope at 10 X eye piece magnification and 40X objective magnification (Olympus model U-CMAD3) with Camera attachment of Samsung (model SDC-313B).

RESULT AND DISCUSSION

The innermost layer is mucosa consists of (CE) columnar epithelium with columnar cells, (GC) goblet cells, and (LP) Lamina propria, Inner to the muscle layer is (SM) sub mucosa, which consists of connective tissue, blood vessels and nerve. It permits mobility of the mucosa. Muscular is below the (S) serosa and it consisting of inner layer of circularly oriented and outer layer of longitudinally oriented smooth muscle, circular muscle layer (CML) and longitudinal layer (LML). The functions are mixing of food with digestive enzymes.

Adventitia is relatively dense layer of areolar connective tissue often blending with connective tissue of

surrounding structures ,when covered by the peritoneum it is also called a (S) Serosa, containing blood vessels, nerves, and lymphatics also consists of flattened cell covering the outer boundary. The histological sections of Figure 1 the control intestine of C. punctatus consists of four layers: Mucosa, sub mucosa, muscularis, adventia. Figure 2. At lower concentrations 5 mg/L for 15 days not many changes were observed, Section of intestine showing loosely arranged of circular muscle layer and necrosis of epithelial cells were also observed. No changes in villi and longitudinal muscular layer were noticed (Figure 3 and 4). However higher concentration of Atrazine 10 mg/L and 15 mg/L for 15 days showed many changes rupturing of muscular layers both and circular muscle layer, Necrosis were prominent, reduction of villi, Reductions in the length of intestinal fold were obvious Figure 5. Slight changes in Intestine were observed after exposure to 5 mg/L for 30 days, degeneration of serosa, vacuolization in certain places and loosely arrangement of longitudinal muscle were also observed, Figure 6 and 7. After exposure to 10 mg/L and 15 mg/L for 30 days, the intestine of C. punctatus showed many strong changes viz, rupturing of longitudinal and circular muscle layers, reduction of villi, Necrosis at the tips of villi and highly shrinkage of sub mucosa at the tips of the villi were observed, Distortion of goblet cells, Blood vessels and Loss of structural integrity of mucosal folds were also seen in the intestine of C. punctatus at higher exposure of Atrazine.

Intestine is the most important part of the fish alimentary canal and plays an important role in digestion and absorption of food materials as well as considered as a sensitive organ for toxicity assessment of xenobiotic substances in fish species as they are directly exposed to complex mixture of toxic substances via ingestion of contaminated food stuffs or indirectly via blood and/or (Muniyan, 1999). In the present histopathology of the intestine of C. punctatus was studied after Atrazine intoxication. A number of studies on histopathological effects of different pesticides on fish intestine have been reported by Velmurugan et al. (2007) who reported degeneration in the tip of villi, loss of structural integrity in mucosal folds, hypertrophy, vacuolation and necrosis in Cyprinus carpio and Cirrhinus mrigala after exposed to atrazine. Sharma et al., (2001) also reported similar results in fish C. Batrachus and C. mrigala after pesticidal exposure. Ravanaiah & Murthy, (2010) also noticed vacuolations, lesions in villi and serosa layer, necrosis, congestion in blood capillaries and severe mucus secretion in Tilapia mossambica when exposed to industrial pollutants. Yildirim et al. (2006) also reported similar result in fish Oreochromis niloticus after exposed to pesticide. Reports on degenerative changes and rupture in tip of villi, loss of structural integrity of mucosal folds and degeneration and necrosis of submucosa in the intestine of C. punctatus after the exposure to carbofuran was also found by Muley et al. (1996).




Figure 7

- **Figure 1.** Transverse section of the intestine of the control fish *Channa punctatus* showing the normal organization of intestine. Columnar Epithelium (CE), sub mucosa (SM) and serosa (S) Bouin, 6 μm, HE. X 400.
- **Figure 2.** Intestine of *C. punctatus* exposed for 15 days to sublethal concentration (SLC) 5 mg/L of Atrazine; Bouin, 6 μm, HE x400.
- **Figure 3.** Intestine of *C. punctatus* exposed for 15 days to sublethal concentration 10 mg/L of Atrazine; Bouin, 6 μm, HE x400.
- **Figure** 4. Intestine of *C. punctatus* exposed for 15 days to sublethal concentration 15 mg/L of Atrazine; Bouin, 6 μm, HE x400.
- **Figure** 5. Intestine of *C. punctatus* exposed for 30 days to sublethal concentration 5 mg/L of Atrazine; Bouin, 6 μm, HE x400.
- **Figure 6.** Intestine of *C. punctatus* exposed for 30 days to sublethal concentration 10 mg/L of Atrazine; Bouin, 6 μm, HE x400.
- **Figure 7.** Intestine of *C. punctatus* exposed for 30 days to sublethal concentration 15 mg/L of Atrazine; Bouin, 6 μm, HE x400.

CONCLUSION

The results obtained by the study on toxic effect of Atrazine clearly demonstrate that, at sublethal concentrations itself the herbicide has deleterious effect on the tissues exposed to it directly or even indirectly. Hence, Atrazine can be considered as a potent toxic pollutant capable of destroying the balance of aquatic ecosystem.

ACKNOWLEDGMENT

The authors express sincere thanks to the head of the Department of zoology, Govt. M. H. College of Home Science and Science for women for the facilities provided to carry out this research work.

REFERENCES

- Bhatnagar, A., & Srivastava, S.K. (1992). Aldose reductase: congenial and injurious profiles of an enigmatic enzyme. *Biochemical Medicine and Metabolic Biology*, 48(2), 91-121.
- Cui, H., Hwang, H.M., Zeng, K., Glover, H., Yu, H., & Liu, Y. (2002). Riboflavin-photosensitized degradation of atrazine in a freshwater environment. *Chemosphere*, 47(9), 991-999.
- Alvarez, D.C.M., & Fuiman, L. A. (2005). Environmental levels of atrazine and its degradation products impair survival skills and growth of red drum larvae. *Aquatic Toxicology*, 74(3), 229-241.
- Hussain, M., Nakahira, A., & Niihara, K. (1996). Mechanical property improvement of carbon fiber reinforced epoxy composites by Al2O3 filler dispersion. *materials letters*, 26(3), 185-191.
- Muley, D., Kamble, G., & Gaikwad, P. (1996). *Endosulfan toxicity in the freshwater fish Tilapia mossambica*. Paper presented at the Proceeding of Academic *Environmental Biolology*, 5(1), 49-55.
- Muniyan, M. (1999). Effect of ethofennprox (trebon) on the biochemical and histological changes in the selected tissues of the freshwater fish, *Oreochromis mossambicus* (Peters). Ph. D. Thesis, Annamalai University, India.
- Rahmathullah, L., Tielsch, J.M., Thulasiraj, R., Katz, J.,

- Coles, C., Devi, S., Edwin, N. (2003). Impact of supplementing newborn infants with vitamin A on early infant mortality: community based randomised trial in southern India. *Bmj*, *327*(7409), 254.
- Ravanaiah, G., & Murthy, C. (2010). Impact of aquaculture and industrial pollutants of Nellore district on the histopathological changes in the liver and intestine tissues of fish, Tilapia mossambica. *National Journal of Life Sciences*, 7(2), 117-122.
- Roberts, I., Schierhout, G., & Alderson, P. (1998). Absence of evidence for the effectiveness of five interventions routinely used in the intensive care management of severe head injury: a systematic review. *Journal of Neurology, Neurosurgery and Psychiatry*, 65(5), 729-733.
- Sharma, R., Pandey, A., & Shukla, G. (2001). Histopathological alterations in fish induced by pesticides toxicity: *Aquaculture*, (2),31-43.
- Steinberg, D. (1997). Low density lipoprotein oxidation and its pathobiological significance. *Journal of Biological Chemistry*, 272(34), 20963-20966.
- Velmurugan, B., Selvanayagam, M., Cengiz, E.I., & Unlu, E. (2007). Histopathology of lambda-cyhalothrin on tissues (gill, kidney, liver and intestine) of Cirrhinus mrigala. *Environmental Toxicology and Pharmacology*, 24(3), 286-291.
- Waring, C. P., & Moore, A. (2004). The effect of atrazine on Atlantic salmon (Salmo salar) smolts in fresh water and after sea water transfer. *Aquatic Toxicology*, 66(1), 93-104.
- Yildirim, M.Z., Benlı, A.Ç.K., Selvı, M., Özkul, A., Erkoç, F., & Koçak, O. (2006). Acute toxicity, behavioral changes, and histopathological effects of deltamethrin on tissues (gills, liver, brain, spleen, kidney, muscle, skin) of Nile tilapia (Oreochromis niloticus L.) fingerlings. *Environmental Toxicology: An International Journal*, 21(6), 614-620.
- Zhou, Q., Brown, J., Kanarek, A., Rajagopal, J., & Melton, D. A. (2008). In vivo reprogramming of adult pancreatic exocrine cells to β-cells. *Nature*, *455*(7213), 627.