



## ADULTICIDAL AND OVICIDAL ACTIVITIES OF *RHODOMYRTUS TOMENTOSA* LEAF EXTRACTS AGAINST DENGUE VECTOR *AEDES AEGYPTI*

Muthu Kasinathan, Jayapal Subramaniam,\* Chakravarthy Elanchezhian,  
Samivel Kanthammal and Mayakrishnan Vijay

<sup>1</sup>Department of Zoology, Annamalai University, Annamalai Nagar, Chidambaram-608002. Tamil Nadu, India.

**Article History:** Received 30<sup>th</sup> March 2018; Accepted 27<sup>th</sup> April 2018; Published 30<sup>th</sup> April 2018

### ABSTRACT

Mosquitoes has developed the resistance to synthetic insecticides, unfavorable effects to human and animal health, beneficial organisms and the environment, there is an urgent need to develop new insecticides of botanical based, which are suitable alternative bio control techniques with effective, safe, biodegradable and target-specific in the future. The adulticidal activities of crude hexane, acetone, and methanol extracts of the leaf of *Rhodomyrtus tomentosa* were assayed for their toxicity against dengue, chikungunya and zika virus vector, *Aedes aegypti*, the adult mortality was observed after 24 h of exposure. All extracts showed moderate adulticidal effects; however, the highest adult mortality observed was found in methanol extract. The LC<sub>50</sub> and LC<sub>90</sub> values of *R. tomentosa* leaf extracts against adulticidal activity of (acetone methanol and aqueous) *A. aegypti*, were the following: LC<sub>50</sub> values were 113.09 ppm for (RTAE), 106.38 ppm for (RTME) and 125.25 ppm for (RTAQE); LC<sub>90</sub> values were 251.18 for (RTAE), 229.94 for (RTME) ppm and 273.38 ppm for (RTAQE), respectively. The results of the ovicidal activity of acetone and methanol extracts of *R. tomentosa* plant at six different concentrations of 60-420 ppm were applied on eggs of *A. aegypti*. The toxicity of leaf extracts was dependent on its concentration against *A. aegypti*. There was zero hatchability (100% mortality) was attained at the concentration of 240 ppm for methanol extract and 300 ppm acetone extract and 360 ppm for aqueous extract. In this observation, this plant crude extracts gave the potential to be used as an ideal eco-friendly approach for the control of mosquitoes. This is the first report on mosquito adulticidal and ovicidal activities of the tribal medicinal plant, *R. tomentosa* against mosquito vectors from Southern India.

**Keywords:** Insecticide, Vector borne diseases, Eco friendly, Public health, Crude extracts.

### INTRODUCTION

Mosquito-borne diseases, such as malaria, Japanese encephalitis, filariasis, yellow fever, dengue and zika remain a major source of illness and death worldwide, particularly in tropical and subtropical countries (Becker *et al.*, 2003). Dengue is a mosquito-borne flavivirus found in tropical and sub-tropical regions of the world, mostly in urban and semi-urban regions. Dengue virus is transmitted by female mosquitoes mainly of the species *Aedes aegypti* and, to a lesser extent, *Ae. albopictus*. This mosquito's also can transmit chikungunya, yellow fever and Zika infection (WHO, 2015). *Aedes aegypti*, the primary carrier for viruses that cause dengue fever, dengue haemorrhagic fever, chikungunya fever, and yellow fever is widespread over large areas of the tropics and

subtropics (Kasinathan *et al.*, 2018; Service & Youdeowei, 1983; Yang *et al.*, 2004). It is the fastest spreading vector borne viral disease and is now endemic in over 100 countries, resulting in 40% of the world's population living in an area at risk for dengue. Dengue fever / dengue hemorrhagic fever (DHF) continues to be of major public health importance in countries of the Western Pacific and Southeast Asia. It is caused by one of four distinct serotypes (DEN -1, DEN - 2, DEN - 3 and DEN - 4) reported by WHO, (2016). The chemical based insecticides are play a very important role in controlling mosquito vector in their breeding sites, but this kind of control methods confirmed a negative impact in areas of beneficial and non target organisms and still mosquito vectors resistance to synthetic insecticides. In the view of the

\*Corresponding Author: Dr. C. Elanchezhian, Associate Professor, Department of Zoology, Annamalai University, Tamilnadu, India. Email: chezhian6@gmail.com, Mobile: +91 98658 25558

recently increased interest in developing plant origin insecticides as an alternative to synthetic insecticides (Amer & Mehlhorn, 2006; Wattanachai & Tintanon, 1999a). The environmental safety and low cost insecticide is considered to be of paramount importance and should not cause mortality on non-target organism in order to be acceptable (Kabaru & Gichia, 2001). However, the environmental threat that these chemicals pose, effects on non-target organisms, and the resistance of mosquitoes to insecticides have all increased during the last five decades (Amer & Mehlhorn, 2006; Wattanachai & Tintanon, 1999).

Biopesticides provide an alternative to synthetic pesticides because of their generally low environmental pollution, low toxicity to humans, and other advantages (Liu *et al.*, 2000). Many herbal products have been used as natural insecticides before the discovery of synthetic organic insecticides. Plants may be an alternative source of mosquito control agents because they constitute a rich source of bioactive chemicals (Kamaraj *et al.*, 2010; Tiwary *et al.*, 2007).

*Rhodomyrtus tomentosa* (Aiton) Hassk is a small shrub belonging to the Myrtaceae family and this herbal plants that are commonly used in traditional medicine in Southeast Asia (Winotai *et al.*, 2005). *R. tomentosa* have been recognized by several researchers. All parts of this plant (leaves, roots, buds, and fruits) have been often used in traditional medicine to treat colic diarrhoea (Ong & Nordiana, 1999), dysentery, abscesses, haemorrhage, and gynecopathy. In addition, it is used to formulate skin-whitening, antiaging, and skin beautifying agents (Miyake & Nojima, 2006). The extract from this plant possess strong inhibitory activity against gram-positive bacteria (Limsuwan *et al.*, 2009; Saising *et al.*, 2008), antimicrobial properties (Limsuwan & Voravuthikunchai, 2008; Sianglum *et al.*, 2011; Voravuthikunchai *et al.*, 2010) the Larvicidal and Pupicidal (Kasinathan *et al.*, 2018). The *R. tomentosa* plant part has been employed in traditional medicine to treat colic diarrhoea, haemorrhage, dysentery, abscesses, and gynaecopathy (Kanner *et al.*, 1994; Lavanya *et al.*, 2012; Sanchez moreno, 2002; Vinson & Hontz, 1995). No data regarding its adulticidal properties are available. Hence, the present investigation was aimed to explore the adulticidal and ovicidal activity of the extracts (ethanol, methanol and aqueous) from *Rhodomyrtus tomentosa* leaves against dengue and zika virus vector, *Aedes aegypti* under the laboratory condition.

## MATERIAL AND METHODS

### Collection of eggs Maintenance of larvae *Aedes aegypti*

The eggs of *Aedes aegypti* were collected from Vector Control Research Centre (VCRC - ICMR, Pondicherry), India without expose to any insecticide and in and around

Coimbatore district, India at different breeding habitats with the help of 'O' type brush. These eggs were brought to the laboratory and were transferred to 18 X 13 X 4 cm size enamel trays containing 500 ml of water and kept for larval hatching. The mosquito larval culture was maintained in our laboratory at  $27 \pm 2^{\circ}\text{C}$ , 75 - 85% RH, under 14L: 10D photoperiod cycles. The mosquito larvae were fed with dog biscuits and yeast at 3:1 ratio. The feeding was continued till the larvae were transformed into pupae.

### Maintenance of pupae and adult *Aedes aegypti*

The pupae were collected from the culture trays and were transferred to plastic containers (12 X 12 cm) containing 500 ml of water. The plastic jars were kept in 90 X 90 X 90 cm sized mosquito cage for adult emergence. The freshly emerged adults were maintained  $27 \pm 2^{\circ}\text{C}$ , 75 - 85% RH, less than 14L: 10D photoperiod cycles. The adults were fed with 10% sugar solution for a period of three days before they were provided an animal for blood feeding.

### Blood feeding of adult Mosquitoes

The female mosquitoes were allowed to feed on the blood of a rabbit (exposed on the dorsal side) for two days. The males were provided with 10% glucose solution on cotton wicks. The cotton was always kept moist with the solution and changed every day. An egg trap (cup) lined with filter paper containing water was placed at a corner of the cage egg collection.

### Collection of plant materials

The tribal medicinal plant, *Rhodomyrtus tomentosa* were collected from Nilgiri hills, Western Ghats, Tamil Nadu, India. The plants were identified (Taxonomy) at BSI (Botanical Survey of India, Coimbatore, Tamil Nadu, India.) and the specimens were deposited at Zoology Department, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu, India. This plant is a small tree growing up to 5 to 10 feet. It is an edible plant of higher altitude. It's used by the native tribal for stomach pain.

### Preparation of plant extracts

*R. tomentosa* plant was washed with tap water and shade dried at room temperature ( $27 \pm 2^{\circ}\text{C}$ ). An electrical blender powdered the dried plant materials (leaves). From the powder, 300 g of the plant materials was extracted with 1 L of organic solvents of Acetone and Methanol for using a Soxhlet apparatus boiling point range 60 - 80°C for 8 h. The extracts were filtered through a Buchner funnel with Whatman number 1 filter paper. The crude plant extracts were evaporated to dryness in rotary vacuum evaporator. One gram of the plant residue was dissolved in 100 ml of acetone (stock solution) and considered as 1% stock

solution. From this stock solution, different concentrations were prepared ranging from 120 to 600 ppm, respectively.

### Adulticidal bioassay

Sugar - fed adult female mosquitoes (5 to 6 days old) were used. The *R. tomentosa* leaf extract were diluted with acetone to make different concentrations. The diluted plant extracts were impregnated on filter papers (140 × 120 mm). A blank paper consisting of only ethanol was used as control. The papers were left to dry at room temperature to let the ethanol evaporate overnight. Impregnated papers were prepared fresh prior to testing. The bioassay was conducted in an experimental kit consisting of two cylindrical plastic tubes both measuring 125 × 44 mm following the method of (WHO, 1981). Mortality of the mosquitoes was recorded after 24 h. The above procedure was carried out in triplicate for plant extract of each concentration.

### Ovicidal Activity

For ovicidal activity, slightly modified method of (Su & Mulla, 1998) was performed. The egg raft / eggs of *A. aegypti* were collected from vector control Laboratory, Annamalai University. The *R. tomentosa* leaves extracts diluted in the appropriate solvent to achieve various concentrations ranging from 60 to 420 ppm. Eggs of these mosquito species (100 nos.) were exposed to each concentrations of leaf extract until they hatched or died. After treatment the eggs from each concentration were individually transferred to distilled water cups for hatching assessment after counting the eggs under microscope. Each experiment was replicated six times along with appropriate control. The hatch rates were assessed 48 h post treatment by following formula.

$$\% \text{ of egg mortality (Ovicidal activity)} = \frac{\text{Number of unhatched eggs}}{\text{Total number of eggs introduced}} \times 100$$

### Statistical analysis

The average adult mortality data were subjected to probit analysis for calculating LC<sub>50</sub>, LC<sub>90</sub>, and other statistics at 95 % fiducial limits of upper fiducial limit and lower fiducial limit, and Chi-square values were calculated using the SPSS Statistical software package 16.0 version was used. Results with  $P < 0.05$  were considered to be statistically significant.

## RESULTS AND DISCUSSION

The results of the adulticidal activity of acetone, methanol and aqueous extracts of *R. tomentosa* against the adult of dengue and chikungunya vector *A. aegypti*, are Tables 1 presented in (Figure 1 and 2). Among the three vectors tested, the highest adulticidal activity was observed in the highest mortality followed by methanol, acetone and aqueous extract. At higher concentrations, the adult showed restless movement for some time with abnormal wagging and died. The rates of mortality were directly proportional to concentration. The LC<sub>50</sub> and LC<sub>90</sub> values of *R. tomentosa*

leaf extracts against adulticidal activity of (acetone, methanol and aqueous) *A. aegypti*, were the following: Acetone extract LC<sub>50</sub> and LC<sub>90</sub> values were 113.09 and 251.18 ppm; Methanol extract LC<sub>50</sub> (LC<sub>90</sub>) values were 106.38 (229.94) ppm; Aqueous extract LC<sub>50</sub> values were 125.25 ppm and LC<sub>90</sub> values were 273.38 ppm, respectively. Similarly, highest adulticidal effect was established from *Piper sarmentosum*, followed by *Piper ribesoides*, and *Piper longum*, with LD<sub>50</sub> values of 0.14, 0.15, and 0.26 µg / female, respectively. Govindarajan *et al.* (2013) have reported that the moderate adulticidal properties were found in methanol extract of *A. paniculata* against the adults of *C. quinquefasciatus* and *A. aegypti* with the LC<sub>50</sub> and LC<sub>90</sub> values were 149.81 and 172.37 ppm, and 288.12 and 321.01 ppm, respectively. Plant borne extracts and essential oils can be tested for their adulticidal properties against a number of mosquitoes of economic importance (Amerasan *et al.* (2012); Govindarajan *et al.*, 2013; Panneerselvam & Murugan, 2013). For example, Subramaniam *et al.* (2015) reported that the adulticidal activity of methanol extracts of seaweeds *D. dichotoma*, *P. pavonica* and *V. pachynema* on the costal malarial vector *Anopheles sundaicus*, LC<sub>50</sub> values were 147.18 ppm, 161.9 pm and 133.79 ppm, respectively. In particular, 88 % adult mortality was observed from *Pelargonium citrosa* leaf extracts at 2 % concentration against *A. stephensi* (Jeyabalan *et al.*, 2003).

The ethanol extract of *Citrus sinensis* showed LC<sub>50</sub> and LC<sub>90</sub> values 320.38 and 524.57 ppm against *A. aegypti* adults (Murugan *et al.*, 2012). Recently, (Subramaniam *et al.*, 2016) have reported that the adulticidal properties of *C. guianensis* flower extract and nanoparticles showed LC<sub>50</sub> and LC<sub>90</sub> of 133.96 and 11.23 ppm and 287.65 and 24.61 ppm, respectively. For instance, (Subramaniam *et al.*, 2015) were highlighted the adulticidal properties of *Mimusops elengi* extract, which are highly toxic against *A. stephensi* and *A. albopictus*, respectively. The percentage of egg hatchability of dengue and zika vector, *A. aegypti* was tested with three different solvents (acetone, methanol and aqueous) at various concentrations of *R. tomentosa* leaf extracts and the results are listed in (Table 2). The toxicity of leaf extracts was dependent on its concentration against *A. aegypti*. There was zero hatchability (100% mortality) was attained at the concentration of 240 ppm for methanol extract and 300 ppm acetone extract and 360 ppm for aqueous extract. At 360 ppm all the three extracts exerted the zero hatchability. Control eggs exerted the hatchability rate ranged from 100%. The hatchability rate of control group with acetone, methanol and aqueous were 97.6%, 96.2% and 98.2%, respectively. The ovicidal activity of *R. tomentosa* against *A. aegypti* revealed that the hatchability rate of control egg rafts was ranged from 96.2% to 100%. The bio-active compound *Azadiracta indica* showed highest ovicidal activity in the eggs of *Culex tarsalis* and *Culex quinquefasciatus* exposed to 10 ppm concentration (Su & Mulla, 1998). Plant extracts have been screened and studied for their ovicidal repellency activity against mosquitoes (Tennyson *et al.*, 2011; Thavara, *et al.*, 2002).

For examples, some researchers have been reported the essential oil and plant extracts, which have highest ovicidal potential against major mosquito vectors, *A. stephensi*, *A. aegypti*, and *C. quinquefasciatus* (Prajapati *et al.*, 2005), *Cinnamomum zeylanicum*, *Zingiber officinale* and *Rosmarinus officinalis* oils (Miyake & Nojima, 2006) *Citrullus vulgaris* (Samidurai *et al.*, 2009). Recently, Amerasan *et al.* (2015) *Morinda tinctoria* and *Pongamia*

*glabra*, Subramaniam & Murugan (2013) *Myristica fragrans* seed and Subramaniam *et al.* (2017) *Chenopodium ambrosioides* and Kasinathan *et al.* (2018) have reported the larvicidal and pupicidal activity of methanolic leaf extract of *R. tomentosa* (RTME) at different concentrations (120 to 600 ppm). The LC<sub>50</sub> values were 263.82, 299.43, 331.64, 386.16 and 419.56 ppm (I - IV instar and pupae), respectively.

**Table 1.** Adulticidal toxicity of *R. tomentosa* leaf extracts against the Dengue and Zika virus vector, *Ae. aegypti*.

| Treatment                               | Dosage (ppm) | Mortality (%)            | LC <sub>50</sub> (LCL-UCL)  | LC <sub>90</sub> (LCL-UCL)  | $\chi^2$  |
|-----------------------------------------|--------------|--------------------------|-----------------------------|-----------------------------|-----------|
| <i>R. tomentosa</i> -acetone extract    | Control      | 0.0 ± 0.0                | 113.09<br>(97.85 - 126.27)  | 251.18<br>(229.10 - 282.37) | 3.29 n.s. |
|                                         | 50           | 31.6 ± 2.0 <sup>e</sup>  |                             |                             |           |
|                                         | 100          | 43.8 ± 1.8 <sup>d</sup>  |                             |                             |           |
|                                         | 150          | 59.0 ± 2.0 <sup>bc</sup> |                             |                             |           |
|                                         | 200          | 76.8 ± 2.2 <sup>ab</sup> |                             |                             |           |
|                                         | 250          | 93.4 ± 2.8 <sup>a</sup>  |                             |                             |           |
| <i>R. tomentosa</i> -methanolic extract | Control      | 0.0 ± 0.0                | 106.38<br>(54.09 - 139.27)  | 229.94<br>(187.06 - 342.47) | 8.60 n.s. |
|                                         | 50           | 33.2 ± 2.6 <sup>e</sup>  |                             |                             |           |
|                                         | 100          | 45.8 ± 2.2 <sup>d</sup>  |                             |                             |           |
|                                         | 150          | 61.0 ± 2.6 <sup>bc</sup> |                             |                             |           |
|                                         | 200          | 79.8 ± 2.0 <sup>ab</sup> |                             |                             |           |
|                                         | 250          | 98.4 ± 1.8 <sup>a</sup>  |                             |                             |           |
| <i>R. tomentosa</i> -aqueous extract    | Control      | 0.0 ± 0.0                | 125.25<br>(110.04 - 138.91) | 273.38<br>(247.92 - 310.24) | 3.92 n.s. |
|                                         | 50           | 29.4 ± 1.2 <sup>e</sup>  |                             |                             |           |
|                                         | 100          | 39.8 ± 1.8 <sup>cd</sup> |                             |                             |           |
|                                         | 150          | 55.4 ± 2.6 <sup>c</sup>  |                             |                             |           |
|                                         | 200          | 69.8 ± 2.2 <sup>b</sup>  |                             |                             |           |
|                                         | 250          | 90.6 ± 1.6 <sup>a</sup>  |                             |                             |           |


Mortality rates are means ± SD of five replicates. No mortality was observed in the control. Within each column means followed by the same letter(s) are not significantly different (P<0.05). Chi-square value followed by an asterisk is significant (heterogeneity factor used in calculation of confidence limits) (P<0.05).

LC<sub>50</sub> lethal concentration that kills 50 % of the exposed organisms, LC<sub>90</sub> lethal concentration that kills 90% of the exposed organisms, LCL lower confidence limit, UCL upper confidence limit,  $\chi^2$  Chi-square test, NS not significant.


**Table 2.** Ovicidal activity of *R. tomentosa* leaf extracts against the eggs of Dengue and Zika virus vector, *Ae. aegypti*

| Treatment                               | Total | Control                  | Egg hatchability (%)     |                          |                          |                          |                        |     |     |
|-----------------------------------------|-------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|------------------------|-----|-----|
|                                         |       |                          | Concentration (ppm)      |                          |                          |                          |                        |     |     |
|                                         |       |                          | 60                       | 120                      | 180                      | 240                      | 300                    | 360 | 420 |
| <i>R. tomentosa</i> -acetone extract    | 100   | 97.6 ± 2.05 <sup>a</sup> | 67.5 ± 1.80 <sup>b</sup> | 48.9 ± 1.50 <sup>c</sup> | 25.7 ± 1.55 <sup>d</sup> | 11.2 ± 1.24 <sup>e</sup> | NH                     | NH  | NH  |
| <i>R. tomentosa</i> -methanolic extract | 100   | 96.2 ± 1.82 <sup>a</sup> | 62.3 ± 1.56 <sup>c</sup> | 44.5 ± 1.32 <sup>d</sup> | 22.2 ± 1.78 <sup>e</sup> | NH                       | NH                     | NH  | NH  |
| <i>R. tomentosa</i> -aqueous extract    | 100   | 98.2 ± 1.82 <sup>a</sup> | 69.3 ± 1.16 <sup>b</sup> | 54.5 ± 1.72 <sup>c</sup> | 36.2 ± 1.18 <sup>d</sup> | 20.4 ± 0.82 <sup>e</sup> | 08 ± 1.82 <sup>f</sup> | NH  | NH  |

Values in a row with a different superscript are significantly different at p < 0.05% level (DMRT test). Each value (X ± SD) represents the mean of six values. NH - No hatchability (100% mortality).



**Figure 1.** Adulticidal toxicity of *R. tomentosa* leaf extracts against the Dengue and Zika virus vector, *Ae. aegypti*.



**Figure 2.** Adulticidal activity of *R. tomentosa* leaf extracts against the eggs of Dengue and Zika virus vector, *Ae. Aegypti*.

## CONCLUSION

The conclude the results showed that the adulticidal and ovidal activities of acetone methanol and aqueous extract of *R. tomentosa* leaves has potential to be developed as an insecticide against dengue, zika and chikungunya vector, *A. aegypti*. However, further studies to evaluate its toxicity and effects on non-target organisms and the environment need to be conducted. These are environmentally safe and eco-friendly approaches for the vector control programs. The results of the present study could be useful in promoting research aimed at the development of new agents for mosquito control based on bioactive chemical compounds from indigenous plant sources.

## ACKNOWLEDGEMENT

We thank Dr. R. Karuppasamy, Professor and Head, Dr. A. Jebanesan, Professor, Department of Zoology,

Annamalai University for the laboratory facilities provided. The authors are grateful to Dr. K. Gunasekaran, Scientist 'G' and Mr. T. Vijay Kumar, Senior Technician, Vector Control Research Centre, Puducherry for their help and suggestions provided for the present work.

## REFERENCES

- Amer, A., & Mehlhorn, H. (2006). Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitology Research*, 99(4), 466 - 472.
- Amerasan, D., Murugan, K., Kovendan, K., Kumar, P. M., Panneerselvam, C., Subramaniam, J., Hwang, J. S. (2012). Adulticidal and repellent properties of *Cassia tora* Linn. (Family: Caesalpinaceae) against *Culex quinquefasciatus*, *Aedes aegypti*, and *Anopheles stephensi*. *Parasitology Research*, 111(5), 1953-1964.

- Amerasan, D., Murugan, K., Panneerselvam, C., Kanagaraju, N., Kovendan, K., & Kumar, P. M. (2015). Bioefficacy of *Morinda tinctoria* and *Pongamia glabra* plant extracts against the malaria vector *Anopheles stephensi* (Diptera: Culicidae). *Journal of Entomological and Acarological Research*, 47(1), 31-40.
- Becker, N., Petric, D., Zgomba, M., Boase, C., Dahl, C., & Lane, J. (2003). *Mosquitoes and their control*. New York, Boston, Dordrecht, London, Moscow: Kulmer Academic: Plenum Press.
- Govindarajan, M., Sivakumar, R., Rajeswary, M., & Yogalakshmi, K. (2013). Chemical composition and larvicidal activity of essential oil from *Ocimum basilicum* (L.) against *Culex tritaeniorhynchus*, *Aedes albopictus* and *Anopheles subpictus* (Diptera: Culicidae). *Experimental parasitology*, 134(1), 7-11.
- Jeyabalan, D., Arul, N., & Thangamathi, P. (2003). Studies on effects of *Pelargonium citrosa* leaf extracts on malarial vector, *Anopheles stephensi* Liston. *Bioresource technology*, 89(2), 185-189.
- Kabaru, J., & Gichia, L. (2001). Insecticidal activity of extracts derived from different parts of the mangrove tree *Rhizophora mucronata* (Rhizophoraceae) Lam. against three arthropods. *African Journal of Science and Technology*, 2(2), 66 -73.
- Kamaraj, C., Rahuman, A. A., Bagavan, A., Zahir, A. A., Elango, G., Kandan, P., Santhoshkumar, T. (2010). Larvicidal efficacy of medicinal plant extracts against *Anopheles stephensi* and *Culex quinquefasciatus* (Diptera: Culicidae). *Tropical Biomedicine*, 27(2), 211-219.
- Kanner, J., Frankel, E., Granit, R., German, B., & Kinsella, J. E. (1994). Natural antioxidants in grapes and wines. *Journal of Agricultural and Food Chemistry*, 42(1), 64-69.
- Kasinathan, M., Subramaniam, J., Elanchezhian, C., Kanthammal, S., & Vijay, M. (2018). Mosquitocidal potential of *Rhodomyrtus tomentosa* leaf extracts against Dengue and Zika virus vector, *Aedes aegypti*. *International Journal of Entomology Research*, 3(1), 95-100.
- Lavanya, G., Voravuthikunchai, S. P., & Towatana, N. H. (2012). Acetone extract from *Rhodomyrtus tomentosa*: a potent natural antioxidant. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-8.
- Limsuwan, S., Trip, E. N., Kouwen, T. R., Piersma, S., Hiranrat, A., Mahabusarakam, W., Kayser, O. (2009). Rhodomyrtone: a new candidate as natural antibacterial drug from *Rhodomyrtus tomentosa*. *Phytomedicine*, 16(6 -7), 645-651.
- Limsuwan, S., & Voravuthikunchai, S. P. (2008). *Boesenbergia pandurata* (Roxb.) Schltr., Eleutherine americana Merr. and *Rhodomyrtus tomentosa* (Aiton) Hassk. as antibiofilm producing and antiquorum sensing in *Streptococcus pyogenes*. *FEMS Immunology & Medical Microbiology*, 53(3), 429-436.
- Liu, S., Shi, J., Cao, H., Jia, F., Liu, X., & Shi, G. (2000). Survey of pesticidal component in plant. *Entomology in China in 21st Century*, 1098-1104.
- Miyake, Y., & Nojima, J. (2006). Skin Cosmetic and Food/Drink for Cosmetological Use. *Maruzen Pharmaceutical, Hiroshima*, Japan Press.
- Murugan, K., Kumar, P. M., Kovendan, K., Amerasan, D., Subramaniam, J., & Hwang, J.S. (2012). Larvicidal, pupicidal, repellent and adulticidal activity of *Citrus sinensis* orange peel extract against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). *Parasitology Research*, 111(4), 1757-1769.
- Ong, H., & Nordiana, M. (1999). Malay ethno-medico botany in Machang, Kelantan, Malaysia. *Fitoterapia*, 70(5), 502-513.
- Panneerselvam, C., & Murugan, K. (2013). Adulticidal, repellent, and ovicidal properties of indigenous plant extracts against the malarial vector, *Anopheles stephensi* (Diptera: Culicidae). *Parasitology Research*, 112(2), 679-692.
- Prajapati, V., Tripathi, A., Aggarwal, K., & Khanuja, S. (2005). Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. *Bioresource technology*, 96(16), 1749-1757.
- Saising, J., Hiranrat, A., Mahabusarakam, W., Ongsakul, M., & Voravuthikunchai, S. P. (2008). Rhodomyrtone from *Rhodomyrtus tomentosa* (Aiton) Hassk. as a natural antibiotic for staphylococcal cutaneous infections. *Journal of Health Science*, 54(5), 589-595.
- Samidurai, K., Jebanesan, A., Saravanakumar, A., Govindarajan, M., & Pushpanathan, T. (2009). Larvicidal, ovicidal and repellent activities of *Pemphis acidula* Forst.(Lythraceae) against filarial and dengue vector mosquitoes. *Academic Journal of Entomology*, 2(2), 62-66.
- Sanchez Moreno, C. (2002). Methods used to evaluate the free radical scavenging activity in foods and biological systems. *Food science and technology international*, 8(3), 121-137.
- Service, M. W., & Youdeowei, A. (1983). *Pest and vector management in the tropics*: Longman Scientific & Technical Press.
- Sianglum, W., Srimanote, P., Wonglumsom, W., Kittiniyom, K., & Voravuthikunchai, S. P. (2011). Proteome analyses of cellular proteins in methicillin-resistant *Staphylococcus aureus* treated with rhodomyrtone, a novel antibiotic candidate. *PLoS One*, 6(2), e16628.

- Su, T., & Mulla, M. (1998). Ovicidal activity of neem products (azadirachtin) against *Culex tarsalis* and *Culex quinquefasciatus* (Diptera: Culicidae). *Journal of the American Mosquito Control Association*, 14(2), 204-209.
- Subramaniam, J., & Murugan, K. (2013). Evaluation of larvicidal, pupicidal, repellent, and adulticidal activity of *Myristica fragrans* against malarial vector *Anopheles stephensi*. Paper presented at the National Conference on Insect Diversity and Systematics, *Aligarh Muslim University Press*, 28-30.
- Subramaniam, J., Murugan, K., Jebanesan, A., Pontheckan, P., Dinesh, D., Nicoletti, M., Canale, A. (2017). Do *Chenopodium ambrosioides*-synthesized silver nanoparticles impact *Oryzias melastigma* predation against *Aedes albopictus* Larvae?. *Journal of Cluster Science*, 28(1), 413-436.
- Subramaniam, J., Murugan, K., Panneerselvam, C., Kovendan, K., Madhiyazhagan, P., Dinesh, D., Rajaganesh, R. (2016). Multipurpose effectiveness of *Couroupita guianensis*-synthesized gold nanoparticles: high antiparasitic potential, field efficacy against malaria vectors and synergy with *Aplocheilus lineatus* predators. *Environmental Science and Pollution Research*, 23(8), 7543-7558.
- Subramaniam, J., Murugan, K., Panneerselvam, C., Kovendan, K., Madhiyazhagan, P., Kumar, P. M., Nicoletti, M. (2015). Eco-friendly control of malaria and arbovirus vectors using the mosquitofish *Gambusia affinis* and ultra-low dosages of *Mimusops elengi*-synthesized silver nanoparticles: towards an integrative approach? *Environmental Science and Pollution Research*, 22(24), 20067-20083.
- Tennyson, S., Ravindran, K. J., & Arivoli, S. (2011). Screening of plant extracts for ovicidal activity against *Culex quinquefasciatus* Say (Diptera: Culicidae). *Elixir Appl Botany*, 40, 5456-5460.
- Thavara, U., Tawatsin, A., & Chompoosri, J. (2002). *Phytochemicals as repellents against mosquitoes in Thailand*. Paper presented at the Proceedings International Conference on Biopesticide. University Press, 2-7.
- Tiwary, M., Naik, S., Tewary, D. K., Mittal, P., & Yadav, S. (2007). Chemical composition and larvicidal activities of the essential oil of *Zanthoxylum armatum* DC (Rutaceae) against three mosquito vectors. *Journal of vector borne diseases*, 44(3), 198-201.
- Vinson, J. A., & Hontz, B. A. (1995). Phenol antioxidant index: comparative antioxidant effectiveness of red and white wines. *Journal of Agricultural and Food Chemistry*, 43(2), 401-403.
- Voravuthikunchai, S. P., Dolah, S., & Charernjitrakul, W. (2010). Control of *Bacillus cereus* in foods by *Rhodomyrtus tomentosa* (Ait.) Hassk. Leaf extract and its purified compound. *Journal of food protection*, 73(10), 1907-1912.
- Wattanachai, P., & Tintanon, B. (1999). Resistance of *Aedes aegypti* to chemical compounds in aerosol insecticide products in different areas of Bangkok, Thailand. *Communicable Diseases Journal*, 25(2), 188-191.
- WHO. (1981). Instructions for determining the susceptibility or resistance of adult mosquitoes to organochlorine, organophosphorous and carbamate insecticides. *Establishment of the base-line*. Geneva: World Health Organization Press.
- WHO. (2015). *Investing to Overcome the Global Impact of Neglected Tropical Diseases: Third WHO Report on neglected Tropical Diseases, 2015*, (3), World Health Organization Press.
- WHO. (2016). *WHO treatment guidelines for drug-resistant tuberculosis 2016 update*: World Health Organization Press.
- Winotai, A., Wright, T., & Goolsby, J.A. (2005). Herbivores in Thailand on *Rhodomyrtus tomentosa* (Myrtaceae), an invasive weed in Florida. *Florida Entomologist*, 88(1), 104-105.
- Yang, Y.C., Park, I.K., Kim, E.H., Lee, H.S., & Ahn, Y.J. (2004). Larvicidal activity of medicinal plant extracts against *Aedes aegypti*, *Ochlerotatus togoi*, and *Culex pipiens pallens* (Diptera: Culicidae). *Journal of Asia-Pacific Entomology*, 7(2), 227-232.