International Journal of Zoology and Applied Biosciences Volume 3, Issue 2, pp: 345-351, 2018 UGC Approved Journal No. 46511



**Research Article** 

# ANTIOXIDANT PROPERTIES OF PORTUNUS SANGUINOLENTUS CHITOSAN FABRICATES CRAB SHELL

<sup>1</sup>Sakthidasan, V., \*<sup>1</sup>Pugazhendy, K., <sup>1</sup>Tamizhazhagan, V. and <sup>2</sup>Jayanthi

<sup>1</sup>Department of Zoology, Annamalai University, Annamalai Nagar, Chidambaram-608 002, Tamilnadu, India <sup>2</sup>Department of Education, Annamalai University, Annamalai Nagar, Chidambaram-608 002, Tamilnadu, India

Article History: Received 22<sup>nd</sup> February 2018; Accepted 28<sup>th</sup> April 2018; Published 30<sup>th</sup> April 2018

## **ABSTRACT**

The chitosan got from the chitin is the second most bottomless polysaccharide delivered after cellulose. It is the fundamental auxiliary segment of the exoskeletons of creatures like shellfish and is found in the cell dividers of growths moreover. Chitosan is delivered primarily from the crab shell squander. The present work is gone for the extraction of chitin from three spotted crab shell *Portunus sanguinolentus* and transformation of chitin into chitosan. The methanolic concentrate of the crab shell of *P. sanguinolentus* was subjected to standard subjective examination and the in-vitro cancer prevention agent movement was assessed by the assurance of aggregate cell reinforcement limit, 1.1-diphenyl-2 picrylhydrazyl (DPPH) radical searching action, Antioxidant action, and Nitric oxide potential. The examinations uncovered that the methanol concentrate of Co could productively rummage the free radicals in a dosage-dependent way. The outcomes were contrasted and the standard cancer prevention agent ascorbic corrosive. Along these lines, additional research might be justified to think about dynamic mixes of *P. sanguinolentus* that give the cancer prevention agent movement. The discoveries displayed here might have suggestions in the populace illness avoidance field.

Keywords: Portunus sanguinolentus, DPPH, Total antioxidant, Nitric oxide scavenge, Methanol.

## INTRODUCTION

Chitin is one of the most to a great extent accessible inexhaustible modest biopolymer acquired from marine birthplace (Synowiecki & Al Khateeb, 1997). It is biocompatible, biodegradable (Amano & Ito, 1978) and bio-absorbable, with antibacterial and wound mending properties and furthermore with negligible immunogenicity. In this manner, there were a few examinations being directed to uncover its biomedical utilization of chitin. Fittingly, an extremely broad determination of projects in different territories, for example, sustenance building, item examine, microbiology, horticulture, wastewater treatment, tissue designing, bionanotechnology and sedate conveyance have just been accounted for late, chitin and its subsidiaries have just been dissected as presumably of utilization pharmaceutical and were regularly found to indicate negligible harmfulness (van der Lubben et al., 2001). Pharmacology and therapeutics to develop various drugs synthesis build new remedies for various diseases (Tamizhazhagan & Pugazhendy, 2017). Asia has abundant species of medicinal and aromatic plants and traditional medicines have practiced in Asia since ancient times. India has made use of medicinal plants to cure ailments of thousands of years (Tamizhazhagan *et al.*, 2017a). Quality can be defined as the status of a drug that is determined by identity, purity, content and another chemical, physical, or biological properties, or by the manufacturing processes (Tamizhazhagan *et al.*, 2017b). Antioxidant enzymes make up the first line of protection alongside oxidative constant worry and smash up caused by free radicals (Tamizhazhagan & Pugazhendy, 2017).

The crabs rank third after shrimps and lobsters for their regarded fish delicacy and furthermore the estimation of fishery they bolster. The crab fishery in India is quick creating and there is a huge degree for the crab meat because of its delicacy and wholesome extravagance. The crab meat contains a rich measure of protein, vitamins A and D, minerals, glycogen and free amino acids. The

greater part of the marine crabs happening along the Indian coasts is having a place with the family portunidae. The monetarily vital portunid crabs found along Parangipettai drift are Scylla serrata, S. tranquebarica, Portunus sanguinolentus, P. pelagicus, Podophthalamus vigil, C. feriata, C. lucifera, C. natator, C. granulata and C. truncata (John Samuel et al., 2004; Varadharajan et al., 2009).

happening Normally cancer prevention substances additionally require security testing. Alerts with respect to a presumption of security of common cancer prevention agents have been more than once prompted since the reality cell reinforcement originates from a characteristic source does not demonstrate its accepted wellbeing (Hattori et al., 1998). Oxidative pressure is associated with the pathology of growth, arteriosclerosis, jungle fever and rheumatoid joint inflammation and could assume a part in neurodegenerative infection and maturing forms (Aruoma, 1997). The increase in free amino acid level suggests that tissues damage probably due to the increased proteolytic activity under toxic stress (Pichaimani et al., 2017). The antioxidant GSH is tripeptide-γ-glutamylcystenineglycine or reduced glutathione. In living organisms GSH plays an important role in normal cell function (Jayalakshmi et al., 2017).

The part of cancer prevention agents has gotten expanded consideration amid the previous decade. Be that as it may, the utilization of such cancer prevention agents has potential wellbeing risk (Park et al., 2004). In this way, as of late, interests have been produced for seeking powerful normal cancer prevention agents, since they can shield human body from free radical and retard the advance of numerous endless sicknesses. The hemolymph, in many scavengers, is a dry liquid containing an expansive number of cells or hemocytes and different natural and inorganic constituents. Around 90% of hemolymph is water, which decides its aggregate volume. The hemolymph fills in as a washing medium for different tissues and organs as they do not have an epithelial coating of a genuine coelom. Accordingly, the hemolymph shapes the gathering spot of both the crude materials required and the results of different physiological exercises of the body. Since the hemolymph isn't specifically associated with the outside condition any adjustment in it can be taken as a measure of the physiological condition of the inward condition of the in-place creature. The haemolymph of shellfish have intense antimicrobial peptide indicated different cluster against a few human pathogens (Arul Prakash et al., 2011) and assumes a part in have resistance reaction including non-self-acknowledgment, self cell correspondence, superoxide anion action, melanisation, phagocytosis, cytotoxicity and exemplification (Sahoo et al., 2005).

Current scenario regular items from marine examples have a wide range of natural exercises and various restorative applications incorporate antiviral, antibacterial, and antitumor. Cyclic and direct peptides found from marine creatures have expanded our insight about new intense cytotoxic, antimicrobial, particle channels particular blockers, and numerous different properties with novel compound structures related to unique instruments of pharmacological movement (Zheng et al., 2011). There is an expanding enthusiasm for cancer prevention agents, especially in those of free radicals in different maladies. These obsessive and clinical foundations have provoked to research novel and intense cancer prevention agent peptides from crab which are at last of restorative utilize. Henceforth the present examination was meant to distinguish the cancer prevention agent action of hemolymph from delicate and hard-shelled crabs of *Charybdis luciferase*. Biomaterials are those non-living materials utilized as a part of the restorative, biomedical and different fields, meaning to communicate with the organic framework (Wang & Chio, 1998).

A large number of these materials, for example, chitosan, are utilized as successful choices for the substitution of tissues, including bone tissue since they don't present dangers of illness transmission or invulnerable dismissal, and also boundless accessibility and minimal effort. A standout amongst the most vital fields of utilizations of common mixes it is prescription. Such materials would have a few favorable circumstances overengineered ones. Materials got from nature, have been appeared to yield speedier recuperating with less contradiction in individuals.

Chitosan is framed by deacetylation of chitin. Chitin is a normally happening polysaccharide found in the exoskeleton of shellfish, for example, crabs (Blumenthal & Roseman, 1957). Chitosan advances cross-linkage with different cross-connecting operators, for example, glutaraldehyde, sodium tripolyphosphate (TPP), genipin and so forth. In the past investigation, we utilized copper as another metal particle and complexed it with chitosan. Cu (II) particles could be effectively adsorbed on the chitosan surface at pH 6. As of late, the superb antibacterial property of copper against gram +ve and ve microscopic organisms and cytotoxicity potential against numerous disease cells have been accounted for (Koleva et al., 2002). Despite the fact that copper has been appeared to have anticancer movement in certain disease cells, there is no examination on its ecacy in osteosarcoma. Chitin and its subordinates including D-glucosamine-carboxymethyl chitin Dihydroxypropyl chitin have safe tweaking (immunomodulatory) effects and can influence intrinsic and greedy resistance which prompts increment cell movement and the discharge of cytokines and chemokines (Wanasundara & Shahidi, 1998).

Chitosan additionally happens normally in a few microorganisms, for example, parasites and yeast and is viewed as the biggest biomaterial after cellulose as far as usage and dissemination. Its concoction name is 2-amino-2-deoxy-b-D-glucopyranose and its recipe is  $(C_6H_{11}O_4N)$ . The advancement of business applications for chitosan has advanced. Chitin and chitosan are delivered monetarily in India, Poland, Japan, Thailand, USA, Norway and Australia (Ghorbel *et al.*, 2011). The present study was carried isolated compound chitosan enhanced fabricated have many oxidative functions were the focus.

### MATERIALS AND METHODS

## **Preparation of extract**

The *Portunus sanguinolentus* were collected from the fish market at Chidambaram (Lat. 11°.39'N and Long. 79° 69'E) near Vandigate. The skeleton was placed in Ziploc cover and frozen overnight and then subsequently cut into smaller pieces using a meat tenderizer. 100 g of crab shells were ground to get coarse powder and cleaned with water and 0.9% saline by blending the shell powder in a 500 ml measuring utensil with water for a couple of minutes. After this, the shell powders are separated off. This procedure is rehashed to the point when sand and different soils are evacuated. The cleaned crab shell (Figure 1) powders are dried overnight in the drying broiler at 80°C.



Figure 1. Portunus sanguinolentus crab shell.

# **Total Antioxidant Activity**

The total antioxidant activity of the fractions was determined according to the method of (Prieto et~al., 1999). 0.3mL of each fraction was mixed with 3.0 mL of reagent solution (0.6M sulphuric acid, 28mM sodium phosphate, and 4Mm ammonium molybdate). The reaction mixture was incubated at 95°C for 90 min in a water bath. The absorbance of all the sample mixtures was measured at 695 nm. Ascorbic acid (100  $\mu$ g/mL) was used as a standard control.

# **Determination of the Total Phenolics**

The total phenolic content of fractions (bioactive compounds) was determined spectrophotometrically with Folin-Ciocalteu reagent, using a slightly modified method by Junaid et al., 2006). The extract of the fraction was mixed with Folin-Ciocalteu reagent (1: 1) and 4mL of sodium carbonate (1M) was added and allowed to stand for absorbance minutes. The spectrophotometrically at 765 nm. A standard curve was plotted using different concentrations of gallic acid (standard,  $0-1000 \mu g/mL$ ) and the total phenolic content of ethenol extract was estimated as ug gallic acid equivalents (GAE)/mg of extract. The reaction was conducted in triplicate, and the results were averaged.

### **Total Flavonoid content**

The Flavonoid content was estimated using the spectrophotometric method (Junaid *et al.*, 2006; Viarengo *et al.*, 1997). The total flavonoid content was estimated as  $\mu g$  quercetin equivalents/mg of extract. Ethyl acetate extracts of *Portunus sanguinolentus* (250  $\mu$ l) was mixed with 1.25 mL of distilled water and 75  $\mu$ l of 5 % NaNO<sub>2</sub> solution. After 5 min, 150  $\mu$ l of 10 % AlCl<sub>3</sub> was added. After 6 min, 500  $\mu$ l of 1M NaOH and 275  $\mu$ l of distilled water were added to prepare the mixture. The solution was mixed well & the absorbance was read at 510 nm using UV-VIS spectrophotometer.

# **DPPH Radical Scavenging Activity**

The free radical scavenging activity of the metabolite, based on the scavenging activity of the stable 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical was determined according to the method of (Kumaran, 2006) and the absorbance was measured at 517 nm spectrophotometrically (Elico SL 159, India). The lower absorbance of the reaction mixture indicates higher free radical scavenging activity. The percent of DPPH radical scavenging activity was calculated as,

Inhibition (%) =  $(A - A0) / A \times 100$ .

Where,

A is the absorbance of the control.

A0 is the absorbance of the metabolite/standard.

## Nitric Oxide Radical Scavenging Activity

Sodium nitroprusside in aqueous solution at physiological pH spontaneously generates nitric oxide, which interacts with oxygen to produce nitrite ions, which can be determined using the Griess Illosvoy reaction. 2ml of 10mM sodium nitroprusside in 0.5ml of phosphate buffer saline (pH 7.4) was mixed with 0.5ml of extract and standard (curcumin) at various concentrations and the mixture was incubated at 25°C for 150min. From the incubated mixture 0.5ml was taken out and added into 1ml of sulfanilic acid reagent and incubated for 5min. Finally, 1ml naphthyl ethylenediamine dihydrochloride (0.1% w/v) was mixed and incubated at room temperature for 30 min before measuring absorbance at 540 nm using a spectrophotometer. Nitric oxide radical scavenging activity was calculated using the formula,

Inhibition (%) =  $(Ao - Ae) / Ao \times 100$ .

Where,

Ao is the absorbance of the control.

Ae is the absorbance of the extract/standard.

# RESULTS

Total Phenol and Flavonoid content were estimated using gallic acid and quercetin as standard references. The Stigmasterol Trimethylsilyl Ether produced  $0.321 \pm 0.001$ 

mg gallic acid equivalents as phenol content and  $0.014\pm0.0002$  mg quercetin equivalents as phenol and flavonoid content (Table 1).

**Table 1.** Total phenol and flavonoid content of Stigmasterol Trimethylsilyl Ether.

	Total flavonoid	Total phenol	
Compound	content mg	content mg	
Name	quercetin	Gallic acid	
	equivalent	equivalent	
Stigmasterol	$0.014 \pm 0.002$	$0.321 \pm 0.001$	
Trimethylsilyl			
Ether			

The value represents Mean  $\pm$  SD.

Free radicals are having a lone pair of electron containing groups which produced harmful effects to the living beings by causing oxidation mechanism which results in the lethal effect. The antioxidants are the agent which prevent oxidation process being produced by this kind of free radical such as OH, O, NO, and Fe<sup>+</sup>.

In total antioxidant activity (Figure 2), the Stigmasterol Trimethylsilyl Ether has produced dose dependent manner increasing activity in the range of 65.853  $\pm$  0.001, 72.097  $\pm$  0.001 and 78.341  $\pm$  0.001 respectively (Table 2).

Stigmasterol Trimethylsilyl Ether exhibited a significant dose-dependent inhibition of DPPH activity. The potential of L-ascorbic acid to scavenge DPPH radical is directly proportional to the concentration. The DPPH assay activity is near to standard as ascorbic acid is presented in (Figure 3). Stigmasterol Trimethylsilyl Ether moderately inhibited nitric oxide in dose-dependent manner (Figure 4), with ascorbic acid.

Table 2. The total antioxidant activity of Stigmasterol Trimethylsilyl Ether

A cook list	Sample name	Total Anti-Oxidant Activity		
Assay list	Sample name	100µg/ml	200µg/ml	300µg/ml
Total antioxidant	Stigmasterol Trimethylsilyl Ether	$65.853 \pm 0.001$	$72.097 \pm 0.001$	$78.341 \pm 0.001$
	Vit – C	$75.099 \pm 1.0$	$82.621 \pm 1.0$	$87.203 \pm 0.57$
DPPH Assay	Stigmasterol Trimethylsilyl Ether	$60.53 \pm 0.001$	$59.85 \pm 0.001$	$58.57 \pm 0.001$
	Vit – C	$59.92 \pm 1.0$	$32.69 \pm 1.0$	$37.75 \pm 0.57$
Nitric Oxide Radical Scavenging	Stigmasterol Trimethylsilyl Ether	$79.34 \pm 0.001$	$82.03 \pm 0.001$	$83.19 \pm 0.001$
	Vit – C	$23.09 \pm 1.0$	$40.21 \pm 1.0$	$56.03 \pm 0.57$

The value represents Mean  $\pm$  SD.





Figure 2. The Total antioxidant activity of Stigmasterol Trimethylsilyl Ether.



**Figure 3.** DPPH Scavenging activity (Each value represents Mean  $\pm$  SD, n=3).




Figure 4. Nitric Oxide Radical Scavenging Activity (RSA).

## DISCUSSION

Antioxidants due to their scavenging activity are useful for the management of those diseases. DPPH stable free radical method is a sensitive way to determine the antioxidant activity of plant extracts (Koleva et al., 2002; Kumar et al., 2008). The therapeutic potential of natural medicinal plants as an antioxidant in reducing such free radical-induced tissue injury, suggests that many plants have antioxidant activities that can be therapeutically useful (Kanatt et al., 2007). India has made use of medicinal plants to cure ailments of thousands of years (Tamizhazhagan & Pugazhendy, 2017). Excellence defined as the status of a drug that is strongly minded by identity, purity, content and another chemical, physical, or biological properties, or by the developed processes (Tamizhazhagan et al., 2017a). Antioxidant enzymes make up the first line of protection alongside oxidative steady worry and shatter up caused by free radicals (Tamizhazhagan et al., 2017a), compared to another source of laboratory animals

like fish's vegetation and vertebrate animals expand pharmacology and therapeutics various drugs synthesis build new remedies for various (Tamizhazhagan et al., 2017b). These differences might be due to their different antioxidant mechanisms or variations in their ability to scavenge free radicals. A fair correlation between total phenolic content and antioxidant activity was also observed. These observations clearly indicated a crosslinkage between phenolics and antioxidant activity. However, a large number of bio active compound groups are implicated for antioxidant activity (Tamizhazhagan et al., 2017a). Ascorbic acid acts as a chain-breaking antioxidant impairs the formation of free radicals in the process of formation of intracellular substances throughout the body, including collagen, bone matrix and tooth dentine. These differences might be due to their different antioxidant mechanisms or variations in their ability to scavenge free radicals. A fair correlation between total phenolic content and antioxidant activity was also observed.

### **CONCLUSION**

These observations clearly indicated a cross-linkage between phenolics and antioxidant activity. However, a large number of bioactive compound groups are implicated for antioxidant activity. The results based on the related to other finding more efficacies have the chitosan fabricated freshwater crab of *Portunus sanguinolentus*.

### ACKNOWLEDGEMENT

The authors are gratefully acknowledged the facility provides Department of Zoology, Dr. R. Karuppasamy, Annamalai University, Tamilnadu, India

## REFERENCES

- Amano, K.I., & Ito, E. (1978). The Action of Lysozyme on Partially Deacetylated Chitin. *The Febs Journal*, 85(1), 97-104.
- Arul Prakash, A., Balasubramanian, S., Gunasekaran, G., Prakash, M., & Senthil Raja, P. (2011). Control of clinical pathogens by the haemolymph of *Paratelphusa hydrodromous*, a freshwater crab. *ISRN Pharmacology*, 1-4. doi:10.5402/2011/642768.
- Aruoma, O.I. (1997). Scavenging of hypochlorous acid by carvedilol and ebselen in vitro. *General Pharmacology*, 28(2), 269-272.
- Blumenthal, H.J., & Roseman, S. (1957). Quantitative Estimation of Chitin In Fungi. *Journal Of Bacteriology*, 74(2), 222.
- Ghorbel Bellaaj, O., Hmidet, N., Jellouli, K., Younes, I., Maâlej, H., Hachicha, R., & Nasri, M. (2011). Shrimp waste fermentation with *Pseudomonas aeruginosa* A2: optimization of chitin extraction conditions through Plackett–Burman and response surface methodology approaches. *International Journal of Biological Macromolecules*, 48(4), 596-602.
- Hattori, M., Yamaji Tsukamoto, K., Kumagai, H., Feng, Y., & Takahashi, K. (1998). Ant oxidative activity of soluble elastin peptides. *Journal of Agricultural and Food Chemistry*, 46(6), 2167-2170.
- Jayalakshmi, S., Pugazhendy, K., Tamizhazhagan, V., Sakthidasan, V., Jayanthi, C., & Sasikala, P. (2017). Therapeutic efficacy of Aloe vera against the antioxidant enzyme effect of cypermethrin in the fresh water fish Cyprinus carpio. International Journal of Current Advanced Research, 6(12), 8054-8060
- John Samuel, N., Thirunavukkarasu, N., Soundarapandian, P., Shanmugam, A., & Kannupandi, T. (2004). Fishery potential of commercially important portunid crabs along Parangipettai coast. *Proceedings of Ocean Life Food and Medicine Expo*, 165-173.

- Junaid, S., Olabode, A., Onwuliri, F., Okwori, A., & Agina, S. (2006). The antimicrobial properties of ocimum gratissimum extracts on some selected bacterial gastrointestinal isolates. *African Journal of Biotechnology*, 5(22), 2315-2321.
- Kanatt, S. R., Chander, R., & Sharma, A. (2007). Antioxidant potential of mint (mentha spicata l.) in radiation-processed lamb meat. *Food Chemistry*, 100(2), 451-458.
- Koleva, I.I., Van Beek, T.A., Linssen, J.P., Groot, A.D., & Evstatieva, L. N. (2002). Screening of plant extracts for antioxidant activity: a comparative study on three testing methods. *Phytochemical Analysis*, 13(1), 8-17.
- Kumar, P.S., Sucheta, S., Deepa, V.S., Selvamani, P., & Latha, S. (2008). Antioxidant activity in some selected indian medicinal plants. *African Journal Of Biotechnology*, 7(12).
- Kumaran, A. (2006). Antioxidant And Free Radical Scavenging activity of an aqueous extract of coleus aromaticus. *Food Chemistry*, 97(1), 109-114.
- Park, P.J., Je, J.Y., & Kim, S.K. (2004). Free Radical Scavenging activities of differently deacetylated chitosans using an esr spectrometer. *Carbohydrate Polymers*, 55(1), 17-22.
- Pichaimani, N., Pugazhendy, K., Tamizhazhagan, V., Sakthidasan, V., Jayanthi, C., & Sasikala, P. (2017). Antioxidant enzyme activity effect of *Solanam Virginianum* against lead acetate toxicity of the fresh water fish *Cyprinus Carpio. International Journal of Current Advanced Research*, 6(12), 8031-8037.
- Prieto, P., Pineda, M., & Aguilar, M. (1999).

  Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Analytical Biochemistry*, 269(2), 337-341.
- Sahoo, B., Sethi, S., Mishra, B.K., & Das, B. (2005). Effects elecitors on prophenonoloxidase and superoxide anion activities of freshwater prawn, *Macrobrachium Malcolmsoni*i. *Asian Fisheries Science*, 18, 345-353.
- Synowiecki, J., & Al Khateeb, N.A.A.Q. (1997). Mycelia of *Mucor rouxii* as a source of chitin and chitosan. *Food Chemistry*, 60(4), 605-610.
- Tamizhazhagan, V., & Pugazhendy, K. (2017).
  Ethnobotanical and phytopharmacological review of Pisonia alba span. Asian Journal of Pharmaceutical and Clinical Research, 10(5), 69-71.
- Tamizhazhagan, V., Pugazhendy, K., Sakthidasan, V., & Jayanthi, C. (2017a). Antioxidant properties of

- Pisonia alba plant leaf extract. International Journal of Zoology and Applied Biosciences, 2(6), 311-314.
- Tamizhazhagan, V., Pugazhendy, K., Sakthidasan, V., & Jayanthi, C. (2017b). Preliminary screening of phytochemical evaluation selected plant of *Pisonia alba*. *International Journal of Biology Research*, 2(4), 63-66.
- Van Der Lubben, I.M., Verhoef, J.C., Borchard, G., & Junginger, H.E. (2001). Chitosan and its derivatives in mucosal drug and vaccine delivery. *European Journal of Pharmaceutical Sciences*, 14(3), 201-207.
- Varadharajan, D., Soundarapandian, P., Dinakaran, G., & Vijakumar, G. (2009). Crab fishery resources from Arukkattuthurai to Aiyammpattinam, south east coast of India. *Current Research Journal of Biological Sciences*, 1(3), 118-122.

- Viarengo, A., Ponzano, E., Dondero, F., & Fabbri, R. (1997). A simple spectrophotometric method for metallothionein evaluation in marine organisms: An application to mediterranean and antarctic molluscs. *Marine Environmental Research*, 44(1), 69-84.
- Wanasundara, U. N., & Shahidi, F. (1998). Antioxidant and pro-oxidant activity of green tea extracts in marine oils. *Food Chemistry*, 63(3), 335-342.
- Wang, S.L., & Chio, S.H. (1998). Deproteinization of shrimp and crab shell with the protease of *Pseudomonas aeruginosa* K-187. *Enzyme and Microbial Technology*, 22(7), 629-633.
- Zheng, L.H., Wang, Y.J., Sheng, J., Wang, F., Zheng, Y., Lin, X.K., & Sun, M. (2011). Antitumor peptides from marine organisms. *Marine Drugs*, 9(10), 1840-1859.