International Journal of Zoology and Applied Biosciences Volume 3, Issue 4, pp: 340-343, 2018 https://doi.org/10.5281/zenodo.1340654

Research Article

FUNGAL INFECTION IN SHRIMPS COLLECTED FROM AGNIAR ESTUARY, TAMIL NADU, INDIA

R. Saraswathi and P. Sumithra*

P.G. & Research Department of Microbiology, Srimad Andavan Arts and Science College, Tiruchirappalli-620 005, Tamil Nadu, India

Article History: Received 15th December 2017; Accepted 27th April 2018; Published 9th August 2018

ABSTRACT

Today shell fishes diseases are of great concern to aqua culturists as they inflict heavy losses by causing large scale mortalities resulting in huge economic losses. Among the various pathogens, fungi come next to bacteria. As knowledge of diseases is basic to their control, the present study was intended to find out the commonly occurring fungi in cultured shrimp farms in Tamil Nadu. Results show that 18 species of fungi could be identified in *Penaeus monodon* and *P. indicus* of which three species *P. monodon*, four species *P. indicus*. However, even though *P. vannamei* also recorded 18 fungal species only two were recorded throughout the period of culture. Nevertheless, in all the three species of *Penaeus*, the maximum number of fungal species (18) was uniformly recorded on the 80th day of culture.

Keywords: Shrimp culture, Estuary, Shell fishes diseases, Fungal infection.

INTRODUCTION

Wetlands although occupying only 4-6% of earth's land area are considered among the most important natural resources of the earth as they are the sources of cultural, economic and biological diversity (Mitra & Majumdar, 2005). Biodegradation of organic matter in this habitat is supported by a wide variety of organisms like bacteria and fungi which are widely distributed in water and sediments of these habitats (Nambiar *et al.*, 2008).

Fungi are important components of the ecosystem typically constituting more of biomass than bacteria and have been considered the largest biotic community after insects (Sarbhoy *et al.*, 1996). Fungi play an indispensable role in industry, agriculture, medicine and biogeochemical cycles (Cowan, 2001; Gates *et al.*, 2005). In addition, fungi are also used in fish industry, textile, bioremediation, biofertilizers, biotechnology, *etc.* (Acharya & Pradhan, 2010; Hunt, 1999; Manoharachary *et al.*, 2005; Pilz & Molina, 2002; Pilz *et al.*, 2006).

Today, the problems of diseases are of great concern to aquaculturists as they inflict heavy losses by causing large scale mortalities resulting in a drastic decrease in yield. Among the various pathogens, fungi come next only to bacteria. As India is poised for a large scale development of aquaculture especially of prawns with a stress on semi

intensive and intensive culture systems, it is natural to expect increasing hazards of diseases in these systems. However, diseases in fish are not understood well. What is known about fish diseases often relates to aquaria fish and more recently to farm fish (Sharma *et al.*, 2012). As knowledge of diseases, their causative agents and etiology is basic to control, the present study was intended to find out the commonly occurring fungal pathogens that occurs in three species of prawns commonly cultured in farms using the same estuarine water.

MATERIAL AND METHODS

Sample Collection

Penaied shrimp (21-30g) sample was collected during August 2017 to July 2018 by trawl net from estuarine waters of Agniar Estuary, Pudukkottai district, Tamil Nadu, India. For each collection, -20 number of fungal brown-gill infected *Penaeus* sp. were collected for further processing.

Isolation and identification of fungi

Penaeus sp. Infected gills were washed three times in sterile physiological saline (0.85% NaCl) and inoculated on plates with Potato Dextrose Agar (PDA) medium (Himedialab). Ampicillin and streptomycin sulphate

(25μg/ml) were added to the medium to inhibit bacterial growth. Plates were incubated for 24 days at 25°C and then subcultured on to fresh PDA plates. The singles poreculture method was applied to obtain pure isolates. The isolates were maintained at 25°C on PDA slants for subsequent experiments. Fungal identification was done by lactophenol cotton blue mount method to observe fungal morphology under light microscopy.

Histopathology method

Routine histological techniques were used. Davidson's fixative was injected into the hepato pancreas and muscle of live fungal fouling infected shrimp in order to avoid autolysis of those organs. After injection, the whole shrimp was immersed in the same fixative for 48h before further processing. The infected portion of brown-gill tissue were dissected and immersed in Davidson's fixative for 48h and transferred to 70% ethanol for further processing. Sections were stained with Ehrlich's hematoxylin and Eosin stains.

RESULTS AND DISCUSSION

The results of the various fungal infections in three species of prawns that were cultured are prevented in Tables 1 and 2. As evident from the table, a total of 18 species of fungi could be identified during the 120-day culture period in the three species of prawns. As far as *Penaeus monodon* was concerned, 18 species of fungi were identified belonging to six genera (*Mucor, Rhizopus, Aspergillus, Fusarium, Penicillium* and *Trichoderma*. Of these, three species were found to occur throughout the period of investigation (*Rhizopus nigricans, Aspergillus niger* and *Fusarium solani*). *Rhizopus nigricans* recorded their overgrowth between the 60th and 80th days, while *Fusarium solani* recorded an overgrowth in the 80th day and *A. niger* was uniformly recorded throughout the period of study.

Among the genus Rhizopus, R. nigricans was noticed throughout and recorded an overgrowth on the 60th and 80th days; R. oryzae which was absent only on the 10th day also recorded an overgrowth on the 60th day and 80th days and R. stolonifer which was recorded from the 80th to 120th day recorded an overgrowth on the 80th day. Thus it appears that they preferred the 80th day to record their highest counts. Within the genus Aspergillus, only A. niger was recorded throughout the period of culture. A perusal of the Table-1 reveals four species of Aspergillus (A. flavus, A. fumigatus, A. terreus and A. vesicolor) recorded overgrowths on the 80th day of culture. However, A. japonicus which was recorded from the 80th day, showed a uniform growth till the 120th day. Among the genus Fusarium, none of the species were recorded throughout. However, all the three species that were observed recorded an overgrowth on the 80th day of culture. Within the genus Penicillium also, none of the species were recorded throughout the culture period. While P. chrysogenum and P. griseofulvum were not recorded between the 60th and 80th days of culture, *P. oxalicum* was not observed during the 10th and 120th day of culture. However, it recorded an overgrowth during the 60th and 80th day of culture.

Among the two species that were recorded in the genus Trichoderma, none of them were recorded throughout. However, both the species recorded an overgrowth on the 80th and 100th days of culture. Thus, an overall comparison reveals that on the 20th day of culture, eight species of fungi were recorded in P. monodon. This increased to 12 species on the 40th day and almost remained stable till the 60th day (11 species). However, the number increased to 16 and 18 species on the 80th and 100th day only to decrease again on the 120th day (9 species). With regard to fungal infection in P. indicus, a total of 18 species belonging to the same six genera that were noticed in P. monodon were recorded. Among the 18 species, only four species (R. nigricans, A. flavus, P. chrysogenum and T. viride) were recorded throughout the period of culture. The genus Mucor was represented by a single species M. racemosus which was not recorded during the 20th and 120th day of culture. On other days, it showed uniform growth.

The genus Rhizopus was represented by three species of which only one (R. nigricaus) was recorded throughout. It recorded an overgrowth on the 80th day of culture. The other two species (R. oryzae and R. stolonifer) when present recorded overgrowth on the 80th day. The genus Aspergillus was recorded by six species of which only one (A. flavus) was recorded throughout. It recorded an increased growth on the 80th day of culture. The remaining species of Aspergillus, eventhough were not recorded throughout the period of culture also recorded their overgrowth on the 80th day when present. Among the three species of Fusarium that were recorded, none were recorded throughout the period of culture. However, here also, all the three species when present, recorded overgrowth on the 80th day of culture. Within the genus Penicillium, among the three species that were recorded, only one (P. chrysogenum) occurred throughout and recorded an overgrowth on the 80th day. The other two species, when present, also recorded their overgrowth on the 80th day of culture. Among the two species of the genus Trichoderma that were observed, T. viride was recorded throughout showing an uniform growth throughout the days of culture. The other species (T. harzianum) recorded overgrowth on the 80th day of culture.

Thus, an overall comparison reveals that there was a gradual increase in the number of fungal infections as the days progressed till about the 80th day of culture (18 species) followed by a decline till the 120th day (6 species). The fungal infections that were recorded on *P. vannamei* are presented in Tables 1 which reveals again that a total of 18 species belonging to 6 genera that were observed in other prawns. However, among these, only two species were recorded throughout the period of culture (*A. fumigatus* and *A. niger*). The genus *Mucor* was represented by a single species *M. racemosus* which was recorded only from the 60th till the 120th day of culture. At that time, it is recorded an average growth.

Table 1. Fungal Infection on *Penaeus* species collected from Agniar Estuary.

S. No.	Species	Penaeus monodon							Penaeus indicus					Penaeus vannamei					
		20	40	60	80	100	120	20	40	60	80	100	120	20	40	60	80	100	120
1.	Mucor racemosus	+	-	-	+	+	-	-	+	+	+	+	-	-	-	+	+	+	+
2.	Rhizopus nigricans	+	+	++	++	+	+	+	+	+	++	+	+	-	-	-	-	-	-
3.	Rhizopus oryzae	-	+	++	++	+	+	+	+	+	++	+	-	+	+	+	+	-	-
4.	Rhizopus stolonifer	-	+	-	++	+	+	-	+	+	++	+	-	+	+	+	++	-	-
5.	Aspergillus flavus	+	-	-	++	+	+	+	+	+	++	+	+	-	-	+	++	+	+
6.	Aspergillus fumigatus	-	+	-	++	+	+	-	-	+	++	+	-	+	+	+	++	+	+
7.	Aspergillus japonicus	-	-	-	+	+	+	+	+	+	++	-	-	+	-	++	+	+	+
8.	Aspergillus niger	+	+	+	+	+	+	-	+	+	++	+	+	+	+	+	++	+	+
9.	Aspergillus terreus	+	+	+	++	+	-	+	+	+	++	+	-	+	+	++	+	-	-
10.	Aspergillus versicolor	-	+	++	++	+	-	-	+	+	++	+	-	-	+	++	+	+	-
11.	Fusarium quaeductum	-	-	+	++	+	-	-	-	+	++	+	-	+	-	-	+	+	-
12.	Fusarium oxysporum	-	-	+	++	+	-	+	+	+	++	-	-	-	-	-	+	+	+
13.	Fusarium solani	+	+	+	++	+	-	-	-	++	++	+	+	+	-	+	+	-	-
14.	Penicillium	+	+	-	-	+	+	+	+	+	++	+	+	+	+	+	++	-	-
	chrysogenum																		
15.	Penicillium	+	+	-	-	+	+	-	-	-	+	-	-	-	-	-	-	-	-
	griseofulvum																		
	Penicillium oxalicum	-	+	++	++	+	-	-	-	+	-	+	-	+	+	+	++	+	-
17.		-	+	+	++	++	-	-	-	-	++	-	-	+	+	+	++	+	-
10	harzianum																		
18.	Trichoderma viride	-	-	+	++	++	-	+	+	+	+	+	+	-	-	+	+	-	

⁻ Represents no growth, + Represents average growth, ++ Represents over growth.

The genus *Rhizopus* was represented by two species none of which were recorded throughout the period of culture. Both the species (*R. oryzae* and *R. stolonifer*) were recorded from the 20th till the 80th day of culture; while *R. stolonifer* recorded an overgrowth on the 80th day; *R. oryzae* recorded a uniformly average growth throughout. Among the six species of genus *Aspergillus* that were noticed, two were recorded throughout the period of culture (*A. fumigatus* and *A. niger*). Further, both the species recorded an overgrowth on the 80th day of culture; while *A. flavus* which was recorded from the 60th to the 120th day of culture also recorded an overgrowth on the 80th day. The other species (*A. japonicus*, *A. terreus* and *A. versicolor*) recorded an overgrowth on the 60th day of culture.

The genus *Fusarium* was represented by three species, none of which were recorded throughout the period of culture. Further, when present, all the three species showed an average growth. The genus *Penicillium* was also represented by two species both of which were not recorded throughout the period of culture. However, when present, both the species (*P. chrysogenum* and *P. oxalicum*) recorded an overgrowth on the 80th day. The genus *Trichoderma* was also represented by two species of which

none occurred throughout the period of culture. While T. viride was recorded only on two occasions, T. harzianum was recorded between the 20^{th} and 100^{th} day of culture recording an overgrowth on the 80^{th} day.

Thus an overall comparison reveals that as the days progressed the number of fungi infecting prawns also increased culminating in a peak on the 80th day recording 16 infections followed by a decreasing trend till the 120th day (6 species). A perusal of literature reveals that (Velmurugan & Ayyaru, 2014) while studying the fungal diversity in three species of cultured prawns reported the presence of 20 fungal species that belonged to six genera. This study is comparable to the present investigation in which 18 species of fungi belonging to 6 genera could be identified.

In the present study, in *Penaeus monodon*, three species were perennial (*Rhizopus nigncans, Aspergillus niger*and *Fusarium solani*) while in *P. indicus*, four species were perennial (*R. nigricans, A. flavus, P. chrysogenum* and *Trichoderma viridae*) and in *P. vannameri*, two species were perennial (*A. fumigatus* and *A. niger*). Thus, in all the three species of prawn, *Aspergillus* was found to dominate.

A similar dominance of Aspergillus in prawn was also reported by (Velmurugan & Ayyaru, 2014). Karunasagar et al. (2004) however, suggested that even though about 500 fungal species have been isolated from marine and estuarine conditions, only a few were found to be pathogenic to prawns. In the present study, Fusarium species could be isolated in all the three species of prawns. Reports reveal that Fusariosis and black gill disease are caused by Fusarium and can affect all the development stages of prawn (Ramaiah, 2006). He also suggested that Fusarium sp. is usually opportunistic pathogens which however can cause mortalities. The present investigation also revealed the presence of Penicillium sp. (Pillai & Vijayaraghayan, 1986) in his observations on various prawn diseases, reported that deep mycosis is produced by Penicillium in P. monodon. Khoa et al. (2004) isolated Fusarium sp. in P. monodon grown in various farms in Vietnam which brought about blackgill disease resulting in high mortality.

CONCLUSION

In the present study, all the three species of prawn recorded maximum growth of fungi at around the 80th day of culture. A similar observation was also reported by Velmurugan & Ayyaru, (2014). This might probably be attributed to the differences in the water exchange that is usually done in various farms. Nevertheless, the incidence of various fungi in all the three species of prawns analysed reveals the need to incorporate mycological examination to be an integral part of monitoring the health of prawns to prevent economic losses as well as prevention of diseases to handlers and human consumers.

ACKNOWLEDGMENT

The authors express sincere thanks to the Management and Principal, P.G. & Research Department of Microbiology, Srimad Andavan Arts and Science College for the facilities provided to carry out this research work.

REFERENCES

- Acharya, K., Rai, M., & Pradhan, P. (2010). Agaricales of Sikkim Himalaya: a review. *Researcher*, 2(5), 29-38.
- Cowan, A. (2001). Fungi-life support for ecosystems. *Essential ARB*, 4, 1-5.
- Gates, G., Ratkowsky, D., & Grove, S. (2005). A comparison of macrofungi in young silvicultural regeneration and mature forest at the Warra LTER Site in the southern forests of Tasmania. *Tasforests Hobart*, 16, 127.
- Hunt, G. (1999). Assessing macrofungi of special concern for conservation in forested ecosystems.

- Proceedings of a Conference on the Biology and Management of Species and Hbitats at Risk, 779-780.
- Karunasagar, I., Karunasagar, I., & Umesha, R. (2004). Microbial diseases in shrimp aquaculture. National Institute of Oceanography, Goa,121-123.
- Khoa, L., Hatai, K., & Aoki, T. (2004). *Fusarium incarnatum* isolated from black tiger shrimp, Penaeus monodon Fabricius, with black gill disease cultured in Vietnam. *Journal of Fish Diseases*, 27(9), 507-515.
- Manoharachary, C., Sridhar, K., Singh, R., Adholeya, A., Suryanarayanan, T., Rawat, S., & Johri, B. (2005). Fungal biodiversity: distribution, conservation and prospecting of fungi from India. *Current Science*, 89, 58-71.
- Mitra, S., & Majumdar, D. (2005). Effect of soil properties, substrate addition, microbial inoculation, and sonication on methane production from three rice topsoils and subsoils in the Philippines. Copyright *International Rice Research Institute Journal*, 546-549.
- Nambiar, G.R., Raveendran, K., Changxing, Z., & Jaleel, C.A. (2008). A glimpse of lignicolous marine fungi occurring in coastal water bodies of Tamil Nadu (India). *Comptes Rendus Biologies*, 331(6), 475-480.
- Pillai, C.T., & Vijayaraghavan, G. (1986). Tail rot disease and cardiac arrest in the culturable cichlid fish, etroplus suratensis. *Current Science*, 55(10), 512-513.
- Pilz, D., & Molina, R. (2002). Commercial harvests of edible mushrooms from the forests of the Pacific Northwest United States: issues, management, and monitoring for sustainability. *Forest Ecology and Management*, 155(1-3), 3-16.
- Pilz, D., Molina, R., & Mayo, J. (2006). Effects of thinning young forests on chanterelle mushroom production. *Journal of Forestry*, 104(1), 9-14.
- Ramaiah, N. (2006). A review on fungal diseases of algae, marine fishes, shrimps and corals. *Indian Journal of Marine Science*, 35(4), 380-387.
- Sarbhoy, A. K., Varshney, J., & Agarwal, D. (1996). *Fungi of India*, (Vol. 7): CBS Publishers & Distributors, 1982-92.
- Sharma, M., Shrivastav, A., Sahni, Y., & Pandey, G. (2012). Overviews of the treatment and control of common fish diseases. *International Journal Research Journal of Pharmacy*, 3(7),123-127.
- Velmurugan, K., & Ayyaru, G. (2014). Original Article Culturable fungal diversity of brown-gill disease in three *Penaeus* species. *International Journal of Reserach Marine Science*, 3, 1-4.