International Journal of Zoology and Applied Biosciences Volume 4, Issue 2, pp: 95-105, 2019 https://doi.org/10.5281/zenodo.2667879

Research Article

EFFECT OF HUMIC ACID, SEAWEED EXTRACT AND ESSENTIAL OILS AS ANTIOXIDANTS ON PRE- AND POST-HARVEST OUALITY OF RED RADISH PLANTS

M.T. Sakr^{1*}, Heba Ibrahim¹, Aml ElAwady² and Amira AboELMakarm²

¹Agricultural of Botany Department, Faculty of Agriculture, Mansoura University, Mansoura, Egypt-35516

²Post harvest and handling vegetables Research Department, Horticulture Research Intuition,

Agricultural Research Center, Giza, Egypt-35516

Article History: Received 14th April 2019; Accepted 26th April 2019; Published 30th April 2019

ABSTRACT

This study was carried out to assess the effect of certain biomodulators, namely seaweed extract, humic acid, peppermint and thyme oils, on quality-related metabolites during both pre- and post-harvest of Red radish. For this purpose, plants were sprayed with tested biomodulators twice, 20 and 30 days after sowing, and chosen biochemical constituents affecting quality were estimated before harvest, and again in roots of harvested plants after storage in standard cold storage. In addition, to evaluate the involvement of each tested parameter in post-harvest quality of stored roots, a correlation matrix between biochemical constituents after storage and quality parameters of stored roots represented by weight loss and post-harvest decay percentage as well as stored root's dry matter percentage. Results indicated that the adopted biomodulators affected the metabolism of treated plants in a way that persisted after storage, hence positively affected roots post-storage quality attributes. Invariably, stored roots of biomodulators-treated plants contained higher concentrations from total anthocyanin, ascorbic acid, total phenols, nutrients concentration as well as flavonoids compared with those of control plants. Moreover, activity of both peroxidase and polyphenol oxidase was higher in stored roots of biomodulators-treated plants. In addition, all applied biomodulators decreased weight loss and post-harvest decay percentage, thus roots of biomodulators-treated plants contained higher dry matter compared with roots of untreated plants. The metabolic and physiological bases of biomodulators-induced quality parameters of roots after storage were discussed.

Keywords: Seaweed extract, Humic acid, Peppermint oil, Raphanus sativus, Post harvest quality.

INTRODUCTION

The *Raphanus sativus* L. is a root vegetable widely grown for its nutritional, culinary and medicinal uses that belongs to Brassicaceae. It grows in temperate climates at altitudes between 190 and 1240 m. Its roots are thick and of various sizes, forms, and colors. Red Radish belongs to the variety radicular. The edible fleshy axis is derived from the hypocotyl and upper radicle tissues. Radish is widely used in salad preparations and contains a considerable amount of antioxidants, vitamin C, and health-promoting compounds such as glucosinolates and phenolic compounds. Various parts of the radish plant, including roots, seeds, and leaves, have been used for medicinal purposes. Its health-promoting properties have been attributed to polyphenolic

compounds. *R. sativus* is popularly used to treat liver and respiratory illnesses (Paredes, 1984), reduce cancer development (Ku, Lee, & Kim, 2008) and contains a range of digestive enzymes (Cho *et al.*, 2009). A comprehensive review of the plants' active constituents and its therapeutic properties could be found in the publication of (Gutiérrez & Perez, 2004). Radish has also been used in naturopathic medicine as a laxative, stimulant, and digestive aid, as well as in the treatment of stomach disorders (Kapoor, 2000). Red radish cultivars are a potential source of natural colorants due to the presence of anthocyanins, which have high stability. Anthocyanins have well-known health benefits, including the ability to scavenge free radicals, inhibit cancer and diabetes, prevent neuronal and

cardiovascular diseases, and suppress inflammation (Hwang *et al.*, 2012). Great attention is being given to post-harvest quality of vegetables due to rising consumer awareness about diet-health connections. Quality parameters of stored vegetables are determined by both pre-and postharvest factors (Kader, 2003). So, it is assumed that manipulation of growth and metabolism of plants during preharvest stage may have significant impact on postharvest quality of their stored products.

Humic acid (HA) are the most significant constituents of organic matter in soils and have a relevant role in the cycling of many elements in the environment and in soil ecological functions (Senesi *et al.*, 1996). Foliar sprays of HA promote growth, increase yield and quality in a number of plant species (Karakurt *et al.*, 2009). Moreover, humic acid influence respiration process, the amount of sugars, amino acids and nitrate metabolism (Boehme *et al.*, 2005). (Nardi *et al.*, 2002) characterized the effects of HA on physiology of higher plants. According to them, HA positively influencing the uptake of some nutrients, especially nitrate, may influence both respiration and photosynthesis, may display a hormone-like activity, and exhibit stimulatory effects on plant cell growth and development.

Seaweed products exhibit growth-stimulating activities, and the use of seaweed formulations as bio modulators in crop production is well established. Seaweed components such as macro- and microelement nutrients, amino acids, vitamins, cytokinins, auxins, and abscisic acid (ABA)-like growth substances affect cellular metabolism in treated plants leading to enhanced growth and crop yield (Ördög et al., 2004). In addition, it has been reported that seaweed extract enhanced the plant's antioxidant defense system of wheat plants subjected to drought stress through elevating both the enzymatic and non-enzymatic components of the plant's antioxidant defense system (Kasim et al., 2015).

Essential oils are well known for their antioxidant and antimicrobial properties that prevent food degradation and alteration (Justesen & Knuthsen, 2001). They possess antimicrobial properties and prevent growth of bacteria, fungus and virus during storage and protect cells from senescence, thereby increase shelf life of vegetables. Thyme oil contains more than 40% phenolic constituents (thymol, carvacrol and caffic acid) which have strong antispectic effects. Essential oils chemical composition of Thymus vulgaris and Mentha piperita were identified by GC-MS where the most components of thyme were thymol (44.7%), cymenin (18.6%) and tripeneine (16.5%) whereas menthol and menthone are the main components of peppermint oil (Porte & Godoy, 2008). Application of 4 % thyme and basil oils had a positive effect on broccoli seed germination, which was ascribed to their antioxidant and disinfecting effects (Nguefack et al., 2005).

The objectives of the current investigation was to assess the effect of bio stimulators seaweed extract (SWE),

humic acid (HA), thyme oil as well as peppermint oil on certain quality-related biochemical constituents during both pre- and post-harvest stages.

MATERIALS AND METHODS

Two field experiments were conducted during the two consecutive growing seasons of 2015 and 2016 at EL-Bramoon Farm, Mansoura city, Dakahlia Governorate. Seeds of *Raphanus sativus* L. (var. Cherry red) were secured from Vegetables Department, Horticulture Research Institute, ARC, Egypt and planted on October 22nd during both growing seasons. Soil samples at 30 cm depth were taken for estimating the experimental soils physical and chemical characteristics (Table 1) according to (Black *et al.*, 1965).

Table 1. Physical and chemical characteristics of the experimental soil (average of the two growing seasons).

Texture	Organic matter%	Total carbon%	Ec (dsm-1)	pН	N%	P%	K%
Clay	1.8	1.82	0.75	8.65	0.19	0.015	0.14

Experimental design

The experimental design was randomized complete block design with three replicates. Each experimental unit contained three rows, 6 meters in length and 0.8 meters in width, with the total experimental area of 14.4 m². Within the row, the distance between plants was 20 cm.

Sampling procedures

Plants were sprayed twice, 20 and 30 days after sowing (DAS) onto foliage with solutions of Seaweed Ecklonia maxima extract (SWE) at the rate of 4 ml L⁻¹, Humic acid (HA) at the rate of 30 ml L⁻¹, Peppermint oil (P.O) at the rate of 50 ml L⁻¹, and Thyme oil (T.O) at the rate of 50 ml L⁻¹. Control plants were sprayed with tap water. Tween 20 was added to sprayed solutions at the rate of 0.05 % as a wetting agent. Seaweed extract and humic acid were obtained from Shoura Company, Egypt. Peppermint and thyme oils were obtained by hydro-distillation for 2-3 h in the Post-harvest Lab., Hort. Res. Station, Agric. Res. Cent., Egypt, using modified Clevenger apparatus according to (Guenter, 1965). Samples were collected 40 DAS to determine the contents of anthocyanin, total phenols, total flavonoids, ascorbic acid, total carbohydrates, total free amino acids as well as the activity of Peroxidase and Polyphenol oxidase.

Anthocyanin determination was determined according to (Chiriboga & Francis, 1973). Total phenols were measured calorimetrically at 650 nm using Folin-ciocalteu reagent according to (Bray & Thorpe, 1954). Total flavonoids content was estimated calorimetrically at 510 nm as described by Heimler *et al.*, (2006). Ascorbic acid was determined by the 2,6-Dichlorophenolindophenol dye procedure of (Freed, 1966). Total carbohydrates were

determined using phenol sulphuric acid method according to (Dubois *et al.*, 1956). Total free amino acids were determined according to the procedure of (Salnikow *et al.*, 1973). Peroxidase extraction and assay were performed according to the procedure of (Maxwell & Bateman, 1967). Polyphenol oxidase (PPO) extraction and assay were performed according to (Galeazzi *et al.*, 1981). Sulfur estimated by using atomic absorption spectroscopy (AAS) was subjected by Kirkbright & Wilson, (1974). Potassium was determined (mg/g) by using flame photometer (Model 400M.X) according to (Chapman & Pratt, 1961).

Storage parameters

Plants were uprooted 60 DAS and samples from roots (ten roots from each replicate) were packed in polyethylene bags and stored at controlled storage conditions (5°C, relative humidity 95%) for three months. Afterwards, postharvest quality and storage parameters represented by weight loss percentage, post-harvest decay percentage and dry matter percentage were estimated. Weight loss percentage (WLP %) was estimated according to the following equation: WLP % = [Initial - (Final Weight /Initial Weight)] × 100 (A.O.A.C, 2007). Post-harvest decay percentage (PDP %) was estimated according to the formula: Number of Decayed Fruits/ Number of Total Fruits x 100 (El-Mougy et al., 2012). Dry matter percentage (DMP %) was estimated as described by Lee, (1981). After drying in the oven at 60°C until weight stability, moisture percentage was calculated according to the equation.

Statistical analysis

Data were subjected to analysis of variance using GENSTATE software (version 11.1.0.1575). To compare differences between means, least significant differences (LSD) at 5% were calculated according to (Gomez *et al.*, 1984). Correlation matrix between biochemical analysis

after storage in refrigerator at (5°C/Rh 95%) and storage parameters at the 0.05 and 0.01 level (2-tailed) on PASW Statistics18 software. The sign of the correlation coefficient determines whether the correlation is positive or negative. The magnitude of the correlation coefficient determines the strength of the correlation subjected by (Evans, 1996) suggests absolute r-value 0.00-0.19 (very weak), 0.20-0.39 (weak), 0.40-0.59 (moderate), 0.60-0.79 (strong) and 0.80-1.0 (very strong).

RESULTS AND DISCUSSION

All applied bio modulators increased total anthocyanin's concentration in the roots either pre- or post-harvest compared to control (Table 2). The highest anthocyanins concentrations were obtained in response to treatment with thymus oil followed by humic acid in pre-harvest stage. In roots that stored in the specified storage experimental conditions, total anthocyanin concentration was decreased. either in biomodulators-treated or untreated plants compared with before storage. However, in post-harvest stage anthocyanins concentrations were highest in response to the treatment with peppermint oil followed by seaweed extract i.e. peppermint oil and seaweed extract caused the least decline in anthocyanins concentration in stored roots, in order. Data presented in table (2) show that ascorbic acid concentration in pre-stored roots was increased in response to thyme oil and seaweed extract, though the increase was insignificant in case of seaweed extract treatment. On the other hand, humic acid and peppermint oil did not affect ascorbic acid concentration in the roots before storage. In stored roots, ascorbic acid concentration was decreased either in bio modulators-treated or untreated plants. However, the magnitude of decline was lower in roots of bio modulators-treated plants. So, after storage, ascorbic acid concentration in roots was significantly higher in all biomodulators treatments compared with control. The lowest loss percentage was recorded in response to peppermint oil (29.7 %) followed by humic acid (31.7 %).

Table 2. Effect of applied bio modulators on total anthocyanin and ascorbic acid content (mg g⁻¹. F.W) in roots of *Raphanus sativus* plants before and after storage.

Parameters	Ant	hocyanin (m	ngg ⁻¹ .F.W)	Ascorbic acid (mgg ⁻¹ .F.W)				
Treatments	Before	After	Total loss%	Before	After	Total loss %		
Seaweed extract	8.34	3.71	55.52	24.13	13.97	42.11		
Humic acid	8.51	2.65	68.86	20.50	14.00	31.71		
Peppermint oil	8.22	4.54	44.77	20.52	14.42	29.71		
Thyme oil	8.55	2.60	69.59	37.10	22.57	39.16		
Control	7.16	1.44	79.89	20.41	9.23	78.54		
LSD at 0.05	$0.60^{\rm c}$	0.29^{d}	-	6.32^{a}	$1.50^{\rm b}$	-		

Table 3. Effect of applied bio modulators on total phenols and flavonoids (mg g⁻¹.D.W) roots of *Raphanus sativus* plants before and after storage.

Parameters	Tot	al phenols (m	gg ⁻¹ .F.W)	Total flavonoids (mg g ¹ .F.W)				
Treatments	Before	After	Total loss%	Before	After	Total loss%		
Seaweed extract	14.49	12.08	16.63	0.33	0.29	12.12		
Humic acid	13.41	10.78	19.61	0.40	0.35	12.50		
Peppermint oil	14.85	12.45	16.16	0.39	0.34	12.82		
Thyme oil	13.00	10.42	19.85	0.47	0.44	6.38		
Control	12.50	9.51	23.92	0.31	0.26	16.13		
LSD at 0.05	2.30^{a}	$1.02^{\rm b}$	-	$0.01^{\rm d}$	0.03°	-		

Table4. Effect of applied biomodulators on total carbohydrates (mgg⁻¹. D.W) and free amino acids (mg/100g.D.W) in dried roots of *Raphanus sativus* plants before and after storage.

Parameters	Total car	rbohydrates (mg	g ⁻¹ .D.W)	Total free amino acid (mg/100g.D.W)			
Treatments	Before	After	Total loss%	Before	After	Total loss%	
Seaweed extract	127.50	104.80	17.80	12.09	5.64	53.35	
Humic acid	127.03	90.83	28.50	12.49	4.99	60.05	
Peppermint oil	126.60	101.92	19.49	13.56	4.34	67.99	
Thyme oil	112.12	84.33	27.46	12.66	5.20	58.93	
Control	110.90	79.00	28.76	12.03	3.30	72.57	
LSD at 0.05	22.64 ^b	25.74 ^a	-	3.73°	0.49^{d}	-	

Table 5. Effect of applied biomodulators on potassium concentration (mgg⁻¹.D.W.) and sulphur percentage in dried roots of *Raphanus sativus* plants before and after storage.

Parameters	Po	otassium (mg	g ⁻¹ D.W.)	Sulfur (%)				
Treatments	Before	After	Total loss%	Before	After	Total loss%		
Seaweed extract	12.48	11.00	11.86	1.43	0.93	34.97		
Humic acid	13.00	10.00	23.08	1.20	0.90	25.00		
Peppermint oil	13.64	10.09	26.03	1.50	1.15	23.33		
Thyme oil	13.26	10.07	24.06	1.29	1.01	21.71		
Control	12.37	9.00	27.24	1.10	0.50	54.55		
LSD at 0.05	0.15^{d}	0.71^{a}	-	0.48^{b}	0.35^{c}	-		

All applied bio modulators increased total flavonoids in roots during both pre- and post-harvest stages as well as total phenols only during post-harvest stage (Table 3). The only bio modulator that increased total phenols in the roots before storage was Peppermint oil. Both total flavonoids and phenols were decreased after storage, but the decrease was lower in case of total flavonoids. The lowest decrease in total phenols was recorded in response to Peppermint oil whereas the least decrease in total flavonoids was recorded in response to thyme oil treatment.

Generally, total carbohydrates concentration in roots before storage was increased in response to bio modulators treatments, though the increase was insignificant (Table 4). In addition, roots of bio modulators-treated plants after storage contained higher concentrations of total carbohydrates compared with control roots, though the differences did not reach the significance level. In all treatments, total carbohydrates concentrations were decreased after storage, with the least decrease in response to seaweed extract followed by peppermint oil treatment.

Total free amino acids concentrations in roots treated with all bio modulators before storage were not significantly different than that in control roots (Table 4). On the other hand, roots of bio modulators-treated plants after storage contained significantly higher concentrations of total free amino acids compared with control. Storage caused a decrease in total free amino acids concentrations in roots of bio modulators-treated as well as control plants. In stored roots of seaweed extract-treated plants, the decrease in concentration of total free amino acids was the least, followed by thyme oil.

Roots of biomodulators-treated plants either before or after storage contained higher concentrations of both potassium (K) and sulphur (S) compared with control roots (Table 5). However, the recorded increase in S concentration before storage was insignificant in response to all biomodulators treatments. In roots of biomodulators-treated as well as control plants, concentrations of both potassium K and S were decreased after storage. The decrease was of the least magnitude in response to seaweed extract treatment in case of K and in response to peppermint oil in case of S. In roots of biomodulators treated plants, the activity of peroxidase (POD) was increased either before or after Storage compared with that

in roots of control plants. A similar trend in the activity of polyphenol oxidase (PPO) was recorded in roots after storage. On the other hand, before storage, PPO activity in roots was increased in response to seaweed extract and peppermint oil treatments whereas decreased in response to humic acid and thyme oil treatments. Storage affected the activity of both POD and PPO, but in different manner. Where POD activity was decreased after storage, PPO activity was increased (Figure 1). The least decrease in POD activity was recorded in response to thyme oil treatment, whereas the highest increase in PPO activity was achieved in response to humic acid treatment.




Figure 1. Quality parameters of stored roots of *Raphanus sativus* plants after storage for three months in refrigerator at 5°C and 95% relative humidity as affected by biomodulators treatments.

Effect of biomodulators on post-harvest quality parameters of stored roots

After two months-storage, stored roots had visible decay symptoms manifested by rotting, water stains and unusual smell due to growth of bacteria and fungus on roots. However, quality parameters of stored roots were differed between biomodulators treatments and control. All applied biomodulators decreased weight loss and post-harvest decay percentages hence, roots of biomodulators-treated plants contained higher dry matter compared with the roots of untreated plants. Weight loss as well as post-harvest decay percentage was minimum in response to peppermint treatment, followed by thyme oil treatment. Nevertheless, dry matter percentage in stored roots was highest due to

seaweed extract followed by humic acid treatments (Figure 2).

Pearson's correlation matrix between roots biochemical analyses after storage and their post-harvest quality parameters

In table (6) showed the output of Pearson's correlation matrix between biochemical analysis after storage in refrigerator at (5°C/Rh 95%) for three months and storage parameters of radish plants in first season 2015. Dry matter and post-harvest decay showed very strong significant negative as follow (-0.90 and -0.99) respectively relationship with sulfur concentration at the 0.05 and 0.01 level. Total anthocyanin revealed very strong significant positive as follow (0.98 and 0.92) relationship with total

phenols and carbohydrates at the 0.05 and 0.01 level. Ascorbic acid, total phenols, flavonoids and anthocyanin recorded very strong significant positive as follow (0.95,

0.97, 0.89 and 0.91) respectively relationship with total flavonoids, carbohydrates, potassium concentration and peroxidase enzyme at the 0.05 and 0.01 level.




Figure 2. Effect of applied bio modulators on the activity (U g⁻¹ F.W.) of Peroxidase (POD) and Polyphenoloxidase (PPO) in roots of *Raphanus sativus* plantsbefore and after storage.

Table 6. Pearson's correlation matrix between roots biochemical analyses after storage and their post-harvest quality parameters.

Biochemical	DM	PDP	WLP	T.A	ASA	T.P	T.F	T.C	T.A	K	S	POD	PPO
Parameters	%	%	%										
DM%													
PDP%	0.88												
WLP%	-0.30	-0.61											
T. A	-0.83	-0.83	0.72										
ASA	-0.57	-0.69	0.00	0.21									
T. P	-0.73	-0.77	0.79	0.98**	0.11								
T. F	-0.52	-0.66	-0.02	0.14	0.95*	0.02							
T.C	-0.57	-0.66	0.85	0.92*	-0.00	0.97**	-0.12						
T.A	-0.35	-0.70	0.66	0.46	0.62	0.51	0.47	0.56					
K	-0.48	-0.72	0.78	0.71	0.38	0.77	0.19	0.83	0.91*				
S	-0.90*	-0.99**	0.61	0.86	0.63	0.79	0.62	0.67	0.63	0.67			
POD	-0.74	-0.81	0.20	0.45	0.78	0.31	0.89*	0.13	0.35	0.21	0.82		
PPO	-0.72	-0.64	0.57	0.86	-0.04	0.81	0.05	0.70	0.05	0.29	0.71	0.49	

Correlation is significant at the 0.05 level (2-tailed). Correlation is significant at the 0.01 level (2 tailed), DM: dry matter PDP: post-harvest decay, WLP, weight loss, TA, total anthocyanin, TP, total phenols, ASA, ascorbic acid, TF, total flavonoids, TC, total carbohydrates, TA, total amino acids, K, potassium, S- sulphur, POD-peroxidase, PPO- polyphenoloxidase.

Senescence is the most important internal factor that causes damage to vegetables whereas low temperature is the best way to delay post-harvest deterioration of fruits and vegetables. In this context, Bayoumi (2008) concluded that the higher post-harvest decay percentage in late harvesting stage of fruits is due to higher rate of respiration, more skin permeability for water loss and high susceptibility to decay. Dry matter acts an index to evaluate the fruit quality due to endure prolonged storage (Jackson & Harker, 1997). Generally, low temperature can maintain organic matter and vitamins levels in short and long term storage because most enzyme activities decrease and the expression of many genes inhibited during cold storage. Low temperature markedly delays senescence

of broccoli during storage (Javanmardi & Kubota, 2006). In addition, Shen *et al.* (2013) reported that refrigeration can reduce deterioration and extend shelf life of fresh fruits by delaying the metabolic processes. Dry matter production represents a balance between photosynthesis and respiration. Respiration rate, which has long been used to measure metabolic process in stored produce (Kittock & Law, 1968; Scholz, 1962), is governed by water availability, temperature, O₂ concentration, microbial contamination, mechanical damage, among other factors. (Porter *et al.*, 2003) reported that respiration rate of chinese cabbage was higher at 20°C than at lower temperatures.

Synthetic chemicals are commonly used in the control diseases of stored plant products however; these chemicals

may cause toxic residues in treated plants (Isman, 2000). Alternatively, the use of natural products like herbal extracts and essential oils can reduce harm to public health and preserve the environment from pollution. These ecofriendly plant-derived products generally possess a broad spectrum of activity against several pathogens and pests (Bakkali et al., 2008) Essential oils are natural complex compounds with a strong smell and produced as secondary metabolites in aromatic plants. In nature, essential oils act as antibacterial, antivirus, antifungal and insecticides. It contains compositions of terpenes, sesquiterpenes, aldehydes, ketones and phenolic compounds (Burt, 2004). The main chemical components of thyme oil are thymol, carvacrol, α- pinene, β-pinene, borneal, linalool, β-simine and camphene (Dew, 1984). The composition of volatile oil of *M. piperita* is menthol, monoterpene and methofuran (Dew, 1984) These constituents act as inhibitor for pathogens growth during cold storage and reduce senescence in treated plants.

In the current investigation, thyme as well as peppermint oils decreased weight loss and post-harvest decay percentages hence, preserved roots dry matter during cold storage (Figure 2). These findings are in harmony with those of previous studies (Aminifard & Mohammadi, 2013; Alikhani et al., 2009; Amin, 2016; Bakry et al., 2013; Geransayeh et al., 2015). Antioxidants play an important role in scavenging reactive oxygen species (ROS) that appear during storage-related senescence of vegetables (Hounsome et al., 2009). Antioxidant capacity is directly correlated with phenolic compounds (Koh et al., 2009; Tavarini et al., 2008). Therefore, plants with enhanced levels of antioxidants can resist oxidative damage (Navarro et al., 2006). The results of the present investigation revealed that generally, the applied biomodulators diminished the decline in concentrations of anthocyanins, ascorbic acid, phenols and flavonoids as well as in the activity of the antioxidant enzyme, peroxidase. This implies that stored roots of biomodulators-treated plants contained higher levels of antioxidants compared with roots of untreated plants. Similar conclusions were made based on the results of previous studies (Del Nobile et al., 2007; Kramchote et al., 2012; Wang et al., 2008; Žnidarčič & Požrl, 2006). How elevated levels from these constituents preserve roots quality during storage could be discussed as follows.

Anthocyanin belongs to flavonoids group and located in a cell vacuole (Friend & Rhodes, 1981). It plays important role as an antioxidant. The anthocyanin pigment inhibits active oxygen radicals formed by exposure to stress conditions that occur during storage periods due to pathological injures i.e., enzymatic browning, molds and water stains on fruits. The results of some in vitro experiments indicated that anthocyanin pigments scavenge superoxide, active oxygen radicals (Yamasaki *et al.*, 1996) and hydrogen peroxide (Leng & Qi, 2003). On the other hand, anthocyanin pigment is affected by the amount of sugar due to a lower water activity (Hounsome *et al.*,

2009). Andreu *et al.*, (1988) reported that anthocyanins concentration was decreased at the end of the cold storage period, which implies that they are consumed in scavenging storage-related ROS.

Ascorbic acid is a major component of the plant defense system (Hodges & Forney, 2000) that protect biological processes of the plant from ROS generated during biotic and abiotic stresses (Foyer & Noctor, 2011). The destruction of ascorbic acid is one of the most serious problems facing plants under stress conditions, whether in the soil or when storing in the short or long term (Jeney et al., 2008). The oxidized form of ascorbic acid is more prone to decomposition during storage in low temperature. Ascorbic acid decreases during storage due to its consumption in the ROS-detoxification process especially at temperature above 0°C (Ajibola et al., 2009) through oxidation in the presence of ascorbate oxidase enzyme, and changes backto its active form (Lee & Kader, 2000). The antioxidant role of ascorbic acid in mitigating stress-related metabolic abnormalities was evident from the results of other studies (Al-Karaki, 2000; Aminifard & Mohammadi, 2013; Raafat et al., 2012; Serrano et al., 2005; Wang et al., 2008).

Post-harvest stresses e.g. mechanical injuries and others during storage led to enzymes activity that responsible for phenolic compounds deterioration (Yang et al., 2011). Phenolic compounds and enzymes do not interact with each other because of their different locations within cells consequently; shelf life of vegetables and fruits is increased. However, when membranes are damaged, destruction and oxidation of phenolic compounds are initiated, causing breakdown of defense line in the cell against free radicals (Toor & Savage, 2006; Yang et al., 2011). Similar conclusions were made by Amin (2016); Wang et al. (2008). Flavonoids are the most important group of phenolic compounds in plants (Hounsome et al., 2009) that assist in scavenging free radicals and inhibiting oxidative stress, thereby decreasing cell membrane deterioration during long storage (Koh et al., 2009) which low temperature could decrease the membrane lipid degradation and POD activity as a result, it can slow down flavonoids oxidation. These results are in agreement with those obtained by DuPont et al. (2000) on lettuce, (Yuan et al., 2010) on broccoli plants and (Geransayeh et al., 2015) on strawberry fruits.

This study showed increased photosynthesis and increased plant growth rate with adding HA and SWE treatments in the open field led to an increase in the rate of carbohydrate content. These results are in harmony with those recorded by Tehranifar & Ameri, (2012) on *Fragaria ananasa*. Antioxidants applied in the field experiment led to a significant increase in storage of fruits due to increased carbohydrates anabolism process within plant cells, (Blunden & Wildgoose, 1977) mentioned that seaweed extract applied on potato plants recorded significantly increased in the yield of tubers for both the varieties King

Edward and Pentland Dell due to plants hormones such as cytokinins represent the most important compound in SWE that affect on all phytochemicals before storage on the other hand, oxidative damage, enzymatic browning is responsible for the degradation of phenolic compounds especially with long storage time at specified temperature where this phenomenon and increased the rate of respiration cause decreasing phytochemicals which increased within storage intervals. Accordingly, the rate of broccoli plants respiration effect on carbohydrate levels when storage in cold temperature at 5°C. These results are in agreement with those reported by Vigna radiate plants that also supported by Majidi et al. (2014) reported that decrease in fruit metabolic activities results in a decrease in fruit water loss and carbohydrates depletion rate and for this reason senescence process delayed in fruits in shortterm storage at specific temperature. The present study supported results of (Kalidass et al., 2010) found the free amino acid concentration were also more in Brassica nigra treated with 10% SWE of P. teriastromatica and S. linearifolium And also, (Bakry et al., 2013) reported the free amino acids may be act as protective agents of membrane that protect the cells from biotic and a biotic stress pre or postharvest on two wheat cultivars. The content of amino acid components in dried roots of radish plants changed with different storage intervals and temperature. The results were showed significantly increase with peppermint oil compared with control plants. Accordingly, application of P.O reduced total amino acids degradation in the roots of radish plants. In general, the percentage of total amino acids decreased due to low respiratory rate and enzymatic activity (Kalidass et al., 2010).

S plays an important role in the growth and development of higher plants (Osman & Rady, 2012). S deficiency affects on synthesis of chlorophyll, nitrogen concentration in leaves and photosynthetic enzymes (Lunde et al., 2008) led to reduce the quality and quantity of yield (Hawkesford, 2000). Adequate S nutrition has regulatory interactions with N assimilation and also can enhance the photosynthesis process (Scherer, 2008). S is an important constituent for N assimilation, protein and co-enzymes synthesis. The results obtained by Mackowiak et al., (2001) revealed that the application of humic and SWE may be increased the uptake of mineral elements. Many study supported these results such as (Tehranifar & Ameri, 2012) on Fragaria ananasa L., and (Boghdady et al., 2016) on chickpea seeds. (Ayas & Gulser, 2005) showed the application of humic acid at rate of (250 g) on spinach led to increase nutrient content, amino acid and sugars concentration. On the other hand after the storage intervals, the respiration rate increase as well as S content and the total amount of protein decreased Potassium is an important nutrient involved in transpiration, stomatel opening, osmoregulation, photosynthesis and enzymes activation; 2005) Also K participated in critical physiological processes such as transport of food so phloem, osmotic potential of cells as well as cell extension wall (Chérel, 2004; Güler & Ibrikci, 2000). Watcharasak & Thammasak, (2005) showed that antioxidants applied on leaves led to significantly increased in potassium concentration in tomato plants. The application of antioxidants as a foliar spray on leaves increased nutrients uptake of the soil helped to form the roots and reach the appropriate size of the harvest that reflected on the storage intervals that maintained the roots from oxidation processes and enzymatic browning (Bombelli & Wright, 2006). Similar results were also reported on radish plants by (Ayub *et al.*, 2013).On *Spinacia oleraceae* plants by Fan *et al.* (2014).

REFERENCES

- Ajibola, V., Babatunde, O., & Suleiman, S. (2009). The effect of storage method on the vitamin C content in some tropical fruit juices. *Trends in Applied Science Research*, 4(2), 79-84.
- Al-Karaki, G. N. (2000). Growth, sodium, and potassium uptake and translocation in salt stressed tomato. *Journal of Plant Nutrition*, 23(3), 369-379.
- Alikhani, M., Sharifani, M., Azizi, M., Mousavizadeh, S., & Rahimi, M. (2009). Increasing shelf life and maintaining quality of strawberry (*Fragaria ananassa* L.) with application of mucilage edible coating and plant essential oil. *Journal of Agricultural Sciences and Natural Resources*, 16(2).
- Amin, A. (2016). Determination of some essential oils effects on the quality traits of the Egyptian Anna apple fruit during its shelf life. *Journal of Horticultural Science and Ornamental Plants*, 8(1), 35-45.
- Aminifard, M. H., & Mohammadi, S. (2013). Essential oils to control Botrytis cinerea in vitro and in vivo on plum fruits. *Journal of the Science of Food and Agriculture*, 93(2), 348-353.
- Andreu, F. L., Mollá, E., Fernández, M., & Álvarez, R. M. E. (1988). Conservación en frío de berenjenas: Modificaciones en su calidad. *Alimentaria: Revista de Tecnología e Higiene De Los Alimentos*(192), 31-36.
- Ayas, H., & Gulser, F. (2005). The effects of sulfur and humic acid on yield components and macronutrient contents of spinach (Spinacia Oleracea Var. Spinoza). *Journal of Biological Science*,5(6),801-804.
- Ayub, R. A., Spinardi, B., & Gioppo, M. (2013). Storage and fresh cut radish. *Acta Scientiarum. Agronomy*, 35(2), 241-245.
- Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils—a review. *Food and Chemical Toxicology*, *46*(2), 446-475.
- Bakry, A., Sadak, M. S., Moamen, H., & Abd El Lateef, E. (2013). Influence of humic acid and organic fertilizer on growth, chemical constituents, yield and quality of two flax seed cultivars grown under newly reclaimed

- sandy soils. *International Journal of Academic Research*, 5(5).
- Bakry, B., Elewa, T., El-Kramany, M., & Wali, A. (2013). Effect of humic and ascorbic acids foliar application on yield and yield components of two wheat cultivars grown under newly reclaimed sandy soil. *International Journal of Agronomy and Plant Production*, 4(6), 1125-1133.
- Bayoumi, Y. A. (2008). Improvement of postharvest keeping quality of white pepper fruits (*Capsicum annuum*, L.) by hydrogen peroxide treatment under storage conditions. *Acta Biologica Szegediensis*, 52(1), 7-15.
- Black, C., Evans, D., White, J., Ensmingher, L., Clarke, F., & Dinauer, R. (1965). Methods of soil analysis. Part 2chemical and microbiological properties, 9 series agronomy. American Society of Agronomy. *Inc, Madison, Wisconsin, USA*.873-895.
- Blunden, G., & Wildgoose, P. B. (1977). The effects of aqueous seaweed extract and kinetin on potato yields. *Journal of the Science of Food and Agriculture*, 28(2), 121-125.
- Boehme, M., Schevtschenko, J., & Pinker, I. (2005). Iron supply of cucumbers in substrate culture with humate. *Acta Horticulturae*, 697, 329.
- Boghdady, M. S., Selim, D. A., Nassar, R. M., & Salama, A. (2016). Influence of foliar spray with seaweed extract on growth, yield and its quality, profile of protein pattern and anatomical structure of chickpea plant (Cicer arietinum L.). Middle East Journal of Applied Sciences, 6(1), 207-221.
- Bombelli, E. C., & Wright, E. R. (2006). Tomato fruit quality conservation during post-harvest by application of potassium bicarbonate and its effect on Botrytis cinerea. Cien. Invention. Agriculture.(In English) 33 (3): 167-172. Ciencia e Investigación Agraria, 33(3), 167-172.
- Bray, H., & Thorpe, W. (1954). Analysis of phenolic compounds of interest in metabolism. *Methods of Biochemical Analysis*, 27-52.
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods—a review. *International Journal of Food Microbiology*, 94(3), 223-253.
- Cakmak, I. (2005). The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *Journal of Plant Nutrition and Soil Science*, 168(4), 521-530.
- Chapman, H., & Pratt, F. (1961). Ammonium vandate-molybdate method for determination of phosphorus. *Methods of Analysis for Soils, Plants and Water, 1*, 184-203.
- Chérel, I. (2004). Regulation of K+ channel activities in plants: from physiological to molecular aspects. *Journal of Experimental Botany*, 55(396), 337-351.

- Chiriboga, C., & Francis, F. (1973). Ion exchange purified anthocyanin pigments as a colorant for cranberry juice cocktail. *Journal of Food Science*, *38*(3), 464-467.
- Cho, E.H., Choi, A., Choi, S.J., Kim, S.Y., Lee, G.S., Lee, S.S., & Chae, H.J. (2009). α-amylase activity of radish and stability in processing. *Journal of the Korean Society of Food Science and Nutrition*, 38(6), 812-815.
- Del Nobile, M. A., Licciardello, F., Scrocco, C., Muratore, G., & Zappa, M. (2007). Design of plastic packages for minimally processed fruits. *Journal of Food Engineering*, 79(1), 217-224.
- Dew, M. (1984). Peppermint oil for the irritable bowel syndrome: a multicentre trial. *Br J Clin Pract*, *38*, 398.
- Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. t., & Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28(3), 350-356.
- DuPont, M. S., Mondin, Z., Williamson, G., & Price, K. R. (2000). Effect of variety, processing, and storage on the flavonoid glycoside content and composition of lettuce and endive. *Journal of Agricultural and Food Chemistry*, 48(9), 3957-3964.
- El-Mougy, N., Abdel-Kader, M., & Aly, M. (2012). Effect of a new chemical formula on postharvest decay incidence in citrus fruit. *Journal of plant protection research*, 52(1), 156-164.
- Evans, J. D. (1996). *Straightforward statistics for the behavioral sciences*: Thomson Brooks/Cole Publishing Co.
- Fan, D., Kandasamy, S., Hodges, D. M., Critchley, A. T., & Prithiviraj, B. (2014). Pre-harvest treatment of spinach with Ascophyllum nodosum extract improves post-harvest storage and quality. *Scientia horticulturae*, 170, 70-74.
- Foyer, C. H., & Noctor, G. (2011). Ascorbate and glutathione: the heart of the redox hub. *Plant Physiology*, *155*(1), 2-18.
- Freed, M. (1966). Methods of vitamin assay. John Wiley and Sons, New York, 1-424.
- Friend, J., & Rhodes, M. J. (1981). *Recent advances in the biochemistry of fruits and vegetables*: Academic Press.
- Galeazzi, M. A., Sgarbieri, V. C., & Constantinides, S. M. (1981). Isolation, purification and physicochemical characterization of polyphenoloxidases (PPO) from a dwarf variety of banana (*Musa cavendishii*, L). *Journal* of Food Science, 46(1), 150-155.
- Geransayeh, M., Sepahvand, S., Abdossi, V., & Nezhad, R. A. (2015). Effect of thymol treatment on decay, postharvest life and quality of strawberry (Fragaria ananassa) Fruit cv. 'Gaviota'. *International Journal of Agronomy and Agricultural Research*, 6, 151-162.

- Gomez, K. A., Gomez, K. A., & Gomez, A. A. (1984). Statistical Procedures for Agricultural Research: John Wiley & Sons.1-680.
- Guenter, E. (1965). The essential oils (Vol. III): New York: Van Nostrand.379-382.
- Güler, S., & Ibrikci, H. (2000). Yield and elemental composition of cucumber as affected by drip and furrow irrigation. Paper presented at the Workshop Towards and Ecologically Sound Fertilisation in Field Vegetable Production, 571.
- Gutiérrez, R. M. P., & Perez, R. L. (2004). Raphanus sativus (Radish): their chemistry and biology. *The Scientific World Journal*, *4*, 811-837.
- Hawkesford, M. J. (2000). Plant responses to sulphur deficiency and the genetic manipulation of sulphate transporters to improve S-utilization efficiency. *Journal of Experimental Botany*, *51*(342), 131-138.
- Heimler, D., Vignolini, P., Dini, M. G., Vincieri, F. F., & Romani, A. (2006). Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. *Food Chemistry*, 99(3), 464-469.
- Hodges, D. M., & Forney, C. F. (2000). The effects of ethylene, depressed oxygen and elevated carbon dioxide on antioxidant profiles of senescing spinach leaves. *Journal of Experimental Botany*, 51(344), 645-655.
- Hounsome, N., Hounsome, B., Tomos, D., & Edwards-Jones, G. (2009). Changes in antioxidant compounds in white cabbage during winter storage. *Postharvest Biology and Technology*, 52(2), 173-179.
- Hwang, J.W., Kim, E.K., Lee, S.J., Kim, Y.S., Choi, D.K., Park, T.K Park, P.J. (2012). Anthocyanin effectively scavenges free radicals and protects retinal cells from H₂ O₂-triggered G2/M arrest. *European Food Research and Technology*, 234(3), 431-439.
- Isman, M. B. (2000). Plant essential oils for pest and disease management. *Crop protection*, 19(8-10), 603-608.
- Jackson, P., & Harker, F. (1997). Changes in firmness of the outer pericarp, inner pericarp, and core of Actinidia species during ripening. New Zealand Journal of Crop and Horticultural Science, 25, 185-189.
- Javanmardi, J., & Kubota, C. (2006). Variation of lycopene, antioxidant activity, total soluble solids and weight loss of tomato during postharvest storage. *Postharvest Biology and Technology*, 41(2), 151-155.
- Jeney-Nagymate, E., & Fodor, P. (2008). The stability of vitamin C in different beverages. *British Food Journal*, 110(3), 296-309.
- Justesen, U., & Knuthsen, P. (2001). Composition of flavonoids in fresh herbs and calculation of flavonoid intake by use of herbs in traditional Danish dishes. *Food Chemistry*, 73(2), 245-250.

- Kader, A. A. (2003). A perspective on postharvest horticulture (1978-2003). *HortScience*, 38(5), 1004-1008.
- Kalidass, C., Jayarani, S., & Glory, M. (2010). Effect of seaweed liquid fertilizer on growth and biochemical constituents of Brassica nigra Linn. *International Journal of Agriculture Environment & Biotechnology*, 3(3), 307-311.
- Kapoor, L. (2000). CRC Handbook of Ayurvedic Medicinal Plants CRC Press. *Inc. Boca Raton, Florida,* 117.
- Karakurt, Y., Unlu, H., Unlu, H., & Padem, H. (2009). The influence of foliar and soil fertilization of humic acid on yield and quality of pepper. Acta Agriculturae Scandinavica Section B Soil and Plant Science, 59(3), 233-237.
- Kasim, W. A., Hamada, E. A., El-Din, N. G. S., & Eskander, S. (2015). Influence of seaweed extracts on the growth, some metabolic activities and yield of wheat grown under drought stress. *Journal of Agronomy and Agricultural Research*, 7, 173-189.
- Kirkbright, G., & Wilson, P. (1974). Application of a demountable hollow cathode lamp as a source for the direct determination of sulfur, iodine, arsenic, selenium, and mercury by atomic absorption flame spectrometry. *Analytical Chemistry*, 46(11), 1414-1418.
- Kittock, D., & Law, A. (1968). Relationship of Seedling Vigor to Respiration and Tetrazolium Chloride Reduction by Germinating Wheat Seeds 1. *Agronomy Journal*, 60(3), 286-288.
- Koh, E., Wimalasiri, K., Chassy, A., & Mitchell, A. (2009). Content of ascorbic acid, quercetin, kaempferol and total phenolics in commercial broccoli. *Journal of Food Composition and Analysis*, 22(7-8), 637-643.
- Kramchote, S., Srilaong, V., Wongs-Aree, C., & Kanlayanarat, S. (2012). Low temperature storage maintains postharvest quality of cabbage (Brassica oleraceae var. capitata L.) in supply chain. *International Food Research Journal*, 19(2),759-763.
- Ku, K.H., Lee, K.A., & Kim, Y.E. (2008). Physiological activity of extracts from radish (*Raphanus sativus L.*) leaves. *Journal of the Korean Society of Food Science and Nutrition*, *37*(3), 390-395.
- Lee, S.K. (1981). A review and background of the avocado maturity standard. *California Avocado Society Yearbook*, 65, 101-109.
- Lee, S. K., & Kader, A. A. (2000). Preharvest and postharvest factors influencing vitamin C content of horticultural crops. *Postharvest Biology and Technology*, 20(3), 207-220.
- Leng, P., & Qi, J. (2003). Effect of anthocyanin on David peach (*Prunus davidiana* Franch) under low

- temperature stress. *Scientia Horticulturae*, 97(1), 27-39
- Lunde, C., Zygadlo, A., Simonsen, H. T., Nielsen, P. L., Blennow, A., & Haldrup, A. (2008). Sulfur starvation in rice: the effect on photosynthesis, carbohydrate metabolism, and oxidative stress protective pathways. *Physiologia Plantarum*, 134(3), 508-521.
- Mackowiak, C., Grossl, P., & Bugbee, B. (2001). Beneficial effects of humic acid on micronutrient availability to wheat. *Soil Science Society of America Journal*, 65(6), 1744-1750.
- Majidi, H., Minaei, S., Almassi, M., & Mostofi, Y. (2014). Tomato quality in controlled atmosphere storage, modified atmosphere packaging and cold storage. *Journal of Food Science and Technology*, 51(9), 2155-2161.
- Maxwell, D., & Bateman, D. (1967). Changes in the activities of some oxidases in extracts of Rhizoctonia-infected bean hypocotyls in relation to lesion maturation. *Phytopathology*, *57*, 132-136.
- Nardi, S., Pizzeghello, D., Muscolo, A., & Vianello, A. (2002). Physiological effects of humic substances on higher plants. *Soil Biology and Biochemistry*, 34(11), 1527-1536.
- Navarro, J. M., Flores, P., Garrido, C., & Martinez, V. (2006). Changes in the contents of antioxidant compounds in pepper fruits at different ripening stages, as affected by salinity. *Food Chemistry*, 96(1), 66-73.
- Nguefack, J., Somda, I., Mortensen, C. N., & Amvam Zollo, P. (2005). Evaluation of five essential oils from aromatic plants of Cameroon for controlling seedborne bacteria of rice (*Oryza sativa L.*). *Seed Science and Technology*, 33(2), 397-407.
- Ördög, V., Stirk, W. A., Van Staden, J., Novák, O., & Strnad, M. (2004). Endogenous Cytokinins in three genera of microalgae from the chlorophyta 1. *Journal of Phycology*, 40(1), 88-95.
- Osman, A. S., & Rady, M. M. (2012). Ameliorative effects of sulphur and humic acid on the growth, anti-oxidant levels, and yields of pea (*Pisum sativum L.*) plants grown in reclaimed saline soil. *The Journal of Horticultural Science and Biotechnology*, 87(6), 626-632.
- Paredes, S. (1984). Etnobotánica Mexicana: Plantas popularmente empleadas en el Estado de Michocán en el tratamiento de enfermedades hepaticas y vesiculares. Tesis Lic. México DF Facultad de Ciencias. UNAM. 96.
- Porte, A., & Godoy, R. L. (2008). Chemical composition of Thymus vulgaris L.(thyme) essential oil from the Rio de Janeiro State (Brazil). *Journal of the Serbian Chemical Society*, 73(3).
- Porter, K. L., Klieber, A., & Collins, G. (2003). Chilling injury limits low temperature storage of 'Yuki'Chinese

- cabbage. Postharvest Biology and Technology, 28(1), 153-158.
- Raafat, S. M., Abou-Zaid, M., Aly, A., & Tohamy, M. (2012). Safety control of strawberry fruits gray mold fungus by plant volatile oils. *Zagazig Journal of Agricultural Research*. 39(2): 181-187.
- Salnikow, J., Liao, T.H., Moore, S., & Stein, W. H. (1973). Bovine Pancreatic Deoxyribonuclease A Isolation, Composition, and Amino acid sequences of the tryptic and chymotryptic peptides. *Journal of Biological Chemistry*, 248(4), 1480-1488.
- Scherer, H. W. (2008). Impact of sulfur on N 2 fixation of legumes *Sulfur Assimilation and Abiotic stress in Plants* 43-54.
- Scholz, B. (1962). Atmungsverluste bei Weizen in Abhängigkeit von Temperatur, Lagerzeit und Wassergehalt. *Landtechnische Forschung*, 12(2).
- Senesi, N., Miano, T., & Brunetti, G. (1996). Humic-like substances in organic amendments and effects on native soil humic substances *Humic substances in Terrestrial Ecosystems*. 531-593.
- Serrano, M., Martinez-Romero, D., Castillo, S., Guillén, F., & Valero, D. (2005). The use of natural antifungal compounds improves the beneficial effect of MAP in sweet cherry storage. *Innovative Food Science & Emerging Technologies*, 6(1), 115-123.
- Shen, Y., Sun, Y., Qiao, L., Chen, J., Liu, D., & Ye, X. (2013). Effect of UV-C treatments on phenolic compounds and antioxidant capacity of minimally processed Satsuma mandarin during refrigerated storage. *Postharvest Biology and Technology*, 76, 50-57.
- Tavarini, S., Degl'Innocenti, E., Remorini, D., Massai, R., & Guidi, L. (2008). Antioxidant capacity, ascorbic acid, total phenols and carotenoids changes during harvest and after storage of Hayward kiwifruit. *Food Chemistry*, 107(1), 282-288.
- Tehranifar, A., & Ameri, A. (2012). Effect of humic acid on nutrient uptake and physiological characteristics of Fragaria× ananassa'Camarosa'. Paper presented at the VII International Strawberry Symposium 1049.
- Toor, R. K., & Savage, G. P. (2006). Changes in major antioxidant components of tomatoes during post-harvest storage. *Food Chemistry*, 99(4), 724-727.
- Wang, C. Y., Wang, S. Y., & Chen, C. (2008). Increasing antioxidant activity and reducing decay of blueberries by essential oils. *Journal of Agricultural and Food Chemistry*, 56(10), 3587-3592.
- Watcharasak, S., & Thammasak, T. (2005). Effect of nitrogen and potassium concentration in fertigation on growth and yield of cucumber. *Kamphaengsaen Academic Journal*, *3*, 18-29.

- Yamasaki, H., Uefuji, H., & Sakihama, Y. (1996). Bleaching of the red anthocyanin induced by superoxide radical. *Archives of Biochemistry and Biophysics*, 332(1), 183-186.
- Yang, Y., Wang, J., Xing, Z.E., Dai, Y.Q., & Chen, M. (2011). Identification of phenolics in Chinese toon and analysis of their content changes during storage. *Food Chemistry*, 128(4), 831-838.
- Yuan, G., Sun, B., Yuan, J., & Wang, Q. (2010). Effect of 1-methylcyclopropene on shelf life, visual quality, antioxidant enzymes and health-promoting compounds in broccoli florets. *Food Chemistry*, 118(3), 774-781.
- Žnidarčič, D., & Požrl, T. (2006). Comparative study of quality changes in tomato cv. 'Malike' (*Lycopersicon esculentum* Mill.) whilst stored at different temperatures. *Acta Agriculturae Slovenica*, 87(2), 235-243.