International Journal of Zoology and Applied Biosciences Volume 4, Issue 6, pp: 275-277, 2019 https://doi.org/10.5281/zenodo.3602582

Research Article

PROTEIN PROFILE OF SOMATIC ANTIGENS OF PARAMPHISTOMUM CERVI

*Iram Abdullah¹, Hidayatullah Tak¹ and Showket Ahmad Gani²

¹Department of Zoology, University of Kashmir, Hazratbal, Srinagar 190006, J & K, India ²Department of Clinical Biochemistry, University of Kashmir, Hazratbal, Srinagar 190006, J & K, India

Article History: Received 27th November 2019; Accepted 23th December 2019; Published 31st December 2019

ABSTRACT

Paramphistomum cervi is one of the most common trematode parasites infecting ruminants across the globe. The current study was carried out to identify the somatic antigens present in *P. cervi* using SDS PAGE. The electrophoretic pattern of somatic proteins of *P. cervi* under reducing conditions in 12% SDS revealed the presence of several bands with the molecular weights ranging between 10-150 kDa. The major bands appeared at 10, 18, 23, 40, 50, 75 kDa. The proteins can be subjected to further experimental analysis for detection of their possible role in serodiagnosis or vaccine development.

Keywords: Paramphistomum, Proteins, SDS PAGE, Somatic antigen.

INTRODUCTION

Paramphistomosis is one of the most important diseases causing heavy economic losses to livestock industry annually in terms of low productivity and mortality (Lefevre et al., 2003). The disease is caused by various Paramphistomum species of among Paramphistomum cervi is the most common specie and is found across the globe. Although there is little information regarding the pathogenicity of adult flukes, however, a severe damage to rumenal mucosa was reported in experimentally induced infection in sheep (Rolfe et al., 1994). Diagnosis of Paramphistomosis is primarily based on microscopical analysis of fecal smears and the infection is usually confirmed during the chronic phase of the disease when the parasites have attained sexual maturity and started to produce eggs. However, nowadays a number of serological tests such as ELISA and western blot are increasingly being used to detect the infection especially at prepatent period. Since these tests are based upon the use of certain reactive proteins which can detect the infection at the earliest stage of development. For this purpose proteomic study of parasites involving the identification of antigenic proteins is the need of the hour for early diagnosis of diseases. Studies on the antigenic composition of different sheep trematode parasites have been carried

by several authors (Arora *et al.*, 2010; Eslami *et al.*, 2012; Goreish *et al.*, 2008; Gul *et al.*, 2017; Jaiswal *et al.*, 2018; Latchumikanthan *et al.*, 2012; Meshgi *et al.*, 2008) However, little information is currently available on the antigenic profile of *P. cervi*. This study was carried out to analyze the Somatic protein profile of *P. cervi* using SDS-PAGE. The results would be used in further trials for evaluation of their antigenicity for diagnostic purposes and for developing a protective vaccine.

MATERIAL AND METHODS

Collection of Parasites

Adult parasites were collected from the rumen of sheep slaughtered at local slaughter houses. The parasites were washed several times in phosphate buffer saline (PBS) and used for preparation of soluble protein extract.

Identification of the Parasite

The parasites were fixed in Carnoy's fixative, stained with aceto-alum carmine, dehydrated in ascending grades of ethanol, cleared in xylene and finally mounted in DPX. The identification was carried out as per the standard keys (Yamaguti, 1958).

Preparation of soluble protein extract

For preparation of soluble protein extracts 5 gm of parasite sample were homogenized in 50 ml of homogenizing buffer (50 mM Tris HCl, 1.15% KCl, = 7.4). The homogenate so obtained was centrifuged at 10,000 rpm for 20 minutes in order to remove the debris. The pellet was discarded and the volume of supernant was measured. To the supernatant appropriate quantity of Ammonium sulphate was added and left overnight for precipitation of proteins. Sample was then transferred into dialysis bags and dialyzed in distilled water followed by phosphate buffer. Sample was then again centrifuged at 12,000 rpm for 20 minutes. The supernatant was transferred into dialysis bags and concentrated under fan. The protein sample so obtained was stored till further use.

SDS-PAGE Analysis

Somatic antigens of the parasites were subjected to SDS-PAGE and Gels were stained with Coomassie brilliant blue G-stain. The molecular weights of proteins were determined by comparing their migration distance against that of a known molecular marker.

RESULTS AND DISCUSSION

In the soluble protein fraction of P. cervi, protein concentration was determined by Lowry et al. (1951) and during our study protein concentration was estimated to be about 11mg/ml. The electrophoretic pattern under reducing conditions of 12% SDS-PAGE revealed the presence of several bands in crude somatic protein extract of P. cervi with molecular weights ranging from 10 to 150kDa as shown in Pg 1. The major bands were about 10, 18, 23, 40, 50, and 75 kDa. The crude protein extract of P. cervi when loaded on gel filtration column got resolved into various peaks. In the present study several bands with molecular weights ranging between 10-150 kDa were detected in soluble protein fraction of P. cervi which is in good agreement with study carried out by Meshgi et al. (2009). Similar findings were made by Anuracpreeda et al. (2013) who reported the presence of several protein bands in P. cervi with molecular weights ranging from 10-170 kDa. Jyoti et al. (2014) from her studies on somatic antigens of P. epiclitum also reported the presence of numerous bands having molecular weights in the range of 8.0-169.3 kDa. Our results are in close confirmity with those of Salib et al. (2015) who reported 14 distinct protein bands by SDS PAGE analysis of Paramphistomum somatic antigen with molecular weights ranging from 11.5 kDa to 174 kDa. In current study major bands appeared at 10, 18, 23, 40, 50, and 75 kDa which are in close conformity with those of Salib et al. (2015) who reported the bands with similar molecular weights of 11.5, 19, 25, 46, 52, 72, and 105 kDa in amphistomes.




Figure 1. SDS PAGE profile of somatic proteins of *Paramphistomum cervi*.

CONCLUSION

From the perusual of literature regarding protein profiling of trematode parasite, diversity in somatic protein bands was noticed between various species of *Paramphistomum* which can be attributed to the subsequent ecological and geographical parameters.

ACKNOWLEDGMENT

I express my sincere thanks to Dr. Showket Ahmad Gani, Assistant Professor, Department of Clinical Biochemistry, University of Kashmir, Srinagar, for the facilities provided to carry out this research work.

REFERENCES

Anuracpreeda, P., Poljaroen, J., Chotwiwatthanakun, C., Tinikul, Y., & Sobhon, P. (2013). Antigenic components, isolation and partial characterization of excretion secretion fraction of *Paramphistomum cervi*. *Experimental Parasitology*, *133*(3), 327-333.

Arora, R., Singh, N., Juyal, P., & Ghosh, S. (2010). Immunoaffinity chromatographic analysis for purification of specific diagnostic antigens of *Paramphistomum epiclitum. Journal of Parasitic Diseases*, 34(1), 57-61.

Eslami, A., Taefie-Nasrabadi, N., & Jalosian, F. (2012). Variations to somatic protein profiles of *Dicrocoelium dendriticum* in Iranian ruminants. *Iranian Journal of Veterinary Medicine*, 6(2), 95-98.

Goreish', I. A., Williams, D., Mc Garry, J., Salam, A., & B, E. (2008). Protein Profile of *Fasciola gigantica* Antigens. The *Sudan Journal of Vetenary Research*. 23, 1-9.

Gul, N., Tak, H., Fazilli, K.M., Abdullah, I., & Sofi, T.A. (2017). Prevalence of Fasciola Infection in Slaughtered Animals in Kashmir. International Journal of Veterinary Sciences and Animal Husbandry, 1(2), 30-36.

- Jaiswal, A.K., Shanker, D., Sudan, V., & Singh, A. (2018). Diagnostic potential of low molecular weight excretory secretory proteins of *Paramphistomum epiclitum* for caprine amphistomosis. *Veterinary Parasitology*, 257, 5-9.
- Jyoti, A.; Prasad, and Singh, N.K. (2014). Evaluation of Antibody Response to Various Developmental Stage Specific Somatic Antigens of Paramphistomum epiclitum in Goats. BioMed Research International.1-6
- Latchumikanthan, A., Soundararajan, C., Dhinakarraj, G., & Basith, S.A. (2012). Purification and characterization of excretory/secretory antigens of Fasciola gigantica. *Tamilnadu Journal of Vetenary and Animal Science*, 8(1), 14-18.
- Lefevre, P.C., Blancou, J., & Chermette, R. (2003). Principales maladies infectieuses et parasitaires du bétail: Europe et régions chaudes. *Journal of Agriculture Organization*, 1-3.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., & Randall, R.J.(1951). Protein measurement with the Folin phenolreagent. *Journal of Biological Chemistry*, 193, 265-275.

- Meshgi, B., Eslami, A., & Halajian, A. (2009). Determination of diagnostic antigens in cattle amphistomiasis using western blotting. *Iranian Journal of Parasitology*, 32-37.
- Meshgi, B., Eslami, A., & Hematzadeh, F. (2008). Determination of somatic and excretory-secretory antigens of *Fasciola hepatica* and *Fasciola gigantic* using SDS-page. *Iranian Journal of Veterinary Research*. Shiraz University, 9(1),1-22.
- Rolfe, P., Boray, J., & Collins, G. (1994). Pathology of infection with *Paramphistomum ichika* wai in sheep. *International Journal for Parasitology*, 24(7), 995-1004.
- Salib, F.A., Halium, M.M.A., Mousa, W.M., & Massieh, E. S.A. (2015). Evaluation of indirect ELISA and Western blotting for the diagnosis of amphistomes infection in cattle and buffaloes. *International Journal of Livestock Research*, 5(3), 71-81.
- Yamaguti, S. (1958). Systema helminthum: the digenetic trematodes of vertebrates. Volume 1, Newyork: Interscience publisher.