



Research Article

PROTECTIVE EFFECT OF *PISONIA ALBA* IN ATRAZINE TOXICITY ON BIOCHEMICAL MARKER ENZYMES IN THE LIVER TISSUE OF ALBINO WISTER RAT *RATTUS NORVEGICUS*

¹A.Revathi and ²K.Pugazhendy

¹Department of Zoology, Annamalai University, Annamalai Nagar - 608 002, Tamil Nadu, India

²Department of Zoology, University of Madras Guindy Campus, Chennai, Tamil Nadu, India

Article History: Received 13th August 2021; Accepted 16th November 2021; Published 16th December 2021

ABSTRACT

The goal of this study was to see if *Pisonia alba* might protect the albino wister rat *Rattus norvegicus* from the herbicide atrazine's toxicity effects on AST, ALT, ACP, and ALP. *Rattus norvegicus* were inebriated with a sublethal dose of atrazine (0.25 mg of atrazine) for 28 days in this experiment. When compared to the control, the biochemical manufacturing enzymes in the liver were found to be higher. During the treatment of atrazine-intoxicated rats with *P. alba*, they were returned to a near-normal level (Group III and IV). The outcomes that were noticed were thoroughly explained.

Keywords: Atrazine, *Rattus norvegicus*, *Pisonia alba* and Biochemical makers enzymes.

INTRODUCTION

Atrazine (IUPAC: 6-chloro-N2-ethyl-N4-isopropyl-1, 3, 5-triazine-2, 4-diamine) is an atriazine herbicide commonly used in corn, sugar cane, and sorghum production. It is widely used in non-EU nations and is one of the most widely used pesticides in the United States. The discovery of atrazine in ground water, surface water, and drinking water (WHO, 1993) has piqued interest in its biological consequences. The irrational use of pesticides is one of the main causes of environmental contamination (Mehdi *et al.*, 2005). Pesticides are distinct from other chemical substances in that they are released into the environment on purpose. The liver is a vital organ that helps the body's metabolism of both endogenous and foreign chemicals. The majority of xenobiotics are detoxified in the liver by a series of activation, conjugation, and clearance pathways (Rouimi *et al.*, 2012). ATR has the ability to cause oxidative stress by increasing the concentration of reactive oxygen species and oxidative damage products like lipid peroxides, and thus influencing antioxidant enzyme activity (Abarikwu, 2014).

Enzymes such as ACP, ALP, AST, and ALT have been employed as bioindicators to evaluate pollution exposure (Lavanya *et al.*, 2011). AST and ALT are two of these enzymes that are commonly employed to diagnose tissue damage and act as stress indicators in aquatic biomonitoring (Jung *et al.*, 2003). ALT and AST are commonly measured clinically as part of a diagnostic evaluation of hepatocellular injury to determine liver health (Wang *et al.*, 2012). The aminotransferases or transaminases (AST and ALT) and the phosphatases: total, acid, and alkaline are the major enzymes of clinical significance in liver function (Raju, 2005). *Pisonia alba* is known in Tamil as Leechi kottai kerai (Khare, 2007). *Pisonia alba* is an evergreen glaborous garden tree with minutely puberulous new branches that belongs to the Nyctaginaceae family. It is native to Hawaii and has spread throughout India. *Pisonia alba* leaves are used in alternative medicine as an analgesic, anti-inflammatory, and diuretic (Radha *et al.*, 2008). Hypoglycemic and antifungal agent (Sripathi & Poongothai, 2008; Sunil,

*Corresponding Author: Dr. K. Pugazhendy, Associate Professor, Department of Zoology, University of Madras Guindy Campus, Chennai, Tamil Nadu, India. Email: pugalendy@gmail.com, Mobile: +91 9865225355.

2009). It can be used to cure ulcers, diarrhoea, and snake bites, among other things. The leaves are edible and used to cure wounds, rheumatism, and arthritis (Prabu *et al.*, 2008). Kim and colleagues (Kim *et al.*, 2002). Leaves are also used as a vegetable and salad ingredient, as well as for cattle feed. The various parts of *Pisonia grandis* are widely used in tribal folk medicines and as an Indian traditional medicine for anti-diabetic (Sunil, 2009), anti-inflammatory (Jayakumar *et al.*, 2006; Radha *et al.*, 2008; Thillaivanan *et al.*, 2013), wound healing (Prabu *et al.*, 2008), diuretic (Prabu *et al.*, 2008), wound healing (Elumalai & Prakash, 2012). Natural remedies derived from traditional herbs are used to treat hepatotoxicity, and they are regarded to be effective and safe (Fakurazi *et al.*, 2012; Huang *et al.*, 2010). The use of medicinal plants as a complement to conventional treatment has recently piqued international interest. Extracts from medicinal plants and other natural sources contain a wide range of chemicals with significant biological activity (Watanabe *et al.*, 2001). As a result, an attempt was made to investigate the effects of atrazine given at a sublethal dose for 28 days to Albino Wistar rats (*Rattus norvegicus*), as well as the chelating activities of *Pisonia alba* against atrazine toxicity.

MATERIALS AND METHODS

Experimental animal

The present study used adult male albino Wistar rats (*Rattus norvegicus*) weighing 150-200 g from the Central Animal House, Rajah Muthiah Medical College (Reg No.160/1999/CPCSEA, Proposal number: 1096/2014), Annamalai University. The Ethics Committee on Animal Experiment, Faculty of Science, Annamalai University, Annamalai Nagar, Tamil Nadu, India, accepted the study protocol.

Experimental chemical

The experimental chemical atrazine was bought from Rallis India Limited in Mumbai (TATA Atrataf 50 percent WP).

Preparation of Ethanolic Extract

The dried powder was extracted (*Pisonia alba*) in a Soxhlet apparatus using ethanol at a temperature range of 55 °C to 60 °C. In a vacuum evaporator, the filtrate was evaporated to dryness under decreasing pressure.

Preparation of samples

Using a homogenizer, a 10% (w/v) tissue homogenate was produced in 50 mM Tris HCL (PH 7.4). The homogenate was centrifuged at 10,000 rpm for 10 minutes at 40°C to obtain post mitochondrial supernatant (PMS). The particle was discarded, and the resulting supernatant was dubbed PMS. The homogenate and post mitochondrial supernatant

of rat liver tissue were tested for a variety of biochemical parameters.

Enzymatic Assay

Tissue aspartate aminotransferase was measured using a diagnostic kit based on Reitman and Frankel's approach (1957). Alanine aminotransferase was measured using a diagnostic kit based on Reitman and Frankel's approach (1957). The diagnostic kit was used to calculate tissue alkaline phosphatase using Kind and King's approach (1954). Tissue acid phosphatase was calculated using King's technique and a diagnostic kit (1956).

Experimental design

A total of 24 animals will be divided into 4 groups of 6 in each. Group 1: Control animals, Group 2: atrazine alone (0.25mg/kg bw), Group 3: atrazine (0.25mg/kg bw) + *Pisonia alba* (1 g/kg bw), Group 4: *Pisonia alba* (1 g/kg bw)

Statistically Analyses

The results of this study were expressed as means SE, percentage changes, and statistically assessed using the student t-test (Milton and Tsokos, 1983) to compare the means of treatment and control data, with the results being judged significant at the (P0.05) and (P0.01) levels.

RESULTS AND DISCUSSION

Hence an attempt has been made to investigate the biochemical makers enzymes activities such AST, ALT, ACP and ALP of liver tissue in albino wister rat *Rattus norvegicus* against atrazine exposed group I, atrazine along with *P. alba* (group III) alone *P. alba* alone (group IV) for the period of 1, 7, 14, 21 and 28 days. When compared to the control group I, the activity of aspartate aminotransferase (AST) in the liver of *Rattus norvegicus* after atrazine administration was considerably lower in group II. - 19.814, - 22 were the percent changes. 435, minus 26 187, minus 32. For the periods of 1, 7, 14, 21, and 28, respectively, 951 and -36.150 were used. In the rat treated with atrazine and *Pisonia alba* extract, group III showed a substantial rise in comparison to group II. For periods of 1 to 28 days, the percent changes were 1.422, 2.221, 12.769, 15.983, and 16.824 correspondingly. When a single dose of *Pisonia alba* extract was given to group IV, the AST activity was higher than in group III, and it was close to that of the control group. For periods of 1 to 28 days, the percent changes were -13.045, -14.226, -14.220, -14.213, and -14.214, respectively. At 1% and 5% levels, the reported AST activity in liver tissue for all four groups was statistically significant (Table 1).

When compared to the control group I, the Alanine aminotransferase (ALT) activity in the liver of *Rattus norvegicus* after atrazine administration was considerably lower in group II. For periods of 1, 7, 14, 21, and 28 days, the % changes were -29.394, -35.086, 40.949, -45.400, and -19.940, respectively. In the rat given atrazine and *Pisonia*

alba extract in group III, the level of atrazine was substantially higher than in group II. For the periods of 1 to 28 days, the percent changes were 26.588, 19.695, 5.649, 12.135, and 19.940, respectively. When the extract of *Pisonia alba* was given to group IV alone, the ALT activity was higher than in group III, and it was close to that of the control group. For the periods of 1 to 28 days, the % changes were 24.702, 19.750, 14.797, 9.843, and 9.834, respectively. At 1% and 5% levels, the reported ALT activity in liver tissue for all four groups was statistically significant (Table 2).

When compared to the control group I, the acid phosphatase (ACP) activity in the liver of *Rattus norvegicus* after atrazine administration was reduced in group II. For the periods 1, 7, 14, 21, and 28, the percent changes were -6.130, -18.480, -26.238, -30.663, and -39.585, respectively. In the rat given atrazine and *Pisonia alba* extract in group III, the level of atrazine was substantially higher than in group II. For the period of 1 to 28 days, the percent changes were 1.548, 14.955, 10.645, 9.116, and 19.034 correspondingly. When a single dose of *Pisonia alba* extract was given to group IV, the ACP

activity was higher than in group III, and it was close to that of the control group. For the period of 1 to 28 days, the percent changes were 0.072, 0.120, 0.217, 0.241, and 0.433, respectively. At 1% and 5% levels, the reported ACP activity in liver tissue for all four groups was statistically significant (Table 3). When compared to the control group I, the activity of alkaline phosphatase (ALP) in the liver of *Rattus norvegicus* after atrazine administration was considerably lower in group II. For periods of 1, 7, 14, 21, and 28 days, the % changes were 111.403, 102.631, 101.754, 102.631, and 93.859, respectively. When rats were given atrazine and *Pisonia alba* extract, the concentration of atrazine in group III was substantially higher than in group II. For the periods of 1 to 28 days, the percent changes were -41.908, -37.229, -43.478, -42.857, and -40.271, respectively. When a single extract of *Pisonia alba* was given to group IV, the ALP activity was higher than in group III, and it was close to that of the control group. For the periods of 1 to 28 days, the % changes were -8.771, -8.781, -8.761, -8.791, and -8.753, respectively. At the 1% and 5% levels, the reported ALP activity in liver tissue for all four groups was statistically significant (Table 4).

Table 1. Changes in the Aspartate Amino Transferase (AST) (IU/L) activities in the liver tissue of albino Wistar rat *Rattus norvegicus* administered to atrazine followed by the extract of *Pisonia alba* administered to 28 days.

Groups		Treatment Duration in Days				
		1	7	14	21	28
AST	Group-I Control	84.288 ± 2.363	84.285 ± 2.244	84.280 ± 2.196	84.277 ± 1.963	84.275 ± 1.972
	Group-II atrazine	67.587 ± 2.006**	65.375 ± 1.872**	62.209 ± 1.960**	56.505 ± 2.786**	53.809 ± 2.509**
		-19.814	-22.435	-26.187	-32.951	-36.150
		83.089 ± 2.472**	82.413 ± 7.904*	73.518 ± 2.680**	70.807 ± 2.872**	70.096 ± 2.387**
	Group-III atrazine + <i>Pisonia alba</i>	22.936	26.068	18.179	25.311	30.268
		73.292 ± 2.804*	72.294 ± 2.009**	72.295 ± 2.373**	72.298 ± 2.904**	72.296 ± 2.707**
		-13.045	-14.226	-14.220	-14.213	-14.214
	Group-IV <i>Pisonia alba</i>					

(Values are mean S.E-Mean of six individual observations; and student t-test. Significant at *P<0.05; Significant at ** P<0.01 levels. (+, -), NS-Non-significant.

Table 2. Changes in the Alanine Amino Transferase (ALT) (IU/L) activities in the liver tissue of albino Wistar rat *Rattus norvegicus* administered to atrazine followed by the extract of *Pisonia alba* administered to 28 days.

Groups		Treatment Duration in Days				
		1	7	14	21	28
ALT	Group-I Control	40.473 ± 1.863	40.469 ± 1.096	40.467 ± 2.084	40.464 ± 2.273	40.461 ± 1.896
	Group-II atrazine	28.576 ± 2.086**	26.270 ± 2.112**	23.896 ± 1.544**	22.093 ± 1.709**	21.544 ± 2.305**
		-29.394	-35.086	-40.949	-45.400	-46.753
		20.978 ± 2.226*	21.096 ± 2.315**	22.546 ± 2.096	24.774 ± 1.880**	25.840 ± 1.196**
	Group-III atrazine + <i>P. alba</i>	26.588	19.695	5.649	12.135	19.940

Group-IV	30.475± 1.715**	32.476± 2.804	34.479±2.317	36.481± 2.326	36.482± 2.445
<i>Pisonia alba</i>	24.702	19.750	14.797	9.843	9.834

(Values are mean S.E-Mean of six individual observations; and student t-test. Significant at *P<0.05; Significant at ** P<0.01 levels. (+,-), NS- Non-significant.

Table 3. Changes of Acid Phosphatase (ACP) (μmole/min/mg protein) activities in the liver of albino Wister rat *Rattus norvegicus* administered to atrazine followed by the supplementary feed of *Pisonia alba* administered to 28 days.

Groups	Treatment Duration in Days				
	1	7	14	21	28
Group-I Control	4.127 ± 0.034	4.134 ± 0.036	4.139 ± 0.044	4.145 ± 0.046	4.148 ± 0.028
Group-II atrazine	3.874** ±	3.370** ±	3.053** ±	2.874** ±	2.506** ±
	0.048	0.037	0.039	0.031	0.044
	-6.130	-18.480	-26.238	-30.663	-39.585
Group-III atrazine + <i>P.alba</i>	3.934** ±	3.874** ±	3.378** ±	3.136** ±	2.983** ±
	0.029	0.037	0.052	0.043	0.036
	1.548	14.955	10.645	9.116	19.034
ACP Group-IV <i>Pisonia alba</i>	4.130 ^{NS} ±	4.139 ^{NS} ±	4.148 ^{NS} ±	4.155 ^{NS} ±	4.166 ^{NS} ±
	0.054	0.046	0.037	0.039	0.054
	0.072	0.120	0.217	0.241	0.433

(Values are mean S.E-Mean of six individual observations; and student t-test. Significant at *P<0.05; Significant at ** P<0.01 levels. (+,-), NS- Non-significant.

Table 4. Changes of Alkaline Phosphatase (ALP) (μmole/min/mg protein) activities in the liver of albino Wister rat *Rattus norvegicus* administered to atrazine followed by the extract of *Pisonia alba* administered to 28 days.

Groups	Treatment Duration in Days				
	1	7	14	21	28
Group-I Control	114.908 ±	114.914 ±	114.921 ±	114.927 ±	114.935 ± 3.056
Group-II atrazine	3.004	4.052	3.037	4.061	
	241.208 ±	231.915	231.212	231.007 ±	221.870±
	6.055**	±4.052**	±4.044**	3.057**	4.053**
Group-III atrazine+ <i>P.alba</i>	111.403	102.631	101.754	102.631	93.859
	140.344 ±	145.126 ±	131.877±	132.334 ±	132.154 ±
	3.061**	3.028**	4.039**	3.033**	4.044**
ALP Group-IV <i>Pisonia alba</i>	-41.908	-37.229	-43.478	-42.857	-40.271
	104.915 ±	104.923	104.931±	104.943±	104.957 ±
	4.062*	±4.046*	5.055*	4.045*	3.036*
	-8.71	-8.781	-8.761	-8.791	-8.751

(Values are mean S.E-Mean of six individual observations; and student t-test. Significant at *P<0.05; Significant at ** P<0.01 levels. (+,-), NS Non-significant.

The pervasive pollution of the environment caused by chemical chemicals such as pesticides is a severe problem for all living things, including people 38-40. Various chemical substances that enter animal bodies are transported to and expelled by organs responsible for detoxification, such as the liver and kidney. The calculation of antioxidant potential and hepatocyte damage are valuable tools in the assessment of liver toxicity in the

current study. An experiment with rats revealed that pesticide had oxidative and inflammatory effects on the brain system and livers of mammals. More than 100 human diseases are caused by oxidative stress (Kosecik *et al.*, 2005), and liver diseases are still one of the most important health problems in the world (Huang *et al.*, 2010). Antioxidants obtained from nature combat the oxidative stress caused by several hepatotoxins (Sundararajan *et al.*,

2006). Enzyme markers are more accurate indications of an organ's health. Many environmental contaminants, including as pesticides and metals, can cause the production of reactive oxygen species (ROS) (Gluszczak *et al.*, 2007).

Any ingested toxicant in an animal or human has a significant target organ in the liver. This could be related to the principal site for all chemical metabolisms (Thangapandiyan & Miltonprabu, 2013). These findings imply that ethanolic extract functions as a natural antioxidant and anti-inflammatory medium (Huang *et al.*, 2010). However, the aspartate transaminase (AST) level in the exposed groups' livers fell at first, and then increased after some time. Lysozymes secrete ALP, which is regarded as a key integrity marker of cellular injury. Bromadiolone-treated rats had significantly higher levels of AST, ALT, ALP, LDH, and bilirubin, which is consistent with a recent report. As a result, changes in these enzymes' activity are often related to the severity of cellular damage. Aspartate aminotransferase (AST) and aspartate aminotransferase (ALT) are often evaluated clinically to determine liver health as part of a diagnostic evaluation of hepatocellular damage. In clinical research, aspartate aminotransferase is a good indication of liver injury. AST was discovered to be secreted into the bloodstream during hepatocellular damage. These enzymes seep into the bloodstream from dead or damaged cells (Mansour & Mossa, 2010). Elevated ALT and AST levels in this study hint to active amino acid consumption in energy-generating metabolic pathways such as gluconeogenesis. In order to escape pesticide-induced oxidative stress, it also represents a genetic aberration in production (Bhushan *et al.*, 2013). Enhanced levels of serum enzymes like AST and ALT suggested increased permeability and hepatocyte injury or necrosis. The current findings are consistent with those who found higher hepatic markers. These findings show that atrazine acts as an enzyme inhibitor, impairing hepatic metabolism, at the dosages used in this study. As the concentration of petrol rises, so do the levels of AST, ALT, and ALP. While Hepatotoxicity was observed in male and female rats exposed to Premium Motor Spirit (PMS) blend unleaded gasoline (UG) vapours.

The treated rats' livers had higher levels of alkaline and acid phosphatases. It's been suggested that an increase in ACP levels is caused by liver cell destruction (Vohra & Khera, 2013). Similarly, an increase in the specific activity of ACP in rat tissues only validated the pesticide-induced increase in the dephosphorylation potential within the rat cells. According to there was a corresponding increase in liver enzyme activity. (Revathi *et al.*, 2016; Tamizhazhagan & Pugazhendy, 2017; Tamizhazhagan *et al.*, 2017a, 2017b) *Pisonia alba* supplementation provides anti-oxidant and anti-inflammatory properties against atrazine-induced oxidative stress. It also protects against hepatic dysfunction caused by atrazine and plays a modulatory role in atrazine-induced free radical production. When compared to the control group I, *Rattus norvegicus* showed a recovery response in enzymological activity after 28 days of supplemental group III (atrazine together with *Pisonia*

alba), which was drastically restored from the atrazine exposure (group II).

CONCLUSION

As a result of our research, it is obvious that after supplementing the atrazine-treated rat with *Pisonia alba*, the activity levels of AST, ALT, ACP, and ALP in the test tissue showed a trend toward normalcy. Vitamin A, Vitamin C, thiamine, riboflavin, nicotinic acid (Vitamin B3), alkaloids, proteins, and fats were found in the active ingredient of *P. alba*, as well as alkaloids, proteins, and lipids. Vitamin C is one of four dietary antioxidants, with Vitamin E, Vitamin A precursor-carotene, and Selenium being the others.

ACKNOWLEDGMENT

The authors express sincere thanks to the head of the University of Madras, Guindy Campus for the facilities provided to carry out this research work.

REFERENCES

- Abarikwu, S. O. (2014). Protective effect of quercetin on atrazine-induced oxidative stress in the liver, kidney, brain, and heart of adult wistar rats. *Toxicology International*, 21(2), 148.
- Bhushan, B., Saxena, P. N., & Saxena, N. (2013). Biochemical and histological changes in rat liver caused by cypermethrin and beta-cyfluthrin. *Arhiv za Higijenu rada i Toksikologiju*, 64(1), 57-66.
- Elumalai, A., & Prakash, Y. G. (2012). Evaluation of anti-arthritis activity of ethanolic extract of *Pisonia grandis* R. Br. *Asian Journal of Pharmaceutical Research*, 2(3), 91-93.
- Fakurazi, S., Sharifudin, S. A., & Arulselvan, P. (2012). *Moringa oleifera* hydroethanolic extracts effectively alleviate acetaminophen-induced hepatotoxicity in experimental rats through their antioxidant nature. *Molecules*, 17(7), 8334-8350.
- Gluszczak, L., dos Santos Miron, D., Moraes, B. S., Simões, R. R., Schetinger, M. R. C., Morsch, V. M., & Loro, V. L. (2007). Acute effects of glyphosate herbicide on metabolic and enzymatic parameters of silver catfish (*Rhamdia quelen*). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 146(4), 519-524.
- Huang, B., Ban, X., He, J., Tong, J., Tian, J., & Wang, Y. (2010). Hepatoprotective and antioxidant activity of ethanolic extracts of edible lotus (*Nelumbo nucifera* Gaertn.) leaves. *Food Chemistry*, 120(3), 873-878.
- Jayakumar, A., Rao, K. R., Murthy, C. R., & Norenberg, M. (2006). Glutamine in the mechanism of ammonia-induced astrocyte swelling. *Neurochemistry International*, 48(6-7), 623-628.

- Jung, S. H., Sim, D. S., Park, M. S., Jo, Q., & Kim, Y. (2003). Effects of formalin on haematological and blood chemistry in olive flounder, *Paralichthys olivaceus* (Temminck et Schlegel). *Aquaculture Research*, 34(14), 1269-1275.
- Khare, C. (2007). *Indian Medicinal Plants* Springer-Verlag Berlin: Heidelberg.
- Kim, S.H., Kwon, H.B., Kim, Y.S., Ryu, J.H., Kim, K.S., Ahn, Y., Choi, K.Y. (2002). Isolation and characterization of a Drosophila homologue of mitogen-activated protein kinase phosphatase-3 which has a high substrate specificity towards extracellular-signal-regulated kinase. *Biochemical Journal*, 361(1), 143-151.
- Kosecik, M., Erel, O., Sevinc, E., & Selek, S. (2005). Increased oxidative stress in children exposed to passive smoking. *International Journal of Cardiology*, 100(1), 61-64.
- Lavanya, S., Ramesh, M., Kavitha, C., & Malarvizhi, A. (2011). Hematological, biochemical and ionoregulatory responses of Indian major carp *Catla catla* during chronic sublethal exposure to inorganic arsenic. *Chemosphere*, 82(7), 977-985.
- Mansour, S. A., & Mossa, A.-T. H. (2010). Oxidative damage, biochemical and histopathological alterations in rats exposed to chlorpyrifos and the antioxidant role of zinc. *Pesticide Biochemistry and Physiology*, 96(1), 14-23.
- Mehdi A, A. H., Naser A, A., & Anis A, A. (2005). Survey of pesticides used in Qat cultivation in Dhale'and Yafe'and their adverse effects. *Journal of Biology, Agriculture and Healthcare*, 4(25), 74-78.
- Prabu, D., Nappinnai, M., Ponnudurai, K., & Prabhu, K. (2008). Evaluation of wound-healing potential of *Pisonia grandis* R. Br: A preclinical study in Wistar rats. *The International Journal of Lower Extremity Wounds*, 7(1), 21-27.
- Radha, R., Arokiyaraj, S., Agastian, P., Balaraju, K., Kumar, R. M., & Bula, P. (2008). Phytochemical analysis and anti-inflammatory activity of *Pisonia grandis* R. Br. *Biomedical & Pharmacology Journal*, 1(1), 127.
- Raju, S. (2005). *Illustrated medical biochemistry*: Jaypee Brothers Publishers.
- Revathi, A., Pugazhendy, K., & Jayanthi, C. (2016). Hepatoprotective effects of *Pisonia alba* against atrazine toxicity on antioxidant enzymes activities in the liver tissue of albino Wistar rat *Rattus norvegicus*. *International Journal For Research & Development In Technology*, 5(1), 24-32.
- Rouimi, P., Zucchini-Pascal, N., Dupont, G., Razpotnik, A., Fouché, E., De Sousa, G., & Rahmani, R. (2012). Impacts of low doses of pesticide mixtures on liver cell defence systems. *Toxicology in Vitro*, 26(5), 718-726.
- Sripathi, S. K., & Poongothai, G. (2008). Bioassay-guided fractionation and anti-fungal activity studies on *Pisonia grandis*. *Wounds*, 7(1), 21-27.
- Sundararajan, R., Haja, N. A., Venkatesan, K., Mukherjee, K., Saha, B. P., Bandyopadhyay, A., & Mukherjee, P. K. (2006). *Cytisus scoparius* link-A natural antioxidant. *BMC Complementary and Alternative Medicine*, 6(1), 1-7.
- Sunil, C. (2009). Effect of ethanolic extract of *Pisonia alba* Span. leaves on blood glucose levels and histological changes in tissues of alloxan-induced diabetic rats. *International Journal of Applied Research in Natural Products*, 2(2), 298-304.
- Tamizhazhagan, V., & Pugazhendy, K. (2017). Ethnobotanical and Phytopharmacological review of *Pisonia alba* Span. *Asian Journal of Pharmaceutical Clinical Research*, 10(5), 69-71.
- Tamizhazhagan, V., Pugazhendy, K., Sakthidasan, V., & Jayanthi, C. (2017a). Antioxidant properties of *Pisonia alba* plant leaf extract. *International Journal of Zoology and Applied Biosciences*, 2(6), 311-314.
- Tamizhazhagan, V., Pugazhendy, K., Sakthidasan, V., & Jayanthi, C. (2017b). Preliminary screening of phytochemical evaluation selected plant of *Pisonia alba*. *International Journal of Biology Research*, 2(4), 63-66.
- Thangapandiyan, S., & Miltonprabu, S. (2013). Epigallocatechin gallate effectively ameliorates fluoride-induced oxidative stress and DNA damage in the liver of rats. *Canadian Journal of Physiology and Pharmacology*, 91(7), 528-537.
- Thillaivanan, S., Samraj, K., & Parthiban, P. (2013). A Review on Anti Arthritic Herbs in Siddha Medicine. *International Journal of Pharmaceutical Research*, 5(4), 13.
- Vohra, P., & Khera, K. (2013). Effect of Imidacloprid on Plasma and Tissue Biochemistry of Albino rat. *Indian Journal of Applied Research*, 3, 12.
- Wang, X., Yang, G., Feng, Y., Ren, G., & Han, X. (2012). Optimizing feeding composition and carbon nitrogen ratios for improved methane yield during anaerobic co-digestion of dairy, chicken manure and wheat straw. *Bioresource Technology*, 120, 78-83.
- Watanabe, C. M., Wolfram, S., Ader, P., Rimbach, G., Packer, L., Maguire, J. J., Gohil, K. (2001). The in vivo neuromodulatory effects of the herbal medicine ginkgo biloba. *Proceedings of the National Academy of Sciences*, 98(12), 6577-6580.
- WHO (1993). *Guidelines for drinking-water quality*: World Health Organization.