International Journal of Zoology and Applied Biosciences Volume 8, Issue 1, pp: 8-19, 2023 https://doi.org/10.55126/ijzab.2023.v08.i01.002

ISSN: 2455-9571

http://www.ijzab.com Rishan

Research Article

ANTIOXIDANT STATUS ANTICANCER AND HEPATO PROTECTIVE EFFICACY OF MACROTYLOMA UNIFLORUM SEEDS AN IN-VITRO APPROACH

¹Manikandan Mathaiyan, *Suriyavathana Muthukrishnan, ¹Sobiya Pradeep and ¹Thamaraiselvi Ganesan

Department of Biochemistry, Periyar University, Salem - 636011, Tamil Nadu, India

Article History: Received 19th November 2022; Accepted 26th December 2022; Published 8th January 2023

ABSTRACT

The *Macrotyloma uniflorum* seed (Horse gram) has been underutilized pulse around the world. Nevertheless, it contains most valuable nutritional and pharmacological phytochemicals. Thus, this research was performed to evaluate the antioxidant, anticancer, and hepatoprotective potential of horse gram methanol extract. Interestingly, methanol extract of horse gram demonstrated dose dependent antioxidant, anticancer, and hepatoprotective activities. Since, at increased (100 mg mL⁻¹) concentration the methanolic extract scavenged the DPPH radical up to 93.5% and reduced the ferric (Fe³⁺) into ferrous (Fe²⁺) up to 95%. Similarly, the methanol extract also possess considerable anticancer activities up to 75.64% at 300 mg mL⁻¹ concentration. Furthermore, at increased concentration (300 mg mL⁻¹), the methanol extract effectively protect (hepatoprotective) the HEK293 cells (95.71%), which exposed to 100 mM glucose (hepatotoxicity). The obtained results conclude that the horse gram methanol extract contains pharmacologically valuable phytochemicals and that regular intake of horse gram can be recommended to improve human health.

Keywords: Macrotyloma uniflorum, Antioxidant, Anticancer, Cytotoxicity, Nephroprotectivity.

INTRODUCTION

Regular consumption of phytochemical-enriched cereals, grains, vegetables, and fruits can help to reduce the harm caused by free radicals and that are responsible for a variety of chronic illnesses including cancer, vascular disease, coronary heart disease, macular degeneration, metabolic rheumatism, immunodeficiency syndrome, syndrome, ageing, and so on (Golzarand et al., 2015). The ability of antioxidant components such as flavonoids, tannins, and phenolic acids to serve as efficient antioxidants has received a lot of attention recently (Gulcin, 2020). Grains of crops play a significant role in several regions of the world's as largest traditional diets since they are low in calories and high in protein, soluble fibres, a wide range of phytonutrients, and phyto-constituents (Chivenge et al., 2015). The extracts derived from grains contains nutritional tannins as well as non-tannin polyphenolic compounds is prominent among villagers in certain regions of India (Taylor and Duodu, 2017).

Consumption of these grains have been associated with a lower threat of diabetes, obesity, as well as an inhibiting role in the prevention of cardiovascular disease (Asgary et al., 2018). A nutritional supplement would be any compound, which is a nutrition or an element of such a food that contains health - related benefits, such as disorder prevention and cure (Giampieri et al., 2014). Nutraceuticals include isolated nutrition, dietary supplements, particular diet plans, herbal supplements, processed foods, and processed refreshments (Mondello, 2013). Polyphenolic components of numerous grains have been investigated by several researchers as well as found to already have prospective medicinal benefits, such as antioxidant activity (Saleh et al., 2013). Thus, the research into the significance of non-nutrient components such as phenols, tannins, and flavonoids found in plant products, as natural antioxidants has increased significantly (Hano and Tungmunnithum, 2020). The Macrotyloma uniflorum is widely grown, particularly in waterless areas of various countries

including India (Fuller and Murphy, 2018). Previous research has shown that M. uniflorum seed is a fine source of amino acids, protein, carbohydrates, low lipid, iron and molybdenum (Bhaskar et al., 2019). It also contains slow metabolizable carbohydrate (starch), which has a reduced postprandial blood glucose reaction while ingested by diabetics (Chahota et al., 2020). Furthermore, a solvent extract of horse gram plants has been reported as hypolipidemic activity in lab animals fed a elevated fat diet (Kumar et al., 2013). The extract administered to rabbits with oxidative stress caused by an elevated fat diet resulted in an increase in superoxide dismutase (SOD) as well as catalase (CAT), as well as an increase in lowered glutathione (GSH) accumulation (Jarukamiorn et al., 2016). Such horse gram has been consumed by a massive population in rural areas of southern India as a whole seed sprouts (Chahota et al., 2020). Notwithstanding, consuming horse gram seeds upon having to process including heating accompanied by preparing meals, together with boiled rice, millet, and sorghum is indeed a most common practise in rural Indians (Platel and Srinivasan, 2016). Furthermore, the food preparation alcohol of horse gram seedlings with ingredients is thought to become a prospective cure for the throat infection, fever, common cold, as well as the soup produced from this plant's seedlings was shown to produce heat and assist solubilize renal stones (Tamang et al., 2020). However, as per the author's knowledge, the cytotoxicity, and hepatoprotective efficiency of methanolic extract of horse gram against HepG2 and HEK 293 cells respectively have not been reported. Thus, to understand the possible health benefits of regular consumption of horse gram is critical to enhance the health of common peoples. Accordingly, this study was framed to evaluate the free radicals scavenging (DPPH, ABTS, OH, and FRAP), cytotoxic (HepG2 cells) and hepatoprotective (HEK 293 cells) activity of methanol extract of horse gram through in-vitro approach.

MATERIALS AND METHODS

Extract preparation

The *M. uniflorum* seed was collected from nearby market and pulverized to obtain fine powdered particles for extract preparation. The standard hot plate extraction method was followed for extraction process. Concisely, about 100 g of well pulverized and sieved *M. uniflorum* seed sample was completely dissolved in 100 mL of methanol solvent in a 250 mL conical flask and kept for overnight extraction at $50 \pm 1^{\circ}$ C. The methanol solvent from extract was then allowed to evaporate until completely dried. Well dried sample was dissolved in dimethyl sulfoxide (DMSO) to obtain various dosages (12.5, 25, 50, 75, and 100 mg mL⁻¹) for further studies.

Antioxidant activity potential analysis

DPPH assay

The free radical scavenging efficiency of 12.5 - 100 mg mL⁻¹ concentrations of methanolic extract was investigated

through the DPPH (2, 2-diphenyl-1-picrylhydrazyl) radical scavenging activity by adopting the protocol of Aksoy et al. with slight modifications (Aksoy et al., 2013). Briefly, about 100 μ L of (each) DPPH reagent and various concentrations of test samples were dissolved and left undisturbed for 30-35 min at ambient temperature. Similarly, 100 mg mL⁻¹ concentrations of ascorbic acid (vitamin C) were utilized as a positive control. Then absorbance was measured with 517 nm using a UV-vis. spectrophotometer. The typical equation was applied to determine the percentage of DPPH scavenging and IC₅₀ value was calculated through linear regression study.

Hydroxyl (OH) radical scavenging activity

The hydroxyl radical scavenging potential of various concentrations (12.5 - 100 mg mL⁻¹) of test samples were studied by adopting the modified protocol of Pavithra and Vadivukkarasi (Pavithra and Vadivukkarasi, 2015). Concisely, 0.2 mL (each) of various concentrations of samples was blend individually with following ingredients: phosphate buffer: 0.8, FeCl₃: 0.2 EDTA: 0.2, 2-deoxy-dribose: 0.2 mL and mixed thoroughly. Such reaction mixtures were then kept over the water bath for few min and then added vitamin C and H₂O₂ (each 0.2 mL). After 60 min of incubation at 37 °C, ice cold thiobarbituric acid (1.5 mL) and 25% HCl (1.5 mL) were mixed and then heated at 100 °C for 15 min. Subsequently (after cooled), absorbance of these reaction mixtures (test and 100 mg mL⁻¹ concentration of Catechin as positive control) were recorded at 532 nm. The hydroxyl radical scavenging % and IC₅₀ values were standard formula and linear regression studies respectively.

ABTS assay

The ABTS free radical scavenging potential of different concentrations (12.5 - 100 mg mL $^{-1}$) of methanolic extract was investigated by adopting the protocol of Proestos et al. with slight modifications (Proestos et al., 2013). In brief, about 100 μL of each dosage of sample was added with 100 μL of ABTS reagent individually and kept for 20 min at normal temperature. The 100 mg mL $^{-1}$ concentration of vitamin C was utilized as reference. The absorbance of each reaction mixtures were then read 734 nm and used the standard formula and linear regression to calculate the ABTS radical scavenging percentage and IC $_{50}$ values of test and positive control.

Ferrous reducing antioxidant power activity (FRAP)

The ferrous reducing competence of various dosage of test sample was analyzed by adopting the procedure of Rahman et al. with slight modifications (Rahman et al., 2015). Concisely, 0.25 mL (each) of various concentrations (12.5 - 100 mg mL $^{-1}$) of sample was added with 2.5 mL (each) of 1% K₄[Fe(CN)₆]·3H₂O and 2.5 mL of potassium buffer. Such mixtures were left undisturbed for 20 min at **50** \pm **1°**C, later 0.5 mL of TCA (10%) was added individually to each reaction mix to stop further reduction and then centrifuged at 5000 rpm for 15 min. Subsequently, 1.8 mL

of test sample (supernatant) was blended with distilled water (0.35 mL) and FeCl₃ (0.1%) reagent. The 100 mg mL⁻¹ dosage of quercetin was utilized as reference. The absorbance of these reaction mixtures were recorded at 700 nm. The standard formula and linear regression analyses were used to calculate the ferric reducing percentage and IC₅₀ values of test sample and positive control.

Anticancer activity potential analysis

The anticancer activity potential of methanol extract test sample was studied by MTT assay based cytotoxicity study on Liver Hepatocellular Carcinoma (HepG-2) cell line by following the methodology of Al-Sheddi et al. with slight modification on Minimal Essential Medium (MEM, GIBCO) (Al-Sheddi et al., 2018). Briefly, about 5000 viable HepG-2 cells were seeded on 96 well plate (Triplicates) and then incubated for overnight at 37 \pm 1 °C. After incubation, the attached cells were gently washed with 100 mL of medium (serum-free) and then left starved for 1 h at 37 °C. Subsequently, 100 mL of various concentrations (12.5 - 300 mg mL⁻¹) of test sample were treated with HepG-2 cells and then incubated for overnight at 37 °C. Then the medium was aspirated then and refilled with MTT reagent and again incubated for 4 h at 37 °C in a CO₂ chamber. After incubation, the medium was removed and attached cells were gently rinsed with PBS. Then the cells were suspended in 100 mL of DMSO and then stained with formazan and mixed well. Later, the cell-loaded 96 well plates were recorded at 570 nm. The anticancer drug Melphalan was used as a reference control. The following equation was used to calculate the percentage of cell viability.

Hepatoprotective potential analysis

Hepatoprotective activity potential of test sample was studied by following the methodology of Amalia *et al.*, with slight modification (Amalia *et al.*, 2021). Briefly, about 50 mL of diluted (5000 cells/well) HEK 293 cells was added in 96 well microplate containing DMEM medium and incubated for overnight at 37 °C in a CO₂ (5%) chamber. After overnight incubation, the (supernatant discarded) monolayer cells was rinsed with DMEM. Then, 50 mL of various concentrations (12.5 - 300 mg mL⁻¹) of

methanolic extract were individually added and then incubated for 1 h in a CO_2 (5%) incubator at 37 °C, and then added 50 mL of 100 mM glucose (as nephrotoxic inducer) solution and incubated for 24 h. Subsequently, 10 mL of water-soluble tetrazolium-8 solution was added and incubated at 37 °C in a CO_2 (5%) incubator for 2 h. Then 100 mL of HCl was blended to each well to inhibit the reaction and then the absorbance of was read at 450 nm.

RESULTS AND DISCUSSION

The antioxidant efficiency of methanolic extract sample was investigated against number of free radical, which including DPPH, OH, ABTS, and FRAP through in-vitro approach (Table 1). The horse gram methanol extract possess considerable DPPH free radicals scavenging potential on dose dependent basis. Since, at increased concentration of 100 mg mL⁻¹ the DPPH free radicals was scavenged up to 93.5%, interestingly it was close to the DPPH radicals scavenging potential (98.95%) of ascorbic acid (positive control) (Figure 1). Moreover, the IC₅₀ was calculated as 41.524 mg mL⁻¹ and it almost close to the IC₅₀ of positive control (38.74 mg mL⁻¹). Such result suggests that the horse gram methanol extract may contain significant quantity of bioactive compounds with antioxidant potential. Interestingly, a report stated that the acetone (70%) extract and followed by methanol extract possess excellent free radicals scavenging on DPPH, and ABTS free radicals as well as fine ferric reducing potential at increased concentration (Siddhuraju and Manian, 2007). The bioactive components present in the horse gram methanol extract can act as a H atoms donor and it can interact with DPPH (unstable) radical and convert it as DPPH-H (stable) substance and it can also reduce the toxic effects (Moukette et al., 2015). The elevated quantity of polyphenolic group bioactive components might enhance the antioxidant activity (Karaś et al., 2017). Since, numbers of reports suggests that the antioxidant efficiency of plant is polyphenols determined by the occurrence of (Hajimahmoodi et al., 2010; Zhang et al., 2018). Such polyphenols can prevent the formation and accumulation of unstable free radicals and that are accountable for the cause of number of chronic diseases and oxidative stress (Losada-Barreiro and Bravo-Diaz, 2017). People who consume a polyphenolic-rich nutrition have a lower risk of heart disease and cancer (Cruz et al., 2016).

Table 1. Absorbance of methanol extract on various free radicals (DPPH, OH, ABTS, and FRAP,) scavenging activity.

| Plant extract | Various concentration of methanol extract (mg mL ⁻¹) | | | | | | |
|---------------|--|-----------------|-----------------|-----------------|-----------------|---|--|
| | 12.5 | 25 | 50 | 75 | 100 | Positive control (100 mg mL ⁻¹) | |
| DPPH | 2.47 ± 0.06 | 2.95 ± 0.03 | 3.41 ± 0.07 | 3.69 ± 0.05 | 3.95 ± 0.10 | 4.23 ± 0.09 (Ascorbic acid) | |
| OH | 0.95 ± 0.09 | 1.24 ± 0.05 | 1.58 ± 0.02 | 1.74 ± 0.07 | 1.98 ± 0.04 | 4.18 ± 0.08 (Catechin) | |
| ABTS | 0.85 ± 0.05 | 1.02 ± 0.02 | 1.28 ± 0.04 | 1.52 ± 0.06 | 1.94 ± 0.07 | 4.04 ± 0.03 (Ascorbic acid) | |
| FRAP | 1.94 ± 0.02 | 2.01 ± 0.05 | 2.61 ± 0.03 | 2.89 ± 0.02 | 3.11 ± 0.04 | 4.01± 0.04 (Quercetin) | |

The values mentioned in the table are mean and standard error (±SE) of triplicates

A most reactive oxygen-centered species is the OH (hydroxyl) radical, which causes significant damage to nearby biomolecules (Raedschelders et al., 2012). An increased concentration (100 mg mL⁻¹) of methanolic extract demonstrated moderate OH radicals scavenging potential as 58.89% and interestingly it was comparatively lower than the antioxidant activity (98.95%) of Catechin (Figure 2). Furthermore, the IC₅₀ value was calculated and recognized as poor (76.55 mg mL⁻¹) than positive control (35.61 mg mL⁻¹). Hydroxyl (OH) radicals are established by developing a Fe+3-EDTA premixture along with vitamin c as well as H₂O₂ at pH 7.4, resulting in the degradation of 2-deoxy-d-ribose and the formation of a malondialdehyde like product (Manjari et al., 2018). This hydroxyl radical can be scavenged by sequential electron and proton transmission from the extract's oxidant molecules (Bayat and Fattahi, 2018). According to this, the methanolic extract of M. uniflorum seed may contain lowest quantity of electron and proton donating oxidant molecules (Naji and Devaraj, 2011), hence moderate hydroxyl radicals scavenging activity was observed. Accordingly the phytochemicals such as anthocyanins, kaempfetol, isoflavones, quercetin, myricetin, and flavonol enriched plant extract showed remarkable hydroxyl radicals scavenging activity (Sreerama et al., 2010). The result obtained from this study revealed that the methanolic

extract of M. uniflorum seed possess moderate ABTS radicals scavenging activity even at 100 mg mL⁻¹ concentration. Since, at this elevated concentration, the methanolic extract scavenged the ABTS radical up to 52.65% and it was comparatively lower than the antioxidant activity (95.21%) of ascorbic acid. The IC₅₀ value for ABTS scavenging activity of methanolic extract was calculated as 80.08 mg mL⁻¹ and it found that as least than the IC₅₀ value of positive control (34.27 mg mL⁻¹). This result suggests that, the methanolic extract may have lowest number of electron donating active components that react with ABTS radicals. Since, this ABTS assay is electron (from bioactive compounds) transfer based assay. It was confirmed by shifting the electrons from bioactive compounds and interact with the dark blue color ABTS+ compound and reduced into colorless ABTS substance (Macchioni et al., 2021). Interestingly, Parikh and Patel reported that the flavonoid, and phenolic enriched various pulses as well as split pulses frequently consumed by people around the world possess remarkable health condition (Parikh and Patel, 2018). Since, these phenol and flavonoid enriched diet can reduce the free radicals formation and accumulation in the cells, which resulting improved health condition in humans (Parikh and Patel, 2018; Wojtunik-Kulesza et al., 2016).




Figure 1. DPPH radicals scavenging potential of methanol extract of horse gram. The values mentioned in the figures are mean and standard error $(\pm SE)$ of triplicates.




Figure 2. Hydroxyl (OH) radicals scavenging competence of methanol extract of horse gram. The values mentioned in the figures are mean and standard error (±SE) of triplicates.




Figure 3. ABTS radicals scavenging potential of methanol extract of horse gram. The values mentioned in the figures are mean and standard error (\pm SE) of triplicates.




Figure 4. Ferric reducing potential of methanol extract of horse gram. The values mentioned in the figures are mean and standard error (±SE) of triplicates.




Figure 5. IC_{50} values of methanol extract against various free radicals: DPPH, OH, ABTS, and FRAP. The values mentioned in the figures are mean and standard error (\pm SE) of triplicates.




Figure 6. Anticancer activity of methanol extract against HepG2 cells: Through cytotoxicity study. The values mentioned in the figures are mean and standard error (±SE) of triplicates.




Figure 7. Hepatoprotective activity of methanol extract on glucose induced nephrotoxicity on HEK 293 cells. The values mentioned in the figures are mean and standard error (±SE) of triplicates.

Table 2. Anticancer activity (Cytotoxicity) profile of methanol extract on HepG2 cells studied through MTT assay.

| Sample | Treated Conc. mg mL ⁻¹ | Absorbance at 570nm (ELISA Reader) | Percentage of HepG2 cell viability | Number of viable HepG2 cell counted | Percentage of cytotoxicity |
|---------------------|--------------------------------------|--|--|---|----------------------------|
| | 12.5 | 1.2564 ± 0.1 | 98.48 ±1.0 | 4924 ± 4 | 1.52 ± 0.56 |
| | 25 | 0.9852 ± 0.1 | 84.16 ± 2.6 | 4208 ± 6 | 15.84 ± 1.1 |
| | 50 | 0.5864 ± 0.2 | 68.02 ± 2.9 | 3401 ± 5 | 31.98 ± 2.5 |
| Methanol extract | 100 | 0.5216 ± 0.4 | 61.08±1.5 | 3054 ± 8 | 38.92 ± 1.7 |
| plant sample | 200 | 0.4758 ± 0.3 | 44.08 ± 1.5 | 2204 ± 6 | 55.92 ± 1.3 |
| | 300 | 0.4214 ± 0.6 | 24.36 ± 1.8 | 1218 ± 8 | 75.64 ± 1.5 |
| Melphalan (control) | 35:5:60 v/v | 0.105 ± 0.2 | 2.14 ± 0.6 | 107 ± 2 | 97.86 ± 0.4 |

The values mentioned in the table are mean and standard error (±SE) of triplicates

Table 3. Hepatoprotectivity profile of various concentration of methanol extract against glucose induced nephrotoxicity on HEK 293 cells.

| | Sample | | Treated Conc. mg mL ⁻¹ | Absorbance at 5 nm (ELIS Reader) | U | Number of viable HEK 293 cell counted |
|-----------------------------|---------|--------|-----------------------------------|----------------------------------|------------------|---------------------------------------|
| | | | 12.5 | 0.3782 ± 0.7 | 13.42 ± 0.98 | 783 ± 9 |
| | | | 25 | 0.4136 ± 0.4 | 19.16 ± 1.7 | 958 ± 7 |
| | | | 50 | 0.4628 ± 0.8 | 23.16 ± 1.6 | 1158 ± 5 |
| Methanol | extract | plant | 75 | 0.5114 ± 0.6 | 38.11 ± 2.4 | 1912 ± 8 |
| sample | | | 100 | 0.5753 ± 0.5 | 53.08 ± 2.0 | 2654 ± 5 |
| | | | 200 | 0.9624 ± 0.3 | 76.22 ± 1.4 | 3811 ± 4 |
| | | | 300 | 1.1652 ± 0.7 | 95.71 ± 2.1 | 4788 ± 3 |
| Glucose (Hepatotoxic agent) | | 100 mM | 0.135 ± 0.5 | 3.7 ± 1.3 | 185 ± 2 | |

The values mentioned in the table are mean and standard error (±SE) of triplicates

dosage of 100 mg mL⁻¹, the sample effectively reduced the ferric molecule up to 95.52% and it was almost close to the ferric reducing potential (99.12%) of quercetin (Figure 4). A nominal IC₅₀ value (44.146 mg mL⁻¹) was found for methanol extract to reduce the ferric into ferrous and it was considerably comparable with the IC₅₀ value (42.371 mg mL⁻¹) of quercetin. The figure 5 represent the IC₅₀ values of sample against various form of free radicals, which including DPPH, OH, ABTS, and FRAP. The obtained result suggests that, the methanol extract of M. uniflorum seed possess sufficient volume of ferric reducing antioxidants (Rajagopal et al., 2017). Since, antioxidant molecule converts (through electron transfer reaction) the ferric (Fe³⁺) into ferrous (Fe²⁺) (Zhou et al., 2021). While this ferric (Fe³⁺) salt is reduced to ferrous (Fe²⁺) salt, it produces a blue colour, the intensity of this can be measured using spectrophotometer (Bhatta et al., 2017). The oxidation state of ferric (Fe³⁺) salt is similar to that of the radical cation ABTS⁺ (Li et al., 2012). The anticancer activity potential of test sample was investigated through cytotoxicity study by MTT assay. Interestingly, the horse gram methanol extract demonstrated a dose dependent

cytotoxicity on HepG2 cells. Such cytotoxicity activity against the HepG2 cells was ranged from 1.52 to 75.64% at a dosage of 12.5 – 300 mg mL⁻¹ (Figure 6). Positive control melphalan showed cytotoxicity up to 97.86% against this HepG2 cells at 300 mg mL⁻¹ concentration (Table 2). The IC₅₀ value of methanol extract of horse gram against the HepG2 cells was found as 180.11 mg mL⁻¹ and this value was moderately comparable with the IC₅₀ value (137.23 mg mL⁻¹) of melphalan. A report stated that the butyrate enriched pulses can act as a protective shield against colorectal cancer (Prasad and Singh, 2015). Since, such bioactive components can minimize the menace of malignant related alteration in the cells (Seidel et al., 2017). Horse gram containing soluble and non-soluble bioactive components can aid the proper metabolic and other activities of cells, thus it may obstruct malignant cell formation (Gupta et al., 2019). The bioactive components belongs to the flavonoid group categories have been well recognized as possess considerable free radicals scavenging activity, anticarcinogenic, anti - allergic, as well as gastroprotective characteristics (Xuan and Khanh, 2016).




Figure 1. Cytotoxicity profile of methanol extract on HepG2 cell line.

Cytotoxicity (anticancer) potential of various concentrations of methanol extract of test plant sample against human liver cancer cell line HepG2. (a) Non treated (b) anticancer drug (melphalan) treated (c) 12.5 μ g mL⁻¹ treated (d) 25 μ g mL⁻¹ treated; (e) 50 μ g mL⁻¹ treated; (f) 100 μ g mL⁻¹ treated (g) 200 μ g mL⁻¹ treated; (h) 300 μ g mL⁻¹ treated. Arrow marks indicates cell lysis.

Interestingly, the methanolic extract demonstrated fine ferric reducing potential as on dose dependent basis. Because, at a The horse gram methanol extract effectively protect the HEK 293 cells from 100 mM glucose induced nephrotoxicity. The concentration of methanol extract determined the hepatoprotective percentage. hepatoprotective activity percentage was ranged from 13.42% to 95.71% at the dosage of 12.5 - 300 mg mL⁻¹ (Table 3). Interestingly, the obtained result suggest that, at elevated dosage (300 mg mL⁻¹) the test sample showed remarkable hepatoprotective efficiency (95.71%) on glucose (nephrotoxic) induced HEK293 cells (Figure 7). Interestingly, the Patel and Acharya reported that the horse gram possess significant biomedical potential to treat the urolithiasis and other kidney related urinary diseases (Patel and Acharya, 2020). In general, the crop seeds are enriched with number of pharmaceutically valuable phytochemicals than other plant parts. Such phytochemicals derived bioactive compounds have been reported as possess fine nephroprotective and hepatoprotective activities (Prasad and Singh, 2015). This suggests that the methanolic extract of M. uniflorum seed might possess sufficient quantity of hepatoprotective bioactive compounds. Sharma et al. reported about the number of nutraceutical properties of horse gram (Sharma et al., 2019). As per the author's knowledge, this is the first report about hepatoprotective activity of horse gram on glucose induced nephrotoxicity in HEK293 cells. Another report stated that the *Dolichous biflorus* seeds possess considerable hepatoprotective activity on Ethylene Glycol Induced Urolithiasis (Kant and Singh, 2021).

CONCLUSION

The methanolic extract of M. uniflorum seed demonstrated remarkable free radical scavenging efficiency. The horse gram methanol extract effectively scavenged the DPPH and reduced the ferric into ferrous than OH and ABTS free radicals scavenging at high concentrations. Furthermore, methanolic extract demonstrated anticancer efficiency in a dose-dependent basis on HepG2 cells. Similarly, the horse gram methanol extract has excellent hepatoprotective activity in HEK293 cells against glucose-induced nephrotoxicity. As a result, the results indicate that the methanol extract of horse gram contains pharmaceutically valuable bioactive components. However, more research is needed to conduct in-vivo experiments to assess their pharmacological potential in the near future.

ACKNOWLEDGEMENT

The authors are thankful to the Department of Biochemistry, Periyar University, Salem, Tamil Nadu, India, for providing sufficient laboratory support for successful completion of this research work.

REFERENCES

- Aksoy, L., Kolay, E., Ağılönü, Y., Aslan, Z., & Kargıoğlu, M. (2013). Free radical scavenging activity, total phenolic content, total antioxidant status, and total oxidant status of endemic *Thermopsis turcica*. Saudi Journal of biological Sciences, 20(3), 235-239.
- Al-Sheddi, E. S., Farshori, N. N., Al-Oqail, M. M., Al-Massarani, S. M., Saquib, Q., Wahab, R., ... & Siddiqui, M. A. (2018). Anticancer potential of green synthesized silver nanoparticles using extract of Nepeta deflersiana against human cervical cancer cells (HeLA). Bioinorganic Chemistry and Applications, 2018.
- Amalia, R., Aulifa, D. L., Zain, D. N., Pebiansyah, A., & Levita, J. (2021). The cytotoxicity and nephroprotective activity of the ethanol extracts of angelica keiskei Koidzumi stems and leaves against the NAPQI-induced human embryonic kidney (HEK293) cell line. Evidence-Based Complementary and Alternative Medicine, 2021.
- Asgary, S., Rastqar, A., & Keshvari, M. (2018). Functional food and cardiovascular disease prevention and treatment: a review. *Journal of the American College of Nutrition*, *37*(5), 429-455.
- Bayat, A., & Fattahi, A. (2018). The free radical scavenging activity of lespedezacoumestan toward OH radical: a quantum chemical and computational kinetics study. *Journal of Physical Organic Chemistry*, 31(2), e3755.
- Bhaskar, B., Ananthanarayan, L., & Jamdar, S. N. (2019). Effect of enzymatic hydrolysis on the functional, antioxidant, and angiotensin I-converting enzyme (ACE) inhibitor properties of whole horse gram flour. *Food Science and Biotechnology*, 28(1), 43-52.
- Bhatta, S. R., Bheemireddy, V., Vijaykumar, G., Debnath, S., & Thakur, A. (2017). An efficient molecular tool with ferrocene backbone: discriminating Fe3+ from Fe2+ in aqueous media. *Organometallics*, *36*(11), 2141-2152.
- Chahota, R. K., Thakur, N., & Sharma, R. (2020). Efficient Improvement in an Orphan Legume: Horsegram, Macrotyloma uniflorum (Lam.) Verdi, Using Conventional and Molecular Approaches. In Accelerated Plant Breeding, Volume 3, pp. 369-388.
- Chivenge, P., Mabhaudhi, T., Modi, A. T., & Mafongoya, P. (2015). The potential role of neglected and underutilised crop species as future crops under water scarce conditions in Sub-Saharan Africa. *International Journal of Environmental Research and Public Health*, 12(6), 5685-5711.
- Cruz, A. B., Pitz, H. D. S., Veber, B., Bini, L. A., Maraschin, M., & Zeni, A. L. B. (2016). Assessment of bioactive metabolites and hypolipidemic effect of polyphenolic-rich red cabbage extract. *Pharmaceutical Biology*, *54*(12), 3033-3039.

- Giampieri, F., Alvarez-Suarez, J. M., & Battino, M. (2014). Strawberry and human health: Effects beyond antioxidant activity. *Journal of Agricultural and Food Chemistry*, 62(18), 3867-3876.
- Golzarand, M., Bahadoran, Z., Mirmiran, P., Sadeghian-Sharif, S., & Azizi, F. (2015). Dietary phytochemical index is inversely associated with the occurrence of hypertension in adults: a 3-year follow-up (the Tehran Lipid and Glucose Study). *European Journal of Clinical Nutrition*, 69(3), 392-398.
- Gulcin, İ. (2020). Antioxidants and antioxidant methods: An updated overview. *Archives of toxicology*, 94(3), 651-715.
- Gupta, N., Jangid, A. K., Pooja, D., & Kulhari, H. (2019). Inulin: A novel and stretchy polysaccharide tool for biomedical and nutritional applications. *International Journal of Biological Macromolecules*, *132*, 852-863.
- Hajimahmoodi, M., Faramarzi, M. A., Mohammadi, N., Soltani, N., Oveisi, M. R., & Nafissi-Varcheh, N. (2010). Evaluation of antioxidant properties and total phenolic contents of some strains of microalgae. *Journal of Applied Phycology*, 22(1), 43-50
- Hano, C., & Tungmunnithum, D. (2020). Plant polyphenols, more than just simple natural antioxidants: Oxidative stress, aging and age-related diseases. *Medicines*, 7(5), 26.
- Jarukamjorn, K., Jearapong, N., Pimson, C., & Chatuphonprasert, W. (2016). A high-fat, high-fructose diet induces antioxidant imbalance and increases the risk and progression of nonalcoholic fatty liver disease in mice. *Scientifica*, 2016.
- Kant, R., & Singh, T. G. (2021). Nephroprotective Effect of Functional Beverage of *Dolichous Biflorus* Seeds on Ethylene Glycol Induced Urolithiasis: A Mechanistic Approach.
- Karaś, M., Jakubczyk, A., Szymanowska, U., Złotek, U., & Zielińska, E. (2017). Digestion and bioavailability of bioactive phytochemicals. *International Journal of Food Science & Technology*, 52(2), 291-305.
- Kumar, D. S., Prashanthi, G., Avasarala, H., & Banji, D. (2013). Antihypercholesterolemic effect of *Macrotyloma uniflorum* (Lam.) Verdc (Fabaceae) extract on high-fat diet-induced hypercholesterolemia in Sprague-Dawley rats. *Journal of Dietary Supplements*, 10(2), 116-128.
- Li, X., Lin, J., Gao, Y., Han, W., & Chen, D. (2012). Antioxidant activity and mechanism of Rhizoma Cimicifugae. *Chemistry Central Journal*, *6*(1), 1-10.
- Losada-Barreiro, S., & Bravo-Diaz, C. (2017). Free radicals and polyphenols: The redox chemistry of neurodegenerative diseases. *European Journal of Medicinal Chemistry*, 133, 379-402.

- Macchioni, V., Carbone, K., Cataldo, A., Fraschini, R., & Bellucci, S. (2021). Lactic acid-based deep natural eutectic solvents for the extraction of bioactive metabolites of *Humulus lupulus* L.: Supramolecular organization, phytochemical profiling and biological activity. *Separation and Purification Technology*, 264, 118039.
- Manjari, V., Murugesan, S., Banumathi, V., & Pattarayan, R. (2018). Anticancer Activity of Pancha Paasana Chendhuram on MCF-7 Cells. *International Journal of Nutrition, Pharmacology, Neurological Diseases*, 8(4), 143.
- Mondello, L. (2013). Nutraceuticals and separations. *Analytical and Bioanalytical Chemistry*, 405(13), 4589-4590.
- Moukette, B. M., Pieme, C. A., Njimou, J. R., Biapa, C. P. N., Marco, B., & Ngogang, J. Y. (2015). In vitro antioxidant properties, free radicals scavenging activities of extracts and polyphenol composition of a non-timber forest product used as spice: Monodora myristica. *Biological Research*, 48(1), 1-17.
- Naji, K. M., & Devaraj, V. R. (2011). Antioxidant and other biochemical defense responses of Macrotyloma uniflorum (Lam.) Verdc.(Horse gram) induced by high temperature and salt stress. *Brazilian Journal of Plant Physiology*, 23, 187-195.
- Parikh, B., & Patel, V. H. (2018). Total phenolic content and total antioxidant capacity of common Indian pulses and split pulses. *Journal of Food Science and Technology*, 55(4), 1499-1507.
- Patel, V. B., & Acharya, N. (2020). Effect of *Macrotyloma* uniflorum in ethylene glycol induced urolithiasis in rats. *Heliyon*, 6(6), e04253.
- Pavithra, K., & Vadivukkarasi, S. (2015). Evaluation of free radical scavenging activity of various extracts of leaves from Kedrostis foetidissima (Jacq.) Cogn. *Food Science and Human Wellness*, 4(1), 42-46.
- Platel, K., & Srinivasan, K. (2016). Bioavailability of micronutrients from plant foods: an update. *Critical Reviews in Food science and Nutrition*, 56(10), 1608-1619.
- Prasad, S. K., & Singh, M. K. (2015). Horse gram-an underutilized nutraceutical pulse crop: a review. *Journal of Food science and Technology*, 52(5), 2489-2499.
- Proestos, C., Lytoudi, K., Mavromelanidou, O. K., Zoumpoulakis, P., & Sinanoglou, V. J. (2013). Antioxidant capacity of selected plant extracts and their essential oils. *Antioxidants*, 2(1), 11-22.
- Raedschelders, K., Ansley, D. M., & Chen, D. D. (2012). The cellular and molecular origin of reactive oxygen species generation during myocardial ischemia and reperfusion. *Pharmacology & Therapeutics*, 133(2), 230-255.

- Rahman, M., Islam, M., Biswas, M., & Khurshid Alam, A. H. M. (2015). In vitro antioxidant and free radical scavenging activity of different parts of Tabebuia pallida growing in Bangladesh. *BMC Research notes*, 8(1), 1-9.
- Rajagopal, V., Pushpan, C. K., & Antony, H. (2017). Comparative effect of horse gram and black gram on inflammatory mediators and antioxidant status. *journal* of Food and Drug Analysis, 25(4), 845-853.
- Saleh, A. S., Zhang, Q., Chen, J., & Shen, Q. (2013). Millet grains: nutritional quality, processing, and potential health benefits. *Comprehensive Reviews in Food science and Food Safety*, *12*(3), 281-295.
- Seidel, D. V., Azcárate-Peril, M. A., Chapkin, R. S., & Turner, N. D. (2017, October). Shaping functional gut microbiota using dietary bioactives to reduce colon cancer risk. In *Seminars in Cancer biology* (Vol. 46, pp. 191-204). Academic Press.
- Sharma, N., Bisht, S. S., Gupta, S., Rana, M., & Kumar, A. (2019). Nutraceutical Evaluation of Horse Gram (Macrotyloma uniflorum) Cultivated in High Altitudes of Uttarakhand Himalaya, India. *International Journal Pure Applied Bioscience*, 7(4), 190-202.
- Siddhuraju, P., & Manian, S. (2007). The antioxidant activity and free radical-scavenging capacity of dietary phenolic extracts from horse gram (Macrotyloma uniflorum (Lam.) Verdc.) seeds. *Food chemistry*, 105(3), 950-958.
- Sreerama, Y. N., Sashikala, V. B., & Pratape, V. M. (2010). Variability in the distribution of phenolic compounds in milled fractions of chickpea and horse gram: evaluation of their antioxidant properties. *Journal of Agricultural and Food Chemistry*, 58(14), 8322-8330.
- Suriyavathana, M., Manikandan, M., Janeesha, K. J., Sandhya, M., & Ram, K. A. (2018). Phytochemical screening and antimicrobial activity of aqueous seed extracts of *Macrotyloma uniflorum*. *World Journal Pharm Pharm* Sciences, 7(9), 703-714.
- Suriyavanthana, M., & Punithavanthi, M. (2017). Phytochemical analysis and antioxidant profile of Ocimum sanctum Linn. In 5th International Conference on Emerging Trends in Engineering, Technology, Science and Management, Institution of Electronics and Telecommunications Engineers, Ganganagar, Bengaluru, Karnataka.
- Tamang, J. P., Cotter, P. D., Endo, A., Han, N. S., Kort, R., Liu, S. Q., & Hutkins, R. (2020). Fermented foods in a global age: East meets West. *Comprehensive Reviews in Food Science and Food Safety*, 19(1), 184-217.
- Taylor, J. R., & Duodu, K. G. (2017). Sorghum and millets: Grain-quality characteristics and management of quality requirements. In *Cereal Grains* (pp. 317-351). Woodhead Publishing.

- Wojtunik-Kulesza, K. A., Oniszczuk, A., Oniszczuk, T., & Waksmundzka-Hajnos, M. (2016). The influence of common free radicals and antioxidants on development of Alzheimer's disease. *Biomedicine & Pharmacotherapy*, 78, 39-49.
- Xuan, T. D., & Khanh, T. D. (2016). Chemistry and pharmacology of *Bidens pilosa*: an overview. *Journal of Pharmaceutical Investigation*, 46(2), 91-132.
- Zhang, D. Y., Wan, Y., Hao, J. Y., Hu, R. Z., Chen, C., Yao, X. H., & Li, L. (2018). Evaluation of the alkaloid,
- polyphenols, and antioxidant contents of various mulberry cultivars from different planting areas in eastern China. *Industrial Crops and Products*, 122, 298-307.
- Zhou, H., Zhang, H., He, Y., Huang, B., Zhou, C., Yao, G., & Lai, B. (2021). Critical review of reductant-enhanced peroxide activation processes: Trade-off between accelerated Fe3+/Fe2+ cycle and quenching reactions. *Applied Catalysis B: Environmental*, 286, 119900.