

Research Article

FIRST REPORT ON FISH-BORNE ZONOTIC CESTODE LARVAE OF *DIBOTHRIOCEPHALUS* LÜHE, 1899 IN FRESHWATER FISH, *OREOCHROMIS NILOTICUS* (LINNAEUS, 1758), FROM THE HEMAVATI RIVER, KARNATAKA (INDIA)

Upasana Bhattacharya, Anindita Ghosh and *Anjum N. Rizvi

Zoological Survey of India, New Alipore, Kolkata-700053, West Bengal, India

Article History: Received 7th March 2024; Accepted 28th April 2024; Published 19th May 2024

ABSTRACT

The Broad Tapeworm, *Dibothriocephalus* Lühe, 1899 is a widespread cestode parasite infecting freshwater fishes worldwide and is the causative agent of zoonotic disease Diphyllbothriasis. Freshwater food fish, *Oreochromis niloticus* (Linnaeus, 1758), is of immense commercial importance and consumed all over the world due to its affordability and palatable nature. The present study records the occurrence of *Dibothriocephalus* larvae in the intestine of *Oreochromis niloticus*, collected from the Hemavati River, Hassan district, Karnataka and forms the first report of *Dibothriocephalus* larvae in *Oreochromis niloticus* and also forms the first report from Karnataka, India.

Keywords: *Diphyllbothrium*, Nile tilapia, Proceroid, Zoonosis.

INTRODUCTION

Dibothriocephalus Lühe, 1899 (Syn. *Diphyllbothrium*) infections have been reported throughout the world from fishes, cats, dogs and humans. Fish eating mammals including humans serve as definitive hosts of the parasite whereas aquatic copepods and one or more freshwater fishes serve as intermediate hosts (ANSES, 2012). Four parasite species belonging to Family Diphyllbothriidae are known to infect humans: *Dibothriocephalus latus* (Syn. *Diphyllbothrium latum*), *D. nihonkaiensis* (Syn. *Diphyllbothrium nihonkaiense*, *D. klebnoviskii*), *Adenocephalus pacificus* (Syn. *Diphyllbothrium pacificum*) (Dupouy-Camet, 2004; WORMS). The majority of tilapia are freshwater fish that live in lakes, ponds, rivers, and small streams. They are less frequently seen in brackish water. Tilapia are now the second most significant farmed fish in the world, after carps (Fitzsimmons and Watanabe, 2010).

Oreochromis niloticus (Linnaeus, 1758), commonly known as Nile tilapia has a wide natural distribution ranging from the Nile River basin in the south through the Eastern and Western Rift Valley lakes in East Africa, and through the Lake Chad, Niger, Benue, Volta, Gambia and

Senegal river basins in the west (Trewavas, 1983). Its introduced range has now widened, covering at least 85 countries of tropics, subtropics and temperate environments (Lowe *et al.*, 2012). On a global scale, it is the most important tilapia species in aquaculture, supporting major capture fisheries (Genner *et al.*, 2018). Nile tilapia constitutes the majority of tilapia currently being produced in India (Fitzsimmons and Watanabe, 2010).

Oreochromis niloticus (Linnaeus, 1758), has become extremely popular in fish culture systems across India because of its ease of maintenance and profitable yield in response to the rising demand for fish. It is widely used as an inexpensive and convenient source of animal protein across the nation. The presence of infective *Dibothriocephalus* larvae in *Oreochromis niloticus* is an alarming situation for both public health and commercial viability, given the fish's widespread consumption and availability in fish markets.

MATERIALS AND METHODS

The host samples (*Oreochromis niloticus*) were collected in early October 2023, from Hemavati River (12°49'52" N,

*Corresponding Author: Dr. Anjum N. Rizvi, Zoological Survey of India (ZSI), New Alipore, Kolkata-700053, West Bengal, India, Email: anrizvi@gmail.com.

7603'15" E) flowing through Hassan district in the southern part of Karnataka. Two of the samples were dissected in order to study the presence of helminth parasites. Numerous procercoid larvae were visible on the outer surface of the intestinal wall, of which 10 were collected from each host. The larval samples were initially kept in normal saline and later they were pressed between a slide and a cover slip for better visibility of morphological characters under a microscope. After fixation in FAA (formaldehyde, alcohol, acetic acid), the specimens were kept in 70% alcohol for further processing. Then they were stained using Borax Carmine and dehydrated through graded alcohol series from 70 % to absolute alcohol and finally in rectified Xylene. After cleaning with clove oil,

permanent slides were made using Canada balsam as mounting medium. The specimens were studied under the Microscope Leica DM 1000.

RESULTS AND DISCUSSION

The recovered procercoid larvae of *Dibothriocephalus* were elongated in shape, measuring 550-900 μ in length and about 150-310 μ wide. The earlier stages were shorter, exhibiting cercomer with hooklets (each about 30 μ long) and a distinct cephalic invagination. The later stages were longer with a retracted scolex-like structure forming a ridge at the anterior end (Figure1).



Figure1. *Dibothriocephalus* sp. a-c. Developing stages of procercoid larvae (HL:hooklets, Cr:cercomer, Ci:cephalic invagination, R:ridge).

The introduction of *Oreochromis niloticus* in Indian aquaculture dates back to 1987 (Singh and Lakra, 2006). In India, it has been reported from several states including Andhra Pradesh (Jesintha *et al.*, 2020), Telangana (Laxmappa, 2016), Karnataka (Naik *et al.*, 2013), Tamil Nadu (Sathish *et al.*, 2021), Kerala (Raj *et al.*, 2016), Madhya Pradesh (Johnson *et al.*, 2020), Chhattisgarh

(Raghav and Dixit, 2009), Maharashtra (Gorghate *et al.*, 2021), Gujarat (Boussou *et al.*, 2024), Uttar Pradesh (Singh *et al.*, 2021), Bihar (Gunasekar and Isaac, 2017), West Bengal (Behera *et al.*, 2018) and the North East (Munilkumar and Nandeesh, 2007). Reports are also available from different river systems of the county including the Ganga river (Singh *et al.*, 2014) and its

tributaries (Dwivedi *et al.*, 2015), from the Yamuna river (Alam *et al.*, 2014) and its tributaries (Dwivedi *et al.*, 2017; Mayank and Dwivedi, 2017).

Oreochromis niloticus has been reported as the intermediate host of several helminth parasites from different parts of the world (Ranzani-Paiva *et al.*, 2005, Ashade *et al.*, 2013, Hussein and Joshy, 2015, Hung *et al.*, 2015, Paredes-Trujillo *et al.*, 2016, Sinaré *et al.*, 2016, Arguedas *et al.*, 2017, Dao *et al.*, 2017, Adugna, 2020). However, reports of *Dibothriocephalus* (Syn. *Diphyllobothrium*) infection in *Oreochromis niloticus* are relatively fewer and are available from Nigeria (Edeh and Solomon, 2016, Omeji *et al.*, 2022), Kenya (MjakaKhamis *et al.*, 2017), Ethiopia (Sorsa *et al.*, 2019) and Bangladesh (Aunamika *et al.*, 2021). Although Kumari *et al.* (2021) have reported the presence of *Diphyllobothrium* in Tilapia from India (Madhepura region of the River Kosi), the species has not been mentioned. Therefore, the present study constitutes the second report of *Dibothriocephalus* larvae in Tilapia from India, and the first report of the parasite in *Oreochromis niloticus* from Karnataka, India.

Diphyllobothriasis, a zoonotic disease in mammals is caused by the ingestion of raw or partially cooked fish containing the plerocercoid larvae of *Dibothriocephalus*. Zoonosis is an animal disease communicable to humans by several modes of transmission (Pal, 2005). The life cycle of the parasite involves definitive hosts such as humans and other fish-eating mammals and usually two intermediate hosts including a planktonic crustacean and one or more freshwater fishes. When environmental conditions are favourable, the eggs released in freshwater with the faeces of the definitive host, complete their maturation and then hatch into a ciliated embryo, the coracidium. The coracidium develops into proceroid larvae when ingested by the crustacean host within their body cavity. When a fish consumes the infected crustacean, the parasitic larva develops into the next stage, called plerocercoid. The plerocercoid takes approximately one month to lay the first batch of eggs which are released with faeces (ANSES, 2012).

Diphyllobothriasis is associated with eating raw fish and is endemic to Serbia, Scandinavia, North America, Japan, and Chile (King, 2002). Although there have been numerous reports of Diphyllobothriasis in humans throughout the world, they are comparatively rare from India. It was reported for the first time in India by Pancharatnam *et al.* (1998), followed by Devi *et al.* (2007), Ramana *et al.* (2011), Sahoo and Devi, (2013) and Ashrafulla *et al.* (2017), all from South India. The common symptoms include weakness, dizziness, salt craving, diarrhoea, abdominal discomfort, and reduced vitamin B12 absorption (Baily, 2003). Measures for prevention of infection in humans include educating the population to abstain from consuming raw or undercooked fish, treating the existing carriers of the pathogen to prevent contamination of the environment, cooking fish to 56°C for 5 minutes or freezing it to -10°C for 48 hours to kill the infective plerocercoids (Pal *et al.*, 2014).

CONCLUSION

The popularity of Tilapia as a food fish in India warrants due awareness regarding its potential to act as a reservoir of the infective stages of cestode parasite *Dibothriocephalus*, responsible for causing Diphyllobothriasis in humans. Moreover, cestode parasites of fishes are known to interfere with the metabolic processes of fishes, hindering optimum growth and causing economic losses in the fish industry. Also, cultured species like Nile tilapia are more susceptible to such infections due to overstocking and poor water quality management (El-seify *et al.* 2011). The presence of *Dibothriocephalus*, a zoonotic cestode parasite in food fish *Oreochromis niloticus* requires urgent attention as it has both economic implications and significant health risk.

ACKNOWLEDGEMENT

The authors are grateful to Dr. Dhriti Banerjee, Director, Zoological Survey of India for providing the facilities to conduct the present study.

REFERENCES

- Adewole, S.O, Oguntoyinbo B.V., Agunbiade, R.O. & Ayeni S.K. (2014). Prevalence of *Diphyllobothrium latum* (Fish Tapeworm) in Fish in Ekiti State, Nigeria. *Journal of Environment and Earth Science*, 4(2), 23-25.
- Adugna, M. (2020). The prevalence of fish parasites of Nile tilapia (*Oreochromis niloticus*) in selected fish farms, Amhara Regional State. *Ethiopian Journal of Agricultural Sciences*, 30(3), 119–128.
- Alam, A., Chadha, N. K., Joshi, K. D., Chakraborty, S. K., Banerjee, P., Sawant, P.B., Kumar, T., Srivastava, K., Das, S.C.S., & Sharma, A.P. (2014). Food and Feeding Ecology of the Non-native Nile Tilapia *Oreochromis niloticus* (Linnaeus, 1758) in the River Yamuna. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. doi: 10.1007/s40011-014-0338-3
- ANSES. (2012). Data sheet on foodborne microbial hazards /*Diphyllobothrium latum*. https://www.anses.fr/en/system/files/MIC2012sa0059_FiEN.pdf
- Arguedas, C. D., Ortega, S. C., Martínez, C.S., & Astroza, C.A. (2017). Parasites of Nile Tilapia larvae *Oreochromis niloticus* (Pisces: Cichlidae) in concrete ponds in Guanacaste, Northern Costa Rica. *Cuadernos de Investigación UNED*, Vol. 9(2), 313-319.
- Ashade, O.O., Osineye, O. M. & Kumoye, E.A. (2013). Isolation, Identification and Prevalence of Parasites on *Oreochromis niloticus* from three Selected River Systems. *Journal of Fisheries and Aquatic Science*, 8(1), 115-121
- Ashrafulla, M.R., Lodhi M. M. I., Kuknoor, G. R. & Katakol, S. (2017). A rare case of Diphyllobothriasis

- and Megaloblastic anaemia in a middle-aged lady from South India, *Indian Journal of Microbiology Research*, 4(4), 471-472.
- Aunamika, S.J., Roy, R.C., Islam, Z., Ahmed, N., Rahman, M.K., Biswas, H., Hasan, M.M., & Taluker, M.H. (2021). Prevalence of fish-borne zoonotic parasites and their molecular identification in Bhola district of Bangladesh. *Bangladesh Journal of Fisheries*, 33(2), 283-294.
- Baily, G. (2003). Other Cestode Infections: Intestinal Cestodes, Cysticercosis, Other Larval Cestode Infections. In: Manson's Tropical Diseases (Ed: Cook GC, Zumla AI.). Philadelphia: Saunders/Elsevier, 1593-1596.
- Behera, B.K., Pradhan, P.K., Swaminathan, T.R., Sood, N., Paria, P., Das, A., Verma, D.K., Kumar, R., Yadav, M.K., Dev, A.K., Parida, P.K., Das, B.K., Lal, K.K. & Jena, J.K. (2018). Emergence of Tilapia Lake Virus associated with mortalities of farmed Nile Tilapia *Oreochromis niloticus* (Linnaeus 1758) in India. *Aquaculture*, 484, 168-174.
- Boussou, C.K., Das, S.P., Mohanty, M., Das, G., Verma, D.K., Sahoo, L., Routray, P. & Das, P. (2024). Morphometric and genetic characterization of cultured and wild populations of tilapia, *Oreochromis niloticus* in India. *3 Biotech*, 14, 51.
- Dao, H.T.T., Dermauw, V., Gabriël, S., Suwannatrai, A., Tesana, S., Nguyen, G.T.T. & Dorny, P. (2017). *Opisthorchis viverrini* infection in the snail and fish intermediate hosts in Central Vietnam. *Acta Tropica*, 170(1), 120-125.
- Devi, C.S., Shashikala, Srinivasan, S., Murmu, U.C., Barman, P., Kanungo, R. (2007). A rare case of Diphyllbothriasis from Pondicherry, South India. *Indian Journal of Medical Microbiology*, 25 (2), 152-4.
- Dupouy-Camet, J. & Peduzzi, R. (2004). Current situation of human diphyllbothriasis in Europe. *EurSurveillance*, 9, 31-5.
- Dwivedi, A.C., Tiwari, A. & Mayank, P. (2015). Seasonal determination of heavy metals in muscle, gill and liver tissues of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) from the tributary of the Ganga River, India. *Zoology and Ecology*, 1-6.
- Dwivedi, A.C., Jha, D.N. & Das, S.C.S. (2017). Population Structure of Nile Tilapia, *Oreochromis niloticus* (Linnaeus 1758) from the Ken River, India. *Journal of Scientific Achievements*, 2(5), 23-27.
- Edeh, C. & Solomon, R.J. (2016). Endoparasites of *Oreochromis niloticus* and *Clarias gariepinus* found in Utako flowing gutter. *Direct Research Journal of Agriculture and Food Science*, 4(12), 361-373.
- El-seify, M. A., Zaki, M. S., Desouky, A. R. Y., Abbas, H. H., Hady, O. K. A., & Zaid, A. A. A. (2011). Seasonal Variations and Prevalence of Some External Parasites Affecting Freshwater Fishes Reared at Upper Egypt. *Life Science Journal*, 8(3), 397-400.
- Fitzsimmons, K. & Watanabe, W. (2010). Tilapia (Family: Cichlidae). In: *Finfish Aquaculture Diversification* (Ed: Le François, N.R., Jobling, M., Carter, C. & Blier, P.U.). C.A.B. International, Oxfordshire, UK, 375-397.
- Genner, M.J., Turner, G.F. & Ngatunga, B.P. (2018). A guide to the tilapia fishes of Tanzania, 29 p.
- Gorghate, N.D., Raut, M.B., & Ingale, P.P. (2021). Diversity and Conservation Status of Fish Fauna in Chichtola Lake, Gondia District, Maharashtra, India. *International Journal of Zoological Investigations*, 7, 16-20.
- Gunasekar, A., & Isaac, S.S. (2017). The Biodiversity of fish fauna in Indrapuri Dam Rohtas district, Bihar (India). *International Journal of Fisheries and Aquatic Studies*, 5(2), 416-419.
- Hung, N.M., Dung, D.T., Lan Anh, N.T., Van, P.T., Thanh, B.N., Van Ha, N., Van Hien, H. & Canh, L.X. (2015). Current status of fish-borne zoonotic trematode infections in Gia Vien district, Ninh Binh province, Vietnam. *Parasites and Vectors*, 8(1), 1-10.
- Hussein, D. & Jossy, B. (2015). Prevalence of Internal Parasites of *Oreochromis niloticus* and *Clarias gariepinus* Fish Species in Lake Ziway, Ethiopia. *Journal of Aquaculture Research & Development*, 6. doi: 10.4172/2155-9546.1000308.
- Jesintha, N., Madhavi, K. & Thriveni, K. (2020). Ichthyofaunal diversity of somasila reservoir, Nellore district, Andhra Pradesh. *International Journal of Fauna and Biological Studies*, 7(4), 183-186.
- Johnson, C., Sarkar, U.K., Koushlesh, S.K. & Das, A.K. (2020). Population structure of Nile tilapia and its impact on fisheries of a tropical impacted reservoir, Central India. *Environmental Science and Pollution Research*, 27(6), 1-9.
- King, C.H. (2002). Cestodes (tapeworms). Chapter 285. In: Mandell, Douglas and Bennett's Principles and Practice of Infectious Diseases (Ed: Mandell, G.L., Bennett, J.B., Dolin, R.). Volume 2, 5th edition, Philadelphia: Churchill Livingstone, 2956-2958.
- Kumari, A., Ramashankar, K. & Kumar, A. (2021). Studies on cestode parasites of freshwater fishes in the Madhepura region of the River Kosi. *Biospectra*, 16(2), 185-188.
- Laxmappa, B. (2016). Exotic fish species in aquaculture and aquatic ecosystem in Telangana state, India. *Journal of Aquatic Biology & Fisheries*, 4, 1-7.
- Lowe, M.R., Wu, W., Peterson, M.S., Brown-Peterson, N.J., Slack, W.T. & Schofield, P.J. (2012). Survival, growth and reproduction of non-native Nile tilapia II: fundamental niche projections and invasion potential

- in the northern Gulf of Mexico. *PLoS One* 7(7), e41580
- Mayank, P.& Dwivedi, A.C. (2017).Resource Use Efficiency and Invasive Potential of Non-Native Fish Species, *Oreochromis niloticus* from the Paisuni River, India.*Poultry, Fisheries and Wildlife Sciences*, 5, 1.
- MjakaKhamis, H., Luswet, D., SagweOrina, P.(2017). Prevalence and Diversity of Internal Cestode Parasites Infected Nile Tilapia (*Oreochromis niloticus*) and African Catfish (*Clarias gariepinus*) in Farmers Fresh Water Ponds in Kenya.*American Scientific Research Journal for Engineering, Technology, and Sciences*,34(1), 123-137.
- Munilkumar, S.&Nandeesh, M.C. (2007). Aquaculture practices in Northeast India: Current status and future directions. *Fish Physiology and Biochemistry*, 33:399–412.
- Naik A.S.K., Benakappa S., Somashekara S.R., Anjaneyappa H.N., Kumar, J., Mahesh V., Hulkoti, S.H. &Rajanna, K.B. (2013).Studies on Ichthyofaunal Diversity of Karanja Reservoir, Karnataka, India. *International Research Journal of Environment Sciences*, 2(2), 1-5.
- Omeji, S., Obande, R.A. &Ogaba, S.E. (2022). Seasonal Variation in Parasitic Prevalence of *Oreochromis niloticus*from Upper River Benue, MutumBiu, Taraba State, Nigeria. *Asian Journal of Basic Science & Research*, 4(3), 31-45.
- Pal, M. (2005). Importance of Zoonoses in public health. *Indian Journal of Animal Sciences* 75, 586-591.
- Pal, M., Girmay, G., Woldemariam, T., Dasgupta, R. (2014). Diphyllbothriasis: An emerging Fishborne helminthic zoonosis. *Journal of Natural History*,10(1),10-18.
- Pancharatnam, S., Jacob, E.& Kang, G. (1998). Human diphyllbothriasis: first report from India. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 92,179-180.
- Paredes-Trujillo, A., Velázquez-Abunader, I., Torres-Irriño, E., Romero, D.,& Vidal-Martínez, V.M. (2016). Geographical distribution of protozoan and metazoan parasites of farmed Nile tilapia *Oreochromis niloticus* (L.) (Perciformes: Cichlidae) in Yucatán, México. *Parasites & Vectors*, 9, 66.
- Raghav, M. & Dixit, S. (2009). Diversity of Fishes in a Freshwater Reservoir of Chhattisgarh, India. *Research Journal of Science and Technology* 1(3), 103-107.
- Smrithy, A., Kumar, B. &Abraham, K.M. (2016). Length-Weight and Length-Length Relationship of Invasive Alien Species, *Oreochromis niloticus*from Lentic Habitats of Palakkad District, Kerala. *Journal of Aquatic Biology & Fisheries*, 4, 41-146.
- Sorsa, M., Mamo, G. &Abera, L. (2019). Major fish-borne bacterial and parasitic zoonoses in Ethiopia: A review.*International Journal of Fauna and Biological Studies*, 6(4), 50-58.
- Ramana, K.V., Rao, S., Vinaykumar, M., Krishnappa, M., Reddy,R.,Sarfraz, M.,Kondle, V.,Ratnamani, M.S.,& Rao, R. (2011). Diphyllbothriasis in a nine-year-old child in India: a case report. *Journal of Medical Case Reports*, 5, 332. doi: 10.1186/1752-1947-5-332.
- Ranzani-Paiva, M.J. T., Felizardo, N.N. & Luque, J.L. (2005). Parasitological and hematological analysis of Nile tilapia *Oreochromis niloticus* Linnaeus, 1757 from Guarapiranga Reservoir, São Paulo State, Brazil. *Acta Scientiarum - Biological Sciences*, 27(3), 231-237.
- Sahoo, R.K. & Devi, S. (2013).Acute Diarrhoea in a Child: Fish Tape Worm InfestationWe All May Miss! *International Journal of Recent Trends in Science and Technology*,3(1), 32-33.
- Sathish, S., Chidambaram, P., Uma, A.& Yuvarajan, P. (2021). Prevalence of parasites in tilapia farms and their management practices in Tamil Nadu, India.*Journal of Entomology and Zoology Studies*, 9(2), 678-689.
- Sinaré, Y.,Boungou, M.,Ouédá, A.,Gnémé, A. &Kabré, G.B. (2016).Diversity and seasonal distribution of parasites of *Oreochromis niloticus* in semi-arid reservoirs (West Africa, Burkina Faso). *African Journal of Agricultural Research*, 11(13), 1164-1170.
- Singh, A.K., & Lakra,W.S. (2006). Alien fish species in India: Impact and emerging scenario. *Journal of Ecophysiology and Occupational Health*, 6(3–4), 165–174.
- Singh, A.K., Verma, P., Srivastava, S.C.& Tripathi, M. (2014). Invasion, Biology and impact of feral population of Nile tilapia (*Oreochromis niloticus* Linnaeus, 1757) in the Ganga River (India). *Asia Pacific Journal of Research*, 1(14), 151-162.
- Singh, A.K., Srivastava, S.C.& Verma, P. (2021). MaxEnt distribution modeling for predicting *Oreochromis niloticus* invasion into the Ganga river system, India and conservation concern of native fish biodiversity. *Aquatic Ecosystem Health and Management*, 24, 43–51.
- Trewevas, E. (1983). Tilapiine Fishes of the Genera *Sarotherodon*, *Oreochromis* and *Danakilia*. In: Publication (British Museum (Natural History)), no. 878. Natural History Museum Library, London.
- WORMS, World Register of Marine Species. Available at:<http://www.marinespecies.org> (Accessed in February 2024).

