



ZOOPLANKTON COMMUNITY OF FRESHWATER PONDS IN PURULIA AT WEST BENGAL

¹Chitra J, ²Abantika Majumder, ³Prabakaran, K and ²R. Arthur James

¹Ecology and Conservation Division, Zoological Survey of India, M-Block, New Alipore, Kolkata -700 053, West Bengal, India

²Department of Marine Sciences, Bharathidasan University, Tiruchirapalli – 620124, Tamil Nadu, India

³Department of Applied Science and Humanities, B.P. Poddar Institute of Management and Technology, Kolkata-700052, India

Article History: Received 9th December 2024; Accepted 17th January 2025; Published 31st January 2025

ABSTRACT

Studies were conducted in the zooplankton and water quality from the freshwater ponds of importance on fish culturing at Purulia. Among various groups of zooplanktonic organisms viz., rotifera, copepoda and cladocera were recorded from the study area. The present investigation was conducted in 12 fishery ponds commonly called as bandhs. Water samples were collected and the following physico-chemical parameters viz., pH, Temperature, Dissolved oxygen, Alkalinity, Electrical conductivity, Hardness, Total dissolved solids, dissolved carbon dioxide, nitrate, nitrite, phosphate- total and silicate were analysed. Zooplankton samples were collected to investigate on their diversity and abundance. The results revealed about 14 species of rotifera belongs to 4 families, 2 classes; 12 species of cladocera belongs to 4 families and 3 species of copepod belongs to 1 family, 1 order respectively. 3 species of rotifera namely *B. rubens*, *B. calyciflorus*, *B. ahlstromi* and 1 species *Moina micrura* of cladocera along with nauplii and copepodites stages belonging to cyclopoid copepods. This preliminary finding would act as baseline information for future research in Purulia.

Keywords: Zooplankton, Rotifera, Copepoda, Cladocera, Water quality.

INTRODUCTION

Zooplankton are heterotrophic, microscopic, aquatic organisms and the larval stages of other life forms of invertebrates (Chakraborty and Mallick, 2020). Their population changes in the aquatic environment are often attributed to anthropogenic disturbances and environmental factors. Due to their higher sensitivity to pollution, they were important bioindicators of aquatic ecosystem (Yakubu *et al.*, 2000; Dudgeon *et al.*, 2006; Contreras *et al.*, 2009; Ravikumar *et al.*, 2011 and Krupa *et al.*, 2020). The changes in hydrologic cycle and influences of various environmental factors to the waterbody always implicates the alteration in food web dynamics in the population of plants, microorganisms, invertebrates and vertebrates. Zooplankton constitutes an important food source for many aquatic organisms (Guy, 1992). They act as primary

consumers to several invertebrates and vertebrates (Forro *et al.*, 2008; Chatterjee *et al.*, 2014). The present investigations were carried out on few ponds at Purulia to reveal the zooplankton diversity and the assessment on the water quality in the ponds reflects the suitability to pisciculture.

MATERIALS AND METHODS

Purulia district in West Bengal, forms the lowest step of the Chotanagpur Plateau, north of the Kansabati river in India. About 12 locations were the selected sites for the study viz., Location -S1 (Bhata Bandh - 23° 33239'N; 86° 35456'E); Location – S2 (Bhutgoria Bandh - 23° 31661'N; 86° 37293'E); Location – S3 (Bibir Bandh - 23° 31967'N

*Corresponding Author: Chitra J, Ecology and Conservation Division, Zoological Survey of India, M-Block, New Alipore, Kolkata – 700 053, West Bengal, India Email: chitrajayapalan@gmail.com.

and 86° 35846'E); Location – S4 (Buchaa Bandh - 23° 32693'N; 86°37653'E); Location – S5 (Dulmi Bandh - 23° 31202'N; 86° 36240'E); Location – S6 (Kamala Bandh - 23° 32100'N; 86° 36343'E); Location – S7 (Mahato Bandh - 23° 32549'N; 86°35547'E); Location – S8 (Moyra Bandh - 23° 33350'N; 86° 34702'E); Location – S9 (Poka Bandh - 23° 33188'N; 86° 37131E); Location – S10 (Punya Bandh - 23° 32287'N; 86° 35770'E); Location – S11 (Ram Bandh - 23° 31202'N; 86°36240'E) ; Location – S12 (Saheb Bandh - 23° 33566'N; 86° 36251'E). The collections were carried out during March 2021 (Figure 1).

Surface water samples were collected from above mentioned 12 localities from 7 am to 9 am in poly propylene containers and were analysed water quality parameters viz., pH, Temperature, Dissolved oxygen, Alkalinity, Electrical conductivity, Hardness, Total dissolved solids, dissolved carbon dioxide, nitrate, nitrite, phosphate- total and silicate (APHA, 2005). Zooplankton samples were collected in polypropylene containers of 100ml capacity using plankton net of mesh size 63 microns. 100 litres of surface water were filtered through plankton net and the zooplankton samples were preserved 4 % formaldehyde for the quantitative analysis. The preserved samples were screened using sedgewick rafter counting chamber to investigate the abundance (ind/m³). The plankton samples were sorted groupwise on protozoa, rotifera, copepoda and cladocera were identified following literatures (Sharma, 1998; Michael & Sharma, 1988; Venkataraman, 1999; Roy, 1999). The process of identification and counting were done with the leica DM 1000 microscope.

RESULTS AND DISCUSSION

Rotifera included 5 families Asplanchnidae (1 species; 1 genus), Brachionidae (10 species; 2 genera) Lecanidae (1 species; 1 genera), testudinellidae (1 species; 1 genera), bdelloidea (1 species; 1 genus) of two orders monogononta and bdelloidea respectively. Of these 5 families brachionidae exhibited higher number of species and the species diversity of the rotifers (14 species) were recorded from the fresh water ponds at Purulia. Among Brachionidae, *B. rubens* in S1, S2 and S7; *Brachionus calyciflorus* at S3, *Brachionus ahlstromi* and *B. forficula* at S7 were found to be abundant. The protozoa were represented by amoebozoan with only one species *Centropyxis ecornis* were recorded at S11. The species richness of rotifers was found in location S3 and at S4. Cladocera holds four families viz., Sididae (1 species; 1 genera), Daphnidae (7 species; 3 genera), Moinidae (2 species; 2 genera) and Chydoridae (2 species; 2 genera) along with the cladoceran neonates, 12 species belong to 8

genera, 4 families were recorded. Cladocerans were higher in diversity at S1 and S8. The copepod shown the record of 3 species of cyclopidae. Among copepoda cyclopoid nauplii and copepodites were occurs in all the locations and cyclopoid nauplii dominated in S5 during the study. *Mesocyclops leuckarti* population were higher in S4 and S9. *M. hyalinus* occurs at S12 and *Microcyclops varicans* in S4. In overall the zooplankton diversity were higher in S8 followed in S1 and S4. *Lecane papuana*, *Philodina* sp. of rotifera, *C. ecornis* of amoebzoa, *M. macleayi*, *C. sphaericus*, *A. verrucosa*, *Diaphanosoma Senegal*, *C. laticaudata*, *D. similis* and *S. serrulatus* of cladocera; *M. hyalinus* and *M. varicans* of copepoda were found to be rare in the study (Table. 1).

The zooplankton abundance was investigated in the few groups viz., rotifer, cladocera and copepoda. Among rotifera, the highest peak was observed in S1 by *B. rubens* (2,240 ind/m³) followed by *B. ahlstromi* (1,800 ind/m³) at S7 and *B. forficula* (1,620 ind/m³) at S7 and *B. calyciflorus* (1,380 ind/m³) at S3 and *B. rubens* (2,240 ind/m³) at S1. These mentioned four species were dominating the waters (Fig. 2). Other rotifers were registered moderate abundance. In cladoceran population, cladoceran neonates showed moderate abundance and occurred in all localities. Higher abundance of cladoceran were noticed at S3 by *Moina micrura* (860 ind/m³) followed by *M. macleayi* (760 ind/m³) at S10, cladoceran neonates occupied maximum representation (550 ind/m³) at S1. *Moina micrura* found in majority of the localities S3, S4, S5, S8, S9, S10 and S12 with moderate abundance (Fig. 3). Among the copepods, cyclopoid nauplii (1,140 ind/m³) hold the maximum representation at S4 and *M. leuckarti* (990 ind/m³) at S4 and their abundance were moderate in most of the localities (Figure 4).

Surface water temperature (21° C to 28° C) showed the marked variation where the higher temperature was noticed at S10 and S11 whereas lower temperature at S4. The concentration pH was between 4.8 to 8.5 (Conc.), most of the locations were alkaline in state where the lower level of pH measured at S12. The water transparency revealed the maximum values at S2 followed by S1, S12 and S10. The electrical conductivity also recorded in the surface water showed wide variation between 273 to 716 values. The dissolved oxygen was ranged between 2.04 mg/L to 5.77 mg/L with wide variations whereas lower values were recorded at S12 and S4 and elevated levels were noticed in the values of free carbon dioxide in same station. Alkalinity was almost higher in all the locations except S4 and S12. And the higher values 256 mg/L of hardness recorded at S5. The total dissolved solids were also higher in concentration in water ranging from 0.12 gm/L to 1.6 gm/L. The nutrients viz., phosphate (1.2 to 1.5 mg/L), nitrate (0.02 to 1.8 mg/L), nitrite 0.02 to 5 mg/L) and ammonia (2 to 12.5 mg/L) and silicate (0 to 10mg/L) were recorded with wide fluctuations where higher concentration of nitrate, nitrite, total dissolved solid, hardness at S5 (Figure 5-17).

Table. 1. List of Zooplankton diversity in 12 (S1-S12) localities at Purulia.

	Taxa	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12
	Phylum Rotifera Class Monogonontae Order Ploima Family Asplanchnidae												
1	<i>Asplanchna brightwellii</i> Gosse, 1850	-	-	+	-	+	-	-	-	-	-	-	-
	Family Brachionidae												
2	<i>Brachionus ahlstromi</i> Lindeman, 1939	-	-	+	-	-	+	++ +	+	-	-	+	+
3	<i>B.angularis</i> Gosse, 1851	+	+		+	+	-	-	-	-	-	+	-
4	<i>B.bidentatus</i> Anderson, 1889	-	-	+	-	-	-	-	-	-	-	-	-
5	<i>B. calyciflorus f.dorcas</i> Gosse, 1851	-	-	-	-	+	-	++	-	-	-	-	-
6	<i>B. calyciflorus</i> Pallas 1766	-	-	++ +	-	+	++	+	++	-	-	-	-
7	<i>B. diversicornis</i> Daday, 1883	-	-	-	-	-	++	-	-	-	+	-	-
8	<i>B.forficula</i> Wierzejski, 1891	-	-	-	-	-	-	++ +	+	-	-	-	+
9	<i>B. quadridentatus</i> Hermann 1783	+	-	-	+	-	-	-	-	-	-	+	-
10	<i>B. rubens</i> Ehrenberg, 1838	++ +	++ +	+	+	-	+	++ +	-	+	++	-	+
11	<i>Keratella tropica</i> Apstein, 1907	-	-	+	+	-	-	++	+	-	-	-	-
	Family Lecanidae												
12	<i>Lecane papuana</i> Murray, 1913	-	-	-	-	-	-	-	-	-	-	+	-
	Family Testudinellidae												
13	<i>Testudinella parva</i> Ternetz, 1892	-	-	-	+	-	-	-	-	-	+	-	-
14	Cyst	+	-	-	-	-	-	-	-	-	-	-	-
	Class Bdelloidea												
15	Phyllodina sp	-	-	-	+	-	-	-	-	-	-	-	-
16	Phylum Protozoa – Amoebozoa Family Centropyxidae												
17	<i>Centropyxis ecornis</i> (Ehrenberg, 1841) Leidy, 1879	-	-	-	-	-	-	-	-	-	-	+	-
	Cladocera Order Ctenopoda Family Sididae												
18	<i>Diaphanosoma senegal</i> Gauthier, 1951	+	-	-	-	-	-	-	-	-	-	-	-
	Order Anomopoda												

	Family Daphnidae												
19	<i>Ceriodaphnia cornuta</i> (Sars,1885)	++	++ +	-	-	-	-	+	+	-	-	-	-
20	<i>C. laticaudata</i> Muller, 1867	-	-	-	-	-	-	-	+	-	-	-	-
21	<i>C. quadrangula</i> O F Muller, 1785	+	-	-	-	-	-	-	-	-	-	-	-
22	<i>C. reticulata</i> Jurine, 1820	+	-	-	+	-	-	-	+	-	-	-	+
23	<i>Daphnia similis</i> Claus 1876	+	+	-	-	+	-	-	-	-	-	-	-
24	<i>Simocephalus serrulatus</i> (Koch, 1841)	-	-	-	-	-	-	-	++	-	-	-	-
25	<i>S.expinosus</i> (Koch, 1841)	-	-	-	-	-	-	-	-	-	-	+	+
	Family Moinidae												
26	<i>Moina micrura</i> (Kurz, 1874)	-	-	++ +	+	+	-	-	+	+	+	-	+
27	<i>Moinodaphnia macleayi</i> King, 1853	-	-	-	-	-	-	-	-	-	++	-	-
	Family Chydoridae												
28	<i>Alona verrucosa</i> Sars, 1901	-	-	-	-	-	-	-	+	-	-	-	-
29	<i>Chydorus sphaericus</i> (O.F. Muller, 1776)	-	-	-	-	-	-	-	-	-	-	+	-
30	Cladocera neonates	+	+	-	+	+	-	-	+	-	-	-	+
	Class Copepoda Order Cyclopoida Family Cyclopidae Sub family: Cyclopinae												
31	Cyclopoid copepodites	++	-	+	+	+	-	-	+	++	+	+	+
32	Cyclopoid nauplii	++	+	+	++ +	+	++	+	++	+	++	+	+
33	<i>Mesocyclops hyalinus</i> (Rehberg, 1880)	-	-	-	-	-	-	-	-	-	-	-	+
34	<i>Mesocyclops leuckarti</i> (Claus,1857)	+	+	-	++	+	-	-	+	++	+	-	-
35	<i>Microcyclops varicans</i> (Sars G.O., 1863)	-	-	-	+	-	-	-	-	-	-	-	-




Figure 1. Study Area in Purulia district at West Bengal.




Figure 2. Rotifer abundance in the freshwater ponds at Purulia.




Figure 3. Cladocera abundance in the freshwater ponds at Purulia.




Figure 4. Copepoda abundance in the freshwater ponds at Purulia.




Figure 5. Temperature




Figure 6. pH




Figure 7. Electrical conductivity.



Figure 8. Dissolved Oxygen.




Figure 9. Free carbon dioxide




Figure 10. Alkalinity




Figure 11. Hardness




Figure 12. Total Dissolved Solids




Figure 13. Phosphate




Figure 14. Nitrate




Figure 15. Nitrite




Figure 16. Ammonia




Figure 17. Silicate

The zooplankton and water quality of 12 ponds at Purulia comprised of rotifera, copepoda, cladocera and protozoans. The fish cultured ponds (study sites) were with noticeable variations in nutrient loading with higher concentrations of Silicate (S1), ammonia (S1 and S7) nitrate, nitrate and phosphate, total dissolved solids (S9 and S12). Similar studies with fluctuating levels of nutrients in ponds of West Bengal (Ghosh & Mandal, 2020; Chattopadhyay & Panda 2022). Reports on higher organic load indicates the nutrient availability and values of the dissolved solid interlink with the higher level of pH, alkalinity and hardness level including electrical conductivity (Lin, 1986). And also, the higher concentration of nutrients may be due to external source of wash offs, detergent, pesticides, fisheries etc., substantiates the zooplankton value of potential survivorship in fluctuating environment (Bhakta *et al.*, 2015; Datta, 1999).

In the present study, 12 localities showed distinct differences in the diversity and species wise abundance. The species richness was remarkably higher in S1, S4 and S8 whereas the diversity showed good registry of rotifers, copepods and cladoceran community among zooplankton. The population abundance was also noticed in rotifera copepoda and cladocera. Similarly, Pradhan & Bandyopadhyay, (2018) reported the zooplankton diversity and water quality parameters from few water bodies at Purulia during the studies in the year 2014-2015 revealed few groups of zooplankton viz., rotifer, copepod, protozoa, ostracoda and cladocera at the level of genera. Chatterjee *et al.* 2014 stated from the three sites at Saheb bandh during 2007 – 2012, 10 species of rotifers, 2 species of cyclopoid copepods 5 species of cladocera and 3 species of ostracoda were recorded at Saheb Bandh at Purulia. Similar to the present observations, Brachionus species were reported in large population in the ponds at Purulia. The species diversity, zooplankton population abundance fluctuations depend on the prey, climatic factors, macrophytes, anthropogenic influences, physico chemical factors and seasonal impacts as stated in several studies (Segers, 2003).

CONCLUSION

Ponds/Bandhs are utilized in fisheries widely in rural areas where the water quality and zooplankton are the factors influencing major factor. In present scenario, most of the waterbodies are polluted due to source of factors through algal blooming, pesticides, fertilizers, industrial runoff, climatic factors and human activities. Majority of the studies on zooplankton observation in terms of diversity and population were still unclear with meager identification details upto species level and the gap areas in monitoring freshwater systems. Freshwater systems are to be conserved by the public due to human livelihood depends on their huge resources. The zooplankton available in freshwaters acts as indicators of pollution as well their successful survivorship in extreme fluctuating climate conditions. Long term monitoring on freshwaters seasonally will impact fresh water conservation.

ACKNOWLEDGEMENT

Author expresses sincere thanks to the Director, Zoological Survey of India, M Block, New Alipore, Kolkata -700 053 for the lab facilities.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

ETHICS APPROVAL

Not applicable

REFERENCES

- APHA. (2005). Standard Methods for the Examination of Water and Wastewater. American Public Health Association/American Water Works Association/Water Environment Federation, Washington DC. 21st Edition.
- Bhakta, J.N., Rana, S., Lahiri, S., Jana, B.B. and Ibrahim, S. (2016). Nutrients profile and fish growth of some fish culture ponds. *International Journal of Environment Technology Science*, 1, 1-7.
- Chakraborty, S. and Mallick, P.H. (2020). Freshwater Cladoceran (Cladocera: Branchiopoda) Diversity of Lateritic Rarh Belt of West Bengal, India: A Review. *Advances in Zoology and Botany*, 8, 188-198
- Chatterjee, N. Mukherjee, M., Bhattacharjee, B. (2014). Abundance and diversity of zooplankton and its seasonal variation in the water of Sahebbandh, Purulia, India: A quantitative Study. *The International Journal of Science and Technology*, 2 (12), 22-28.
- Chatterjee, N., Mukherjee, M. and Bhattacharjee, B. (2014). Abundance and Diversity of Zooplankton and Its Seasonal Variation in the Water of Saheb bandh, Purulia, India: A Quantitative Study. *The International Journal of Science and Technology*, 2, 22-28
- Chattopadhyay, D and Panda, S. (2021). Diversity and abundance of zooplankton at Saheb Bandh, Purulia, West Bengal”, *International Journal of Scientific Research in Biological Sciences*, 8 (5), 32 -34.
- Chattopadhyay, D and Panda, S. (2022). Establishment of significant correlation between seasonal Physicochemical parameters and zooplankton diversity in Saheb Bandh, at Purulia, West Bengal. *International Journal of Scientific Research in Biological Sciences*, 9 (2), 17 -23.
- Contreras, J.J., S.S. S. Sarma, M. Merino Ibarra and Nadini, S. (2009). Seasonal changes in the rotifer (Rotifera) diversity from a tropical high altitude reservoir (Valle de bravo, Mexico). *Journal of Environmental Biology*, 30, 191-195.
- Datta, S. (1999). Diversity pattern of tropical pond ecosystem: Nutrient dynamics and functional relationships. PhD Thesis, Aquaculture and Applied

- Limnology Research Unit, Department of Zoology, University of Kalyani, India.
- Forro, L., Korovchinsky, N.M., Kotov, A.A., Petrusek, A. (2009). Global diversity of cladocerans (Crustacea) in freshwater (Review). *Hydrobiologia*, 595 (1), 177-184.
- Guy, D. (1992). The ecology of the fish pond ecosystem with special reference to Africa. Pergamon Press. 220 – 230.
- Krupa, E., Romanova, S., Berkinbaev, G., Yakovleva N., Sadvakasov, E. (2020). Zooplankton as indicator of the Ecological State of Protected Aquatic Ecosystems (Lake Borovoe, Burabay National Nature Park, Northern Kazakhstan). *Water*, 12(9): 2580.
- Lin, C.K. (1986). Acidification and Reclamation of acid sulphate soil fish ponds in Thailand in *Proc. of the First Asian Fisheries Forum*, McLean. Dizon, L.B. and L.V. Hosjhilos, eds. *Asian Fisheries Society*, manila. 71.
- Michael R.G. and B.K. Sharma. (1988). Indian Cladocera (Crustacea: Branchiopoda: Cladocera) Fauna of India and Adjacent countries 263pp. (Edited by Director, Zoological Survey of India).
- Pradhan, A. and Bandyopadhyay, N. (2018). Seasonal variation of Water quality parameters and Zooplankton diversity in waterbodies of Purulia District, West Bengal, India. *International Journal of Zoology and Applied Biosciences.*, 3 (3), 356-461.
- Ravikumar, P., Mehmood, M.A., Somasekhar, R.K. (2013). Water quality index to determine the surface water quality of Sankey tank and Mallathahalli lake, Bangalore urban district, Karnataka, India. *Applied Water Science*, 3 (1), 247-261.
- Roy, T. (1999). Crustacea: Copepoda: Calanoida and Cyclopoida. State Fauna series 3: *Fauna of West Bengal*. Part 10: 251-284. (Edited by Director, Zoological Survey of India).
- Segers, H. (2003). A biogeographical analysis of rotifers of the genus *Trichocerca* Lamarck, 1801 (Trichocercidae, Monogononta, Rotifera) with notes on taxonomy. *Hydrobiologia.*, 500 (1-3), 103 -114.
- Sharma, B.K. (1998). Freshwater Rotifers (Rotifera: Eurotatoria). State Fauna series 3: *Fauna of West Bengal*, Part 11, 341-461. (Edited by Director, Zoological Survey of India).
- Venkataraman, K. (1999). The Freshwater Cladocera, State Fauna series 3: *Fauna of West Bengal*. Part 10, 251-284. (Edited by Director, Zoological Survey of India).
- Yakubu, A.F. Sikoki, F.D., Abowei, J.F.N. and Hart, S.A. (2000). A comparative study of phytoplankton communities of some rivers, creeks and borrow pits in the Niger Delta Area. *Journal of Applied Science, Environment and Management*, 4 (2), 41-46.