

ISSN: 2455-9571

Research Article

# NEW STATE RECORDS OF PYRRHOCOROIDEA (INSECTA: HEMIPTERA: HETEROPTERA) FROM DALMA WILDLIFE SANCTUARY, JHARKHAND, INDIA

\*Rakhi Das, P.C. Saha, Susmita Khanra and Swetapadma Dash

Zoological Survey of India, M-Block, New Alipore, Kolkata 700053, West Bengal, India.

Article History: Received 21st January 2025; Accepted 24th February 2025; Published 31st March 2025

# **ABSTRACT**

The species diversity of superfamily Pyrrhocoroidea (Hemiptera) from Dalma WLS, Jharkhand, India was studied. Eleven species belong to six genera within the families Pyrrhocoridae and Largidae under the superfamily Pyrrhocoroidea were identified, including new distributional records for ten species from Jharkhand. Distributional data for all listed species from Dalma Wildlife Sanctuary, Jharkhand, India, as well as from other regions across India and abroad, have been provided.

**Keywords:** Largidae, New distributional records, Pyrrhocoridae, Species diversity.

# INTRODUCTION

Pyrrhocoroidea is a relatively small superfamily within the order Hemiptera, specifically classified under sub order Heteroptera under infra-order Pentatomomorpha. This superfamily consists of 663 species belonging to 69 genera pertaining to two families worldwide. These two families are Largidae, which includes 213 species under 23 genera, and Pyrrhocoridae, which comprises 450 species under 46 genera (Hemala-Kment, 2015). Generally, these bugs are red in colour and commonly known as "Red Cotton Bugs". Largidae can be distinguished from Pyrrhocoridae by the seventh abdominal sternum of female split medially whereas the seventh abdominal sternum entire in female in case of Pyrrhocoridae. The Indian fauna of Largidae consists of 18 species under 5 genera and Pyrrhocoridae consists of 49 species under 13 genera (Praveen K. et.al.,2024). This paper deals with 11 species under 6 genera belonging to 2 families of the superfamily Pyrrhocoroidea. All of the 11 species are reported for the first time from the Dalma Wildlife Sanctuary and all except Dysdercus koenigii (Fabr,1775) are reported for the first time from the state of Jharkhand.

Dalma Wildlife Sanctuary is a prominent wildlife sanctuary located around the Dalma Hills that is approximately 10 km from Jamshedpur in Jharkhand, India.

The sanctuary's forests, categorized as "Dry peninsular Sal" and "Northern Dry Mixed Deciduous Forest," that cover 193 km² in the East Singhbhum and Saraikela-Kharsawan districts. The sanctuary is safe home for several animals including wild elephants, pangolin, sloth bear, Indian grey wolf, snakes like Bamboo Pit Viper and copper headed trinket snakes, birds like crested serpent Eagle, plum headed parakeets, paradise flycatchers. Although numerous studies have been conducted on various animal groups in this Sanctuary and across the state of Jharkhand, no research has yet focused on the Hemipteran diversity within this Sanctuary. This study specifically investigated the species diversity of the superfamily Pyrrhocoroidea within the sanctuary.

#### MATERIAL AND METHODS

During the survey of the Dalma WLS, Jharkhand by the team of Zoological Survey of India Kolkata, altogether collected Pyrrhocoroid bugs by handpicking, net trap and light tarp methods. Then the specimens were pinned, dried and identified with the help of available literature. For species confirmation, genitalia of certain specimens were dissected following the method described by Ahmad & Qadri (2009). The process began by removing the label from the pinned dry specimen and immersing it in boiling water for 4-5 minutes. Once softened, the specimen was removed from the pin. Under a microscope, the genital capsule (pygophore) was carefully extracted using fine watchmaker's forceps (size 5 or finer). The genital capsule was then placed in a 10% KOH solution and heated to 40°C for 5-10 minutes in a cavity block. Afterward, the internal contents were cleared by thoroughly washing the capsule in distilled water 2-3 times. The parameres and aedeagus were then carefully detached from the genital capsule using fine forceps. Following dissection, the parameres and aedeagus

were preserved in 70% alcohol, stored in an Eppendorf tube, and placed beneath the corresponding specimen.

# RESULTS AND DISCUSSION

From the present study area, 15 examples were collected and identified as 11 species under six genera of superfamily Pyrrhocoroidea from Dalma. All the species except *Dysdercus koenigii* (Fabr,1775) were found to be new report from the state of Jharkhand. Distributional data provided for all the reported species wherever available.




Figure 1. Macrocheraia grandis (Gray,1832), Dorsal view; Fig.2 Iphita limbata (Stal,1870), Dorsal view; Fig.3 Physopelta schlansbuschi (Fabricius,1787), Dorsal view; Fig.4 Physopelta gutta (Burmister, 1834), Dorsal view; Fig.5 Dysdercus evanescens (Dist,1902), Dorsal view; Fig.6 Dysdercus olivaceus (Fabr,1798), Dorsal view; Fig.7 Dysdercus koenigii (Fabr,1775), Dorsal view; Fig.8 Antilochus russus (Stal,1863), Dorsal view; Fig.9 Antilochus coquebertii (Fabr,1803), Dorsal view; Fig.10 Euscopus indicorus (Walk,1872), Dorsal view; Fig.11 Euscopus albatus (Dist,1909), Dorsal view.

#### Systematic account

Superfamily: Pyrrhocoroidea

Key to the Family under the Superfamily Pyrrhocoroidea

- 1. Sixth visible(seventh actual) ventral segment in female always cleft at base.....Largidae

Family I. Largidae

Subfamily: Physopeltinae

Key to the Tribe under the Subfamily Physopeltinae

- 1. Antennae in male very long. First joint of antennae longer than head and pronotum together.....Lohitini

Tribe: Lohitini

Genus: Macrocheraia Guérin-Méneville, 1835

1. Macrocheraia grandis (Gray, 1832) (Fig.1)

1832.Lohita grandis, Gray (Lygreus), in Griff. An. King. xv; Ins. ii, p. 242

1840. Macrocerrea longicornis, Spin. Ess. p. 177

1870. Var. longissima, Stal, En. Hern. i, p. 98

1882. sumatrana, Dz'st. Ent. Month. Mag. xix, p. 158

Lohitini is monotypic includes only the species called *Macrocheraia grandis* (Gray,1832)

Material Examined: 1ex, Forest Guest House, Dalma WLS, East Singhbhum District, Jharkhand, 22.912579N, 86.143106E, 8.vii. 2021, Coll: Chitra. J; 1 ex., Makula Kocha, Dalma Wls, Ranchi, Jharkhand, 24.91187N, 86.13818E, 09.vii. 2021, coll: Chitra J. & Party.

Distribution: India: Maharashtra, Andhra Pradesh, Assam, Uttarakhand, West Bengal, Meghalaya, Uttarpradesh, Tamil Nadu.

Elsewhere: Myanmar, Sumatra, Philippines.

Tribe: Physopeltini

Key to the genus of the Tribe Physopeltinae

Genus: Iphita (Stal, 1870)

2. Iphita limbata (Stal, 1870) (Fig. 2)

1870. Iphita limbata Stal, En.Hem., 1:99.

1902. Iphita limbata Distant, Fauna Brit. India, Rhynchota, 2: 96.

Material Examined: 1ex, Forest Guest House, Dalma WLS, East Singhbhum District, Jharkhand, 22.912579N, 86.143106E, 8.vii. 2021, Coll: Chitra J. & Party.

Distribution: India: Maharashtra, Assam,Andhra Pradesh, Bihar, Karnataka, Kerala, Meghalaya, Tamil Nadu, Uttarakhand, West Bengal.

Elsewhere: China, Bangladesh, Cambodia, Indonesia, Malaysia, Myanmar, Laos, Nepal, Vietnam, Thailand.

Genus: *Physopelta* (Amy. & Serv.,1865) Key to the species of the genus *Physopelta* 

- 1 Two large transverse spot present at anterior margin of the posterior lobe of pronotum, body sanguineous in colour, a comparatively smaller round spot at the apical angles of the corium.....slanbuschii (Fabricius, 1787)
- 3. Physopelta schlansbuschi (Fabricius, 1787) (Fig. 3)

1787. Physopelta schlanbuschi Fabricius, Mant. Ins., 2: 299.

1902. Physopelta schlanbuschi Distant, Fauna.Brit. India. Rhynchota., 2: 99.

Material Examined: 2exs, Tuling, Dalma WLS, East Singbhum, Jharkhand, 22.8987N, 86.1158E, 8.vii.2021, Coll: J.Chitra & Party.

Distribution: India: Maharashtra, Madhya Pradesh, Assam, Uttarakhand, West Bengal, Chhattisgarh, Bihar

Elsewhere: China, Hong Kong, Mayanmar.

4. Physopelta gutta (Burmister, 1834) (Fig.4)

1834. Lygaeus (Pyrrhocoris) gutta Burmeister, Nova. Acta. Acad. Leop. Carolxvi, suppl., 424.

1902. Physopelta gutta: Distant, Fauna . Brit. India, Rhynchota., 11:97.

Material Examined: 2exs, Tuling, Dalma WLS, East Singbhum, Jharkhand, 22.8987N, 86.1158E, 8.vii.2021,Coll: Chitra J. & Party.

Distribution: India: Maharashtra, Assam, Andhra Pradesh, Uttarakhand, West Bengal, Karnataka, Madhya Pradesh, Arunachal Pradesh, Rajasthan.

*Elsewhere*: Australia, Borneo, Myanmar, China, Japan, Java, Philippines, Sumatra, Sri Lanka.

Family II. Pyrrhocoridae Key to the genera of the family Pyrrhocoridae

- 2. Head transversely constricted or impressed beneath near base..... *Antilochus* (Stal, 1863)
- 3. Anterior area of pronotum completely circumscribed by continuous impression or punctures. Head with eyes not broader than anterior margin of pronotum. Eyes not pedunculated.............. Euscopus (Stal,1870)

Genus: *Dysdercus* Amy. & Serv (1843) Key to the species of the genus *Dysdercus* 

- 2. Thorax wider posteriorly. White ventral fasciae varying from slightly to strongly bifid.
- 3. Vertical process of ninth segment broad and transparent.......Olivaceus (Fabr,1798)
- 5. Dysdercus evanescens (Dist,1902) (Fig.5)

1902. Dysdercus evanescens Distant, A.M.N.H., (7)9:43.

1904. Dysdercus evanescens Distant, Fauna. Brit. Ind. Rhynchota, 2: 120.

Material Examined: 1ex, Tuling, Dalma WLS, East Singbhum, Jharkhand, 22.8987N, 86.1158E, 8.vii.2021, Coll: Chitra J. & Party.

Distribution: India: Maharashtra, Assam, Meghalaya, Sikkim, Uttarakhand, West Bengal, Nagaland.

Elsewhere: Myanmar.

6. Dysdercus olivaceus (Fabr, 1798) (Fig. 6)

1798. Dysdercus olivaceous Fabricius, Ent. Syst. Suppl., p. 540.

1868. Dysdercus olivaceous Stal, Hem, Fabr, 1-84.

1904. Dysdercus olivaceous Distant, Fauna. Brit. Ind. Rhynchota, 2:119-120.

Material Examined: 1ex, Forest Guest House, Dalma WLS, East Singhbhum District, Jharkhand, 22.912579N, 86.143106E, 8. vii. 2021, Coll: Chitra J. & Party.

Distribution: India: Maharashtra, Tamil Nadu, Karnataka.

Elsewhere: Sri Lanka.

7. Dysdercus koenigii (Fabr, 1775) (Fig. 7)

1775. Cimex koenigii Fabricius, Syst. Ent. 720.

1904. Dysdercus cingulatus Distant, Fauna. Brit. Ind. Rhynchota, 2: 118.

1914. Dysdercus koenigi Bergroth, Ent. Mitt, 3:355.

Material Examined: 1ex, Tuling, Dalma WLS, East Singbhum, Jharkhand, 22.8987N, 86.1158E, 8.vii.2021, Coll: Chitra J. & Party; 1ex, Forest Guest House, Dalma WLS, East Singhbhum District, Jharkhand, 22.912579N, 86.143106E,8.vii.2021, Coll: Chitra J. & Party.

Distribution: India: Jharkhand, Maharashtra, Assam, Andaman & Nicobar Inlands, Madhya Pradesh, Karnataka, Meghalaya, Nagaland, Sikkim, Uttarakhand, West Bengal, Chhattisgarh.

Elsewhere: Australia, Myanmar, Sri Lanka, Pakisthan, Singapore, Bangkok, Sumatra, Java. Philippine, Solomon Island, Papua New Guinea.

Genus: Antilochus (Stal, 1863)

Key to the species of the genus Antilochus

- 1. Rounded black spot near base of membrane ....... russus (Stal,1863)
- 8. Antilochus russus (Stal, 1863) (Fig. 8)

1863. Antilochus russus Stal, Burl. Ent. Zeitschr., 7:394.

1904. Antilochus russus Distant, Fauna Brit. India, Rhynchota, 2: 101.

Material Examined: 1 ex., Makulakocha, Dalma Wls, Ranchi, Jharkhand,24.91187N, 86.13818E, 09.vii.2021, Coll: Chitra J. & Party;

Distribution: Andhra Pradesh, West Bengal, Sikkim, Assam, Meghalaya,

Elsewhere: Australia, Malayan Archipelago, Myanmar, Sri Lanka

9. Antilochus coquebertii (Fabr, 1803) (Fig. 9)

1803. Lygaeus coqueberti Fabricius, Syst. Rhyng.,: 222.

1904. Antilochus coqueberti Distant, Fauna Brit. India, Rhynchota, 2: 101

Material Examined: 1ex, Tuling, Dalma WLS, East Singbhum, Jharkhand, 22.8987N, 86.1158E, 8.vii.2021, Coll: Chitra J. & Party.

Distribution: India: Maharashtra, Andaman Islands, Madhya Pradesh, Andhra Pradesh, Assam, Jammu & Kashmir, Meghalaya, Nagaland, West Bengal. Elsewhere: Myanmar, Sri Lanka, Bhutan.

Genus: Euscopus (Stal, 1870)

Key to the species of the genus Euscopus

- 1 Greyish black or black in colour. A small yellowish spot near the apical margin of corium......indicorus (Walk,1872)
- 10. Euscopus indicorus (Walk, 1872) (Fig. 10)
- 1872. Pyrrhocoris indecorus Walker, Cat. Heter., 5: 172.
- 1902. Euscopus indecorus Distant, Ann . Mag .Nat .Hist.,  $\mathbf{9}$ : 37 .
- 1909. Euscopus indecorus: Lefroy, Rec .Indian Mus.,  ${\bf 3}$ : 326.

Material Examined: 1ex, Forest Guest House, Dalma WLS, East Singhbhum District, Jharkhand, 22.912579N, 86.143106E, 8. vii. 2021, Coll: Chitra J. & Party.

Distribution: India: Maharashtra, Assam, Meghalaya, Uttarakhand.

Elsewhere: Myanmar, Sri Lanka, Thailand.

11. Euscopus albatus(Dist,1909) (Fig.11)

1909. Euscopus albatus Distant, Ann . Mag . Nat . Hist., (8) · 496

1910. Euscopus albatus Distant, Fauna. Brit .India .Rhynchota., 5 : 94 .

Material Examined: 1 ex., Makula Kocha, Dalma Wls, Ranchi, Jharkhand,24.91187N, 86.13818E, 09.vii.2021, Coll: Chitra J. & Party

Distribution: India: Maharashtra, West Bengal, Madhya Pradesh.

Chattopadhyay (2024) reported only one species of Pyrrhocoridae from the state of Jharkhand, with no species documented from the Dalma Wildlife Sanctuary. This study marks the first effort to document the Pyrrhocoroidea diversity within the Dalma WLS. A total of 4 species of Largidae belonging to 3 genera and 7 species of Pyrrhocoridae belonging to 3 genera have been recorded from this sanctuary. Since the survey was conducted in only one season, collecting specimens in different seasons may reveal additional species, further enriching the Pyrrhocoroidea diversity in this region.

# **ACKNOWLEDGEMENT**

The authors are grateful to Dr. Dhriti Banerjee, Director, Zoological Survey of India for encouragements and laboratory facilities and to Dr. Guru Pada Mandal,

Divisional-in-charge, Entomology Division, Zoological Survey of India for his valuable suggestions and support. Thanks are due to Mr. Amartya Pal, SRF of Hemiptera Section, ZSI, Kolkata and Biju Mukherjee for their support and help to accomplish the work.

# CONFLICT OF INTERESTS

The authors declare no conflict of interest

# ETHICS APPROVAL

Not applicable

#### REFERENCE

- Ahmad, I., Qadri, S.S., (2009), Male and Female Genitalial Characters in the Establishment of a New Subgenus of Cotton Stainer's Genus *Dysdercus* Guérin-Méneville (Hemiptera: Pyrrhocoridae). *Pakistan Journal of Zoology*, 41(2),125-130.
- Chandra, K., Biswas, B., Kushwaha, S., (2015), Diversity of Pyrrhocoroidea (Hemiptera: Heteroptera) of Madhya Pradesh, India. *Munis Entomology & Zoology*, 10 (1), 169-171.
- Chattopadhyay, S., (2024), Some new records and full checklist of hemipteran insects on shisham (*Dalbergia sissoo* Roxb.) from Jharkhand, India. *Journal of Entomology and Zoology Studies*, 12(3), 210-213.
- Distant, W. L., (1902), The fauna British India including Ceylon and Burma, Rhynchota Vol. II. Taylor and Francis, London, 196-430 pp.
- Distant, W. L., (1910), Fauna Brit. India Including Ceylon and Burma, Rhynchota Vol. V. Taylor and Francis, London, 92:100 pp.
- Hassan, M. E., Biswas, B., Saha, P. C., Mukherjee,
  P., Praveen, K., (2019), (Insecta: Hemiptera: Heteroptera).
  In Fauna of Maharashtra, State Fauna Series, (Eds. Director of Zoological Survey of India) 20(Part-3):
  Zoological Survey of India, Kolkata. 41-87 pp.
- Hemala, V., Kment, P., Malenovský, I., (2015), Morphology and phylogeny of the true bug superfamily Pyrrhocoroidea (Heteroptera: Pentatomomorpha)—a preliminary report.7<sup>th</sup> European Hemiptera Congress, Graz. Austria.
- Praveen, K., Hassan, M.E., Saha, P.C., Kushwaha, S., Dubey, A.K., Dash, S., Lyngdoh J., Pal, A., Jehamalar, E., Khanra, S., Fauna of India Checklist. Arthropoda: Insecta: Hemiptera. Version 1.0. Zoological Survey of India.

Stehlík, J. L., Kment, P., (2014), Riegeriana, a new genus for Physopelta apicalis, and checklist of the genus Iphita (Hemiptera: Heteroptera: Largidae) with description of one new species. Zootaxa, 3860 (2), 167-183.

Stehlík, J. L, (2013), Review and reclassification of the Old World genus Physopelta (Hemiptera: Heteroptera: Largidae). Acta Entomologica Musei Nationalis Pragae,53(2), 505-584.