



ASSESSMENT OF AN ANTIMICROBIAL PROPERTY OF SELECTED ANTIBIOTICS AGAINST THE SELECTED BACTERIAL FOOD PATHOGENS

Anand Pundlik Bothe

Regional Public Health Laboratory, Near NIT Sadar Nagpur - 440001

Article History: Received 10th November 2025; Accepted 8th January 2026; Published 31st January 2026

ABSTRACT

Foodborne bacterial pathogens are a leading cause of food-related illnesses worldwide, posing significant public health and economic challenges. The increasing prevalence of antibiotic resistance among these pathogens necessitates continuous evaluation of antimicrobial agents. The present study investigates the antimicrobial efficacy of selected antibiotics against major bacterial food pathogens, including *Escherichia coli*, *Salmonella enterica*, *Staphylococcus aureus*, *Listeria monocytogenes*, and *Bacillus cereus*. Antibiotics belonging to different classes ampicillin, ciprofloxacin, tetracycline, chloramphenicol, and gentamicin—were assessed using agar disc diffusion, agar well diffusion, and minimum inhibitory concentration (MIC) assays. The results demonstrated differential susceptibility patterns among the tested pathogens. Ciprofloxacin and gentamicin exhibited the highest antimicrobial activity, while reduced sensitivity and resistance were observed against ampicillin and tetracycline, particularly in Gram-negative bacteria. Gram-positive pathogens showed comparatively higher susceptibility to most antibiotics. The observed resistance trends highlight the potential risks associated with the indiscriminate use of antibiotics in food production systems. This study emphasizes the importance of routine antimicrobial surveillance and rational antibiotic use to ensure food safety and mitigate the spread of resistant foodborne pathogens.

Keywords: Foodborne bacteria, Antibiotics, Antimicrobial activity, MIC, Antibiotic resistance, Food safety.

INTRODUCTION

Foodborne diseases remain a significant global public health challenge, affecting millions of people annually and resulting in substantial economic losses due to medical expenses, loss of productivity, and food recalls (<https://www.who.int/news-room/fact-sheets/detail/food-safety>). Bacterial pathogens are among the most common causative agents of foodborne illnesses, with *Escherichia coli*, *Salmonella spp.*, *Staphylococcus aureus*, *Listeria monocytogenes*, and *Bacillus cereus* frequently implicated in outbreaks associated with contaminated food and water (Yang *et al.*, 2025). These microorganisms can contaminate food at various stages of production, processing, storage, and distribution, making effective control measures essential for ensuring food safety (Grace, 2025). Antibiotics have long been used as a primary means of treating bacterial infections in humans and animals (Enshaie *et al.*, 2025). In food production systems,

antibiotics have also been employed for therapeutic, prophylactic, and growth-promoting purposes, particularly in livestock (Odey *et al.*, 2024). However, the widespread and often indiscriminate use of antibiotics has contributed to the emergence and dissemination of antibiotic-resistant bacteria. Foodborne pathogens exhibiting resistance to commonly used antibiotics pose a serious threat, as infections caused by such strains are more difficult to treat and may result in increased morbidity and mortality (Manyi-Loh *et al.*, 2018).

Gram-negative bacteria such as *E. coli* and *Salmonella* are particularly concerning due to their intrinsic resistance mechanisms, including the presence of an outer membrane that restricts antibiotic entry and the ability to acquire resistance genes through horizontal gene transfer (Breijyeh *et al.*, 2020). Gram-positive pathogens like *S. aureus* and *L. monocytogenes*, although generally more susceptible, have also developed resistance to several antibiotics, including

*Corresponding Author: Anand Pundlik Bothe, Regional Public Health Laboratory, Near NIT Sadar Nagpur – 440001. Email: bothepatil@gmail.com.

β -lactams and tetracyclines. The persistence of resistant strains in food matrices increases the risk of their transmission to humans through the food chain (Punchihewage-Don *et al.*, 2024; Galgano *et al.*, 2025).

Assessment of the antimicrobial activity of antibiotics against foodborne pathogens is crucial for understanding current resistance trends and guiding appropriate therapeutic strategies (Gajic *et al.*, 2022; Farrukh *et al.*, 2025). In vitro susceptibility testing provides valuable insights into the effectiveness of antibiotics and helps in identifying emerging resistance patterns. Therefore, the present study aims to evaluate the antimicrobial properties of selected antibiotics against common bacterial food pathogens isolated from food sources (Yamin *et al.*, 2023; Belay *et al.*, 2024; Bayot *et al.*, 2025). The findings of this study will contribute to a better understanding of antibiotic efficacy and support the development of informed policies for antibiotic usage in food safety and public health.

MATERIALS AND METHODS

Bacterial Strains

The bacterial food pathogens used in this study included *Escherichia coli*, *Salmonella enterica*, *Staphylococcus aureus*, *Listeria monocytogenes*, and *Bacillus cereus*. Standard reference strains were obtained from a recognized microbial culture collection and maintained on nutrient agar slants at 4 °C until use.

Antibiotics Tested

The antibiotics selected for this study were ampicillin (β -lactam), ciprofloxacin (fluoroquinolone), tetracycline (tetracycline class), chloramphenicol (phenicol), and gentamicin (aminoglycoside). These antibiotics represent different classes and are commonly used in clinical and veterinary medicine (Nhung *et al.*, 2016; Afroze *et al.*, 2025).

Preparation of Inoculum

Bacterial cultures were grown overnight in nutrient broth at 37 °C. The turbidity of the cultures was adjusted to match the 0.5 McFarland standard, corresponding to approximately 1×10^8 CFU/mL. Cultures were maintained on nutrient agar and sub-cultured prior to experimentation (Gayathiri *et al.*, 2018; Bhirdwaj *et al.*, 2025).

Agar Disc Diffusion Method

The antimicrobial activity of antibiotics was evaluated using the Kirby–Bauer disc diffusion method. Mueller–Hinton agar plates were inoculated with standardized bacterial suspensions using sterile cotton swabs. Antibiotic discs were placed on the surface of the agar and plates were incubated at 37 °C for 18–24 hours. Zones of inhibition were measured in millimeters and interpreted according to Clinical and Laboratory Standards Institute (CLSI) guidelines (Bauer *et al.*, 1966; Bayot *et al.*, 2024).

Agar Well Diffusion Method

Sterile wells of 6 mm diameter were punched into inoculated agar plates. Antibiotic solutions of known concentrations were added into the wells, and plates were incubated under the same conditions. The diameter of the inhibition zones was measured to assess antimicrobial activity (Balouiri *et al.*, 2016).

Minimum Inhibitory Concentration (MIC)

The MIC of each antibiotic was determined using the broth dilution method. Serial dilutions of antibiotics were prepared in nutrient broth, inoculated with bacterial suspensions, and incubated at 37 °C for 24 hours. The lowest concentration showing no visible growth was recorded as the MIC (Kowalska-Krochmal *et al.*, 2021).

Statistical Analysis

All experiments were performed in triplicate. Results were expressed as mean $\pm$ standard deviation. Statistical analysis was carried out using appropriate software, and differences were considered significant at $p < 0.05$.

RESULTS AND DISCUSSION

The results of the antimicrobial susceptibility tests revealed significant variations in the sensitivity of the tested foodborne pathogens to the selected antibiotics. Ciprofloxacin and gentamicin exhibited the highest antimicrobial activity against all tested bacteria, as indicated by larger zones of inhibition and lower MIC values. These findings are consistent with previous studies reporting the broad-spectrum efficacy of fluoroquinolones and aminoglycosides (Shariati *et al.*, 2022). Table 1 below represents the zones of inhibitions shown by test antibiotics against the selected pathogens from foods.

Table 1. Zone of inhibition (mm) of antibiotics against foodborne pathogens.

Pathogen	AMP	CIP	TET	CHL	GEN
<i>E. coli</i>	10 $\pm$ 0.5	26 $\pm$ 1.2	12 $\pm$ 0.6	18 $\pm$ 0.9	24 $\pm$ 1.0
<i>Salmonella</i>	11 $\pm$ 0.7	25 $\pm$ 1.1	13 $\pm$ 0.8	17 $\pm$ 0.7	23 $\pm$ 0.8
<i>S. aureus</i>	18 $\pm$ 0.9	28 $\pm$ 1.4	20 $\pm$ 1.0	22 $\pm$ 1.1	26 $\pm$ 1.3
<i>L. monocytogenes</i>	17 $\pm$ 0.8	27 $\pm$ 1.3	19 $\pm$ 0.9	21 $\pm$ 1.0	25 $\pm$ 1.2
<i>B. cereus</i>	16 $\pm$ 0.6	26 $\pm$ 1.1	18 $\pm$ 0.7	20 $\pm$ 0.8	24 $\pm$ 1.0




Figure 1. Zone of inhibition in mm for antibiotics against test foodborne pathogens.

The figure 1, illustrates the antimicrobial efficacy of five standard antibiotics ampicillin (AMP), ciprofloxacin (CIP), tetracycline (TET), chloramphenicol (CHL), and gentamicin (GEN) against selected bacterial food pathogens (*Escherichia coli*, *Salmonella* sp., *Staphylococcus aureus*, *Listeria monocytogenes*, and *Bacillus cereus*), expressed as zone of inhibition in millimeters. Among the tested compounds, ciprofloxacin exhibited the highest antibacterial activity across all pathogens, with inhibition zones ranging approximately from 25–28 mm, indicating its broad-spectrum and potent efficacy. Gentamicin also showed strong inhibitory effects (22–26 mm), followed by chloramphenicol (15–22 mm), suggesting moderate to high sensitivity of both Gram-positive and Gram-negative bacteria. Tetracycline demonstrated comparatively lower activity, particularly against *E. coli* and *Salmonella* sp., while ampicillin showed the least effectiveness overall, especially against Gram-negative strains. Notably, *S. aureus* and *L. monocytogenes* were generally more susceptible to most antibiotics than *E.*

coli and *Salmonella* sp. The presence of error bars indicates experimental variability but does not alter the overall trend, highlighting ciprofloxacin and gentamicin as the most effective antibiotics against the selected foodborne pathogens.

Ciprofloxacin and gentamicin consistently produced the largest inhibition zones, indicating strong antimicrobial activity. Ampicillin and tetracycline showed reduced effectiveness, particularly against Gram-negative bacteria. Ampicillin showed limited effectiveness, particularly against Gram-negative bacteria such as *E. coli* and *Salmonella*, which may be attributed to the production of β -lactamase enzymes. Tetracycline resistance was observed in both Gram-positive and Gram-negative bacteria, reflecting its extensive use in agriculture and clinical settings. Chloramphenicol demonstrated moderate activity; however, its use has declined due to associated toxicity concerns (Kakoullis *et al.*, 2021; Sitohy *et al.*, 2024). Table 2 represents the minimal inhibitory concentrations (MIC) against tested foodborne pathogens.

Table 2. MIC values ($\mu\text{g/mL}$) of antibiotics.

Pathogen	AMP	CIP	TET	CHL	GEN
<i>E. coli</i>	32	0.5	16	8	1
<i>Salmonella</i>	32	0.5	16	8	1
<i>S. aureus</i>	8	0.25	4	4	0.5
<i>L. monocytogenes</i>	8	0.25	4	4	0.5
<i>B. cereus</i>	8	0.5	4	4	1

Lower MIC values for ciprofloxacin and gentamicin confirm their superior efficacy.




Figure 2. MIC values for test antibiotics against selected foodborne pathogens.

The figure 2, depicts the minimal inhibitory concentrations (MICs, $\mu\text{g/mL}$) of five antibiotics—ampicillin (AMP), ciprofloxacin (CIP), tetracycline (TET), chloramphenicol (CHL), and gentamicin (GEN)—against major foodborne pathogens (*Bacillus cereus*, *Listeria monocytogenes*, *Staphylococcus aureus*, *Salmonella sp.*, and *Escherichia coli*). Ampicillin exhibited the highest MIC values (approximately 30–35 $\mu\text{g/mL}$), particularly against *Salmonella sp.* and *E. coli*, indicating comparatively lower antibacterial potency. Tetracycline showed moderate MICs, with *Salmonella sp.* and *E. coli* requiring higher concentrations (~15–17 $\mu\text{g/mL}$) than Gram-positive bacteria. Chloramphenicol demonstrated improved efficacy with lower MICs (around 6–10 $\mu\text{g/mL}$), reflecting broader activity across both Gram-positive and Gram-negative strains. Ciprofloxacin and gentamicin displayed the lowest MIC values (generally <2 $\mu\text{g/mL}$) for all tested organisms, highlighting their superior antibacterial effectiveness. Overall, the MIC trends corroborate the strong potency of ciprofloxacin and gentamicin, while ampicillin appears least effective against the selected food pathogens. Gram-positive bacteria, especially *Staphylococcus aureus* and *Bacillus cereus*, were generally more susceptible to the tested antibiotics compared to Gram-negative bacteria. The structural differences in cell walls likely contribute to this observed variation. The presence of resistant strains among foodborne pathogens underscores the need for judicious antibiotic use and regular surveillance (Karaman *et al.*, 2020; Alhumaid *et al.*, 2021).

The differential susceptibility patterns observed in this study align with earlier reports on foodborne pathogens. The reduced effectiveness of ampicillin may be attributed to β -lactamase production, while tetracycline resistance likely reflects its extensive use in agriculture. Gram-positive bacteria showed higher susceptibility due to simpler cell wall structures, whereas Gram-negative bacteria exhibited increased resistance due to outer membrane barriers. These findings reinforce the urgent

need for antimicrobial stewardship and continuous resistance monitoring in food safety systems (Mwansa *et al.*, 2022; Grudlewska-Buda *et al.*, 2023; Wojnarowski *et al.*, 2025). The findings highlight the importance of continuous monitoring of antibiotic resistance patterns in food-associated bacteria to prevent the spread of resistant strains through the food chain.

CONCLUSION

The present study demonstrates that selected antibiotics exhibit varying degrees of antimicrobial activity against common bacterial food pathogens. Ciprofloxacin and gentamicin were the most effective antibiotics, while resistance to ampicillin and tetracycline was evident in several bacterial strains. The emergence of antibiotic-resistant foodborne pathogens poses a serious threat to public health and emphasizes the need for responsible antibiotic usage and robust food safety practices. Regular surveillance and susceptibility testing are essential to guide effective treatment strategies and minimize the risk of resistance development.

ACKNOWLEDGMENT

The authors express sincere thanks to the head of the Regional Public Health Laboratory, Near NIT Sadar Nagpur for the facilities provided to carry out this research work.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

ETHICS APPROVAL

Not applicable

FUNDING

This study received no specific funding from public, commercial, or not-for-profit funding agencies.

AI TOOL DECLARATION

The authors declares that no AI and related tools are used to write the scientific content of this manuscript.

DATA AVAILABILITY

Data will be available on request

REFERENCES

- Afroze, S., Faisal, M., Khan, M. N. A., & Barua, H. (2025). A comprehensive review of antibiotics and antimicrobial resistance in the aquaculture sector of the world and Bangladesh. *International Journal of Microbiology*, 2025, 8818516. <https://doi.org/10.1155/ijm/8818516>.
- Alhumaid, S., Al Mutair, A., Al Alawi, Z., Alzahrani, A. J., Tobaiqy, M., Alresasi, A. M., Bu-Shehab, I., Al-Hadary, I., Alhmeed, N., Alismail, M., Aldera, A. H., AlHbabi, F., Al-Shammari, H., Rabaan, A. A., & Al-Omari, A. (2021). Antimicrobial susceptibility of gram-positive and gram-negative bacteria: A 5-year retrospective analysis at a multi-hospital healthcare system in Saudi Arabia. *Annals of Clinical Microbiology and Antimicrobials*, 20(1), 43. <https://doi.org/10.1186/s12941-021-00450-x>.
- Balouiri, M., Sadiki, M., & Ibnsouda, S. K. (2016). Methods for in vitro evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis*, 6(2), 71–79. <https://doi.org/10.1016/j.jpah.2015.11.005>.
- Bauer, A. W., Kirby, W. M. M., Sherris, J. C., & Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology*, 36, 493–496.
- Bayot, M. L., & Bragg, B. N. (2025). Antimicrobial susceptibility testing. In *StatPearls* [Internet]. Treasure Island, FL: StatPearls Publishing. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK539714/>.
- Belay, W. Y., Getachew, M., Tegegne, B. A., Teffera, Z. H., Dagne, A., Zeleke, T. K., Abebe, R. B., Gedif, A. A., Fenta, A., Yirdaw, G., Tilahun, A., & Aschale, Y. (2024). Mechanism of antibacterial resistance, strategies and next-generation antimicrobials to contain antimicrobial resistance: A review. *Frontiers in Pharmacology*, 15, 1444781. <https://doi.org/10.3389/fphar.2024.1444781>.
- Bharadwaj, A., Nayariseri, A., & Singh, S. K. (2025). Characterization of natural therapeutic compounds producing novel bacterial strains isolated from Hyderabad, India. *Scientific Reports*, 15(1), 25681. <https://doi.org/10.1038/s41598-025-11615-5>.
- Breijyeh, Z., Jubeh, B., & Karaman, R. (2020). Resistance of gram-negative bacteria to current antibacterial agents and approaches to resolve it. *Molecules*, 25(6), 1340. <https://doi.org/10.3390/molecules25061340>.
- Enshaie, E., Nigam, S., Patel, S., & Rai, V. (2025). Livestock antibiotics use and antimicrobial resistance. *Antibiotics*, 14(6), 621. <https://doi.org/10.3390/antibiotics14060621>.
- Farrukh, M., Munawar, A., Nawaz, Z., Hussain, N., Hafeez, A. B., & Szveda, P. (2025). Antibiotic resistance and preventive strategies in foodborne pathogenic bacteria: A comprehensive review. *Food Science and Biotechnology*, 34, 2101–2129. <https://doi.org/10.1007/s10068-024-01767-x>.
- Gajic, I., Kabic, J., Kekic, D., Jovicevic, M., Milenkovic, M., Mitic-Culafic, D., Trudic, A., Ranin, L., & Opavski, N. (2022). Antimicrobial susceptibility testing: A comprehensive review of currently used methods. *Antibiotics*, 11(4), 427. <https://doi.org/10.3390/antibiotics11040427>.
- Galgano, M., Pellegrini, F., Catalano, E., Capozzi, L., Del Sambro, L., Sposato, A., Lucente, M. S., Vasinioti, V. I., Catella, C., Odigie, A. E., Tempesta, M., Pratelli, A., & Capozza, P. (2025). Acquired bacterial resistance to antibiotics and resistance genes: From past to future. *Antibiotics*, 14(3), 222. <https://doi.org/10.3390/antibiotics14030222>.
- Gayathiri, E., Bharathi, B., & Priya, K. (2018). Study of the enumeration of twelve clinically important bacterial populations at 0.5 McFarland standard. *International Journal of Creative Research Thoughts*, 6, 880–893.
- Grudlewska-Buda, K., Bauza-Kaszewska, J., Wiktorczyk-Kapischke, N., Budzyńska, A., Gospodarek-Komkowska, E., & Skowron, K. (2023). Antibiotic resistance in selected emerging bacterial foodborne pathogens—An issue of concern. *Antibiotics*, 12(5), 880. <https://doi.org/10.3390/antibiotics12050880>.
- World Health Organization. (n.d.). *Food safety*. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/food-safety>.
- Kakoullis, L., Papachristodoulou, E., Chra, P., & Panos, G. (2021). Mechanisms of antibiotic resistance in important gram-positive and gram-negative pathogens and novel antibiotic solutions. *Antibiotics*, 10(4), 415. <https://doi.org/10.3390/antibiotics10040415>.
- Karaman, R., Jubeh, B., & Breijyeh, Z. (2020). Resistance of gram-positive bacteria to current antibacterial agents and overcoming approaches. *Molecules*, 25(12), 2888. <https://doi.org/10.3390/molecules25122888>.
- Kowalska-Krochmal, B., & Dudek-Wicher, R. (2021). The minimum inhibitory concentration of antibiotics: Methods, interpretation, clinical relevance. *Pathogens*, 10(2), 165. <https://doi.org/10.3390/pathogens10020165>.
- Manyi-Loh, C., Mamphweli, S., Meyer, E., & Okoh, A. (2018). Antibiotic use in agriculture and its consequential resistance in environmental sources: Potential public

- health implications. *Molecules*, 23(4), 795. <https://doi.org/10.3390/molecules23040795>.
- Mwansa, T. N., Kamvuma, K., Mulemena, J. A., Phiri, C. N., & Chanda, W. (2022). Antibiotic susceptibility patterns of pathogens isolated from laboratory specimens at Livingstone Central Hospital in Zambia. *PLOS Global Public Health*, 2(9), e0000623. <https://doi.org/10.1371/journal.pgph.0000623>.
- Nhung, N. T., Cuong, N. V., Thwaites, G., & Carrique-Mas, J. (2016). Antimicrobial usage and antimicrobial resistance in animal production in Southeast Asia: A review. *Antibiotics*, 5(4), 37. <https://doi.org/10.3390/antibiotics5040037>.
- Odey, T. O. J., Tanimowo, W. O., Afolabi, K. O., Jahid, I. K., & Reuben, R. C. (2024). Antimicrobial use and resistance in food animal production: Food safety and associated concerns in Sub-Saharan Africa. *International Microbiology*, 27, 1–23. <https://doi.org/10.1007/s10123-023-00462-x>.
- Punchihewage-Don, A. J., Ranaweera, P. N., & Parveen, S. (2024). Defense mechanisms of *Salmonella* against antibiotics: A review. *Frontiers in Antibiotics*, 3, 1448796. <https://doi.org/10.3389/frabi.2024.1448796>.
- Shariati, A., Arshadi, M., Khosrojerdi, M. A., Abedinzadeh, M., Ganjalishahi, M., Maleki, A., Heidary, M., & Khoshnood, S. (2022). The resistance mechanisms of bacteria against ciprofloxacin and new approaches for enhancing the efficacy of this antibiotic. *Frontiers in Public Health*, 10, 1025633. <https://doi.org/10.3389/fpubh.2022.1025633>.
- Sitohy, M., Enan, G., Abdel-Shafi, S., El-Wafa, N. A., El-Gazzar, N., Osman, A., & Sitohy, B. (2024). Mapping pathogenic bacteria resistance against common antibiotics and their potential susceptibility to methylated white kidney bean protein. *BMC Microbiology*, 24, 49. <https://doi.org/10.1186/s12866-024-03202-x>.
- Wojnarowski, K., Cholewińska, P., Zhao, D., Pacoń, J., & Bodkowski, R. (2025). Antibiotic resistance genes in food animal production: Environmental implications and One Health challenges. *Environments*, 12(11), 427. <https://doi.org/10.3390/environments12110427>.
- Yamin, D., Uskoković, V., Wakil, A. M., Goni, M. D., Shamsuddin, S. H., Mustafa, F. H., Alfouzan, W. A., Alissa, M., Alshengeti, A., Almaghrabi, R. H., Fares, M. A. A., Garout, M., Al Kaabi, N. A., Alshehri, A. A., Ali, H. M., Rabaan, A. A., Aldubisi, F. A., Yean, C. Y., & Yusof, N. Y. (2023). Current and future technologies for the detection of antibiotic-resistant bacteria. *Diagnostics*, 13(20), 3246. <https://doi.org/10.3390/diagnostics13203246>.
- Yang, Y., Xu, F. R., Zhou, Y. B., Hu, L. Q., Lu, W., Zhang, S. H., Hu, H., & Huang, X. E. (2025). Epidemiological characteristics of foodborne disease outbreaks in a hospital: A 5-year retrospective study. *International Journal of General Medicine*, 18, 1529–1542. <https://doi.org/10.2147/IJGM.S496799>.