

TEMPORAL CHANGES IN CHEMICAL AND HEAVY METAL CHARACTERISTICS OF CRUDE OIL POLLUTED TOP SOILS OF IZOMBE, SOUTHEASTERN, NIGERIA

¹Ihem, E.E, ¹Aririguzo, B.N, ²Ogueri, C, ²Anyanwu, C.N, ¹Agim, L.C, ²Nwaka, D, ²Chinnah, U.K, ³Madu, H.C, ⁴Nwaka, S.U, ³Agorua, U.N, ⁵Nwokeke, B.C, ⁶Ihezue, C.I, ^{7*}Ndukwu. R. I, ¹Chukwu, E.D, ¹Osisi, A and ⁸Chikaire, J.U

¹Department of Soil Science Technology, Federal University of Technology, Owerri, Imo State

²Department of Fisheries and Aquaculture, Federal University of Technology, Owerri, Imo State

³Department of Fisheries Technology, Imo State, Polytechnic, Omuma, Imo State

⁴Department of Fisheries Technology, Federal Polytechnic, Nekede, Owerri; Imo State

⁵Department of Food Science and Technology, University of Agriculture and Environmental Sciences, Umuagwo, Owerri, Imo State

⁶National Root Crops Research Institute, Umudike, Abia State

⁷Department of Geoinformatics and Surveying University of Nigeria, Enugu Campus, Enugu State

⁸Department of Agricultural Extension, Federal University of Technology, Owerri, Imo State, Nigeria

Article History: Received 22nd December 2025; Accepted 19th February 2026; Published 1st March 2026

ABSTRACT

Crude oil pollution is a significant environmental issue in Izombe, Imo State, Nigeria, and an oil-producing region experiencing frequent oil spillage in the area. Izombe lies in the northeast of Niger Delta and covers about 59 km² in Njaba River. The study was aimed at investigating the physico-chemical and heavy metal characteristics of top soils (0-15 cm) depth of Izombe oil spill fields, 2024. A free survey technique was adopted in the field sampling of seven (7) stations (5 polluted and 2 control) in the months of April and June, 2024. Standard laboratory procedures were followed to determine some physical, chemical and heavy metal concentrations of the study area. Results showed that crude oil pollution increased the Electrical Conductivity (EC) by 148% and 290% in April and June respectively; calcium by 209% and 171%; potassium by 208% and 156%; magnesium by 211% and 170% in April and June respectively. Crude oil pollution increased zinc (Zn) by 150% and 142% in April and June respectively; copper (Cu) by 28% and 20%, lead (Pb) by 815 and 89% in April and June respectively. The study showed that crude oil pollution affected most of the basic soil properties significantly and increased with time. Consequently, heavy metal values from the results were found to be below the critical limits, and as such may not pose a serious threat to the terrestrial and aquatic environment. However, standard procedures should be adopted in the oil producing activities to prevent soil and water against contamination beyond permissible limits.

Keywords: Oil spillage, Heavy metals, Permissible limits, Crude oil, Izombe.

INTRODUCTION

Petroleum industries in Nigeria have eventually created economic boom and remains the backbone of the Nigerian economy contributing over 90% of its revenue and about 45% of its GDP (Akinlo, 2012; Ekeghe, 2022). Izombe, Imo State being one of the nine major states in southern

states in Nigeria produce crude oil in very large quantities (Ejechi, 2022). Available data showed that over 9,500 cases of oil spillage occurred in Nigeria from 2011 to 2021 with an estimate of about 450,000 barrels of oil released into the environment (Akinpelu, 2021). The release of large quantity of crude oil in the environment in form of spillage significantly result to substantial level of soil pollution,

*Corresponding Author: NDUKWU, R.I, Department of Agricultural Extension, Federal University of Technology, Owerri, Imo State, Nigeria Email: raphael.ndukwu@unn.edu.ng

deforestation, toxic discharges into available water sources, biodiversity loss, as well as recurring and extensive oil spills (Owugah, 2006; Chijioke *et al.*, 2018; Akinpelu, 2021). Chijioke *et al.* (2018) in a review opined that farmers have been compelled to evacuate their farmlands and properties due to increasing soil sterility caused by the depletion of soil microbes and diminishing crop yields. It is obvious that crude oil contains metals and their mobility depends on the concentration and soil properties. Similar studies on crude oil polluted areas indicated that oil spillage ruined the subsistence economic activities of the affected areas, drinking water sources lost, agricultural and fishing operations hampered and the ecology of the region deteriorated. According to Moneke *et al.* (2022), air in the affected areas is also polluted by oil spillage. Impact of oil spillage on coastal and aquatic environments can be both short and long term (Kujawinski *et al.*, 2020; Al-Hamadani *et al.*, 2022). It also affects generally the ecosystem, soil properties and food production (Wokocha *et al.*, 2011), seed germination and vegetation cover hampered (Gulshen and Datsi, 2012).

Soil health, soil and water toxicity increased with high heavy metal concentration (Ihem *et al.*, 2015). Research works have been carried out on the effects of oil spillage

without much emphasis on time variations. Therefore, this research finding was aimed at determining the physico-chemical, and heavy metal characteristics of the crude oil polluted surface soils with time in the study sites.

MATERIALS AND METHODS

Site Description

Izombe lies in the Northeast of Niger Delta. It covers about 59 km² in Njaba River basin, Imo State. It comprises of Izombe and Ossu oil fields (Figure 1). The experiment was conducted at seven (7) oil and gas exploration sites in Izombe, Imo State, Nigeria as shown in Table. 1 in April and June, 2024.

A free survey technique was adopted in the choice of sampling sites. Surface soil samples were collected at a depth of 0 – 15cm in each of the seven sites (five polluted and two unpolluted as control). Samples collected in each of these sites were Pseudo-replicated and the means were then used for statistical analysis. Soil auger (2 auger bores) was used to collect the soil samples, homogenized to form a composite sample. The samples were collected in foil packs and nylon bags, properly labelled and preserved for analysis in line with the world standard methods.

Table 1. Sampling location sites of Izombe oil fields.

S. No	Sampling sites	Longitude	Latitude
1.	Flare Area (200m away)	05°37'08.6"N	006°49'02.1"E
2.	Process Area	05°37'04.6"N	006°48'59.3"E
3.	Generator Area	05°37'03.4"N	006°49'00.4"E
4.	Compressor Area	05°37'03.9"N	006°49'02.8"E
5.	Accommodation Area	05°37'06.6"N	006°49'07.6"E
6.	Community Control Point (1)	05°38'32.7"N	006°50'57.4"E
7.	Community Control Point (2)	05°38'36.1"N	006°50'56.7"E




Figure 1. Map showing Sampling points.

Legend: SS = Soil Sample Points , AQ = Air Quality Points, HDW = Hand Dug Well Points, SW = Surface Water Points

However, soils from Izombe are derived from coastal plain sands and are dominated by ultisols and inceptisols (Onweremadu and Peter, 2016). The study sites belong to the lowland area of Nigeria, with a humid tropical climate having a mean annual temperature range of 26 - 31°C and annual rainfall ranges between 1500mm to 2500mm which runs from March to December with its peak in July, through October/November to March with a dry dust and cold intervals, locally termed as Harmattan. The vegetation of the area is tropical rainforest zone mostly dominated by grasses like *Sida acuta*, *Panicum maximum*, *Pennisetum purpureum*, *Cromolima odorata*, *Centrosima purpescens* among others.

Laboratory analytical methods

Soil pH: Soil pH was determined electrometrically with a Horiba Multi-parameter water checker (U-50 series) in a solution ratio of 1:2.5 according to Bremner, (1996). **Electrical Conductivity (us/cm) (EC)** was determined in a ratio of 1:2 soil-to-water ratio by APHA 2510B electrometric method using a calibrated Horiba Multi-parameter water checker as described by Hendershot *et al* (1993). **Total Organic Carbon (%) (TOC)** was determined using the titrimetric dichromate oxidation method (Walkley-Black's method) as described by Nelson and Sommers, (1982). **Exchangeable Cations (Na⁺, K⁺, Ca²⁺ and Mg²⁺) mg/kg:** The level of cations was determined in the soil samples using standard procedure with an Atomic Absorption Spectrophotometer (Shimadzu AA-6650) and (ASTM D1971/4691) as described by Sparks (1996). **Total Hydrocarbon Content (mg/kg) ASTM D3921** was employed for the determination of the THC. Solutions of total hydrocarbon content was further examined by infrared spectroscopy. **Exchangeable Anions (SO₄²⁻, NO₃⁻ & PO₄³⁻):** This was analyzed using international standard methods (APHA 4500- NO₃⁻, E for Nitrate, APHA 4500- P.C for phosphate). **Heavy metals (mg/kg):** Heavy metal

concentration in the studied soils were measured using standard procedure with an Atomic Absorption Spectrophotometer (Shimadzu AA-6650) and a standard method (ASTM D1971/4691) as described by Beckhof *et al* (2006). **Hydrocarbon Utilizing Bacteria, cfu/g:** Bacteria in the soil capable of utilizing hydrocarbon as their sole source of carbon and energy were determined using APHA 9215C (Spread plate) test Methods.

RESULTS AND DISCUSSION

The results of the Physico-chemical properties of the top soils samples in Izombe oil spill fields, 2024 is presented in Table 2. The mean value of the soil pH in April was 5.85 and 5.84 in June. T-test analysis showed that there were no significant differences in the soil pH from the seven sampling oil spill fields. However, the pH value is rated moderately acidic according to the ratings of Babalola *et al.* (1998). Soils of the study area are from coastal plains (ultisols), highly weathered as a result of rainfall. The electrical Conductivity showed a mean value of 182.2 (us/cm) in April and 177.4 (us/cm) in June, indicating an increase by 148% in April and 290% in June influenced by oil spillage. High EC value in the soil is an indication of salinity (EC >4 ds/m) problems, which may impede crop growth and yield as well as microbial activities (Abu-Hassanein *et al.*, 1996; Choo *et al.*, 2022; Comma *et al.*, 2011; Moral *et al.*, 2008). The mean value for Total Organic Carbon (TOC) in April was 0.60% and 0.2% in June, showing an increase of 63% in April and only 4% in June. It was generally observed that TOC relatively were higher in April than in June. possible, there could be cases of more spillage occurring in the month of April than in June resulting to more accumulation of hydrocarbon in the affected fields where immobilization of these substances occurred. This finding agrees with that of Ihem *et al.* (2015).

Table 2. Characteristics of surface soil samples in Izombe oil spill field 2024.

	Sampling Sites (0-15cm) Depth																	
	APRIL, 2024									JUNE, 2024								
Soil parameters	SL1	SL2	SL3	SL4	SL5	Mean	SL6	SL7	Mean	SL1	SL2	SL3	SL4	SL5	Mean	SL6	SL7	Mean
pH	5.87	5.63	5.84	5.90	5.79	5.81	5.98	5.95	5.97	5.92	5.88	5.86	5.79	5.82	5.85	5.73	5.94	5.84
Electrical conductivity μ s/cm	53.0	145	72.0	269	372	182.2	65.0	82.0	5.97	42.0	138	64.0	286	357	177.4	51.0	40.0	45.5
Total Org. Carbon (%)	0.078	0.314	0.629	0.94	1.06	0.60	1.10	1.10	73.5	0.314	0.197	0.393	0.943	0.708	0.53	0.943	0.157	0.55
Total hydrocarbon (mg/kg)	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	0.98	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10
Av. phosphorus (mg/kg)	<0.095	0.724	1.65	2.73	3.84	1.81	2.11	3.72	2.92	<0.095	<0.095	3.80	6.08	5.12	3.04	0.623	1.95	1.23
Total Nitrogen (mg/kg)	0.032	0.046	0.68	0.087	0.098	0.066	0.84	0.099	0.092	0.047	0.039	0.052	0.089	0.073	0.06	0.189	0.037	0.063
Calcium	3.06	10.0	4.66	25.9	30.2	14.7	4.17	5.39	4.78	2.99	9.84	4.45	25.6	30.0	14.5	4.68	6.17	5.38

(mg/kg)						6									8		5	
Sodium, (mg/kg)	17.6	57.8	26.8	149	174	85.04	24.0	31.0	27.5	17.2	56.6	25.6	148	172.83	83.88	25.7	33.2	29.45
Potassium , mg/kg	1.39	4.57	2.10	11.8	13.7	6.71	1.90	2.45	2.18	1.36	4.48	2.02	11.6	13.6	6.61	2.24	2.92	2.58
Magnesium, mg/kg	7.04	23.1	10.7	59.6	69.3	33.95	9.60	12.2	10.90	6.88	22.6	10.3	58.8	68.8	8.33	10.8	14.0	12.4

Table 3. Heavy metal concentrations of surface soil samples in Izombe oil spills fields, 2024.

Element	Sampling Sites (0-15cm) Depth																	
	APRIL, 2024									JUNE, 2024								
	SL1	SL2	SL3	SL4	SL5	Mean	SL6	SL7	Mean	SL1	SL2	SL3	SL4	SL5	Mean	SL6	SL7	Mean
Manganese mg/kg	0.056	0.190	0.216	0.311	0.480	0.250	0.128	0.405	0.267	0.060	0.187	0.206	0.300	0.488	0.248	0.138	0.410	0.274
Zinc, mg/kg	0.073	0.162	0.216	0.436	0.336	0.245	0.080	0.116	0.098	0.070	0.170	0.214	0.428	0.340	0.244	0.078	0.124	0.101
Copper, mg/kg	0.150	0.227	0.144	0.220	0.208	0.190	0.152	0.146	0.149	0.148	0.226	0.140	0.215	0.206	0.187	0.158	0.153	0.156
Vanadium, mg/kg	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002
Iron, mg/kg	46.7	31.8	22.0	31.3	29.2	32.2	32.4	250	28.7	47.0	31.2	21.6	30.6	31.88	32.1	25.5	25.5	28.8
Cadmium, mg/kg	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002	<0.002
Lead, mg/kg	0.240	0.166	0.111	0.313	0.146	0.215	0.082	0.166	0.119	0.211	0.154	0.114	0.405	0.168	0.210	0.082	0.140	0.111

Effects of oil spillage on Total Hydrocarbon Content (THC) in the field sampled were found to be less than 0.01 in both April and June. This value according to EGASPIN (2018) is found to be low, hence below the critical limit, indicating that both plants and animals in terrestrial and aquatic habitats are safe. Available phosphorus (AV. P) in April was 1.81 showing an increase by 61% and 3.04 in June with an increase of 147% due to spillage effect. Low AV. P with crude oil contamination could be linked to microbes in the soils, which utilize petroleum hydrocarbons as a carbon source (Wang *et al.*, 2009; Li *et al.*, 2020). Generally, it was observed that the mean values for both AV. P and total nitrogen in the studied soils were low according to the ratings of Esu, (1991). Crude oil pollution encourages nutrient elements imbalance in the soils (Ihem *et al.*, 2015). Results showed that the exchangeable bases mean values were all within the fertility ratings of Esu (1991) and Babalola *et al.* (1998). The high values of these cations could be as a result of favourable mineralization and mobilization of these nutrients by microbes (Eneje and Ebomotie, 2011). However, the oil spillage increased the exchangeable cations at various periods under study. Calcium (Ca) was increased by 209% in April and 171% in June, Sodium (Na) increased by 208% in April and 156% in June and Mg increased by 211% in April and 170% in June.

Effects of oil spillage on the studied soils on heavy metal concentration are presented in Table 3. Higher values of heavy metals were recorded in polluted soils in the two months under study compared to the control in 2024.

According to Osuji and Onajake (2004), heavy metals in crude oil show significant mobility in polluted soils. oil spillage increased zinc (Zn) by 150% in April and 142% in June, copper (Cu) by 28% in April and 20% in June, lead (Pb) by 81% in April and 89% in June. However, all the heavy metals observed were below the permissible limits according to EGASPIN limits, Denneman and Robberse, (1990); Ministry of Housing Netherlands, (1994) and WHO, (1996). With this plausible result, it shows that soils of this area are not under threat perhaps for this short period.

CONCLUSION

This study revealed that the soils were negatively impacted by the oil spillage. Soils were rendered unproductive and less fertile with oil pollution based on the standard fertility ratings. Heavy metal concentrations were observed to be increased with oil pollution compared to the control sites, though the values were below the permissible limit. These results indicate that soils of the studied locations require proper attention hence application of soil amendments and adopting different measures of bioremediation. Precautionary measures should be taken and world standard procedures followed in crude oil processing and handling to prevent extensive and severe spillage into the soil and water environment. The findings revealed that there was no significant change in soil characteristics affected by the time possibly the duration was too short. It is recommended that the effect of spillage based on the duration should be extended.

ACKNOWLEDGMENT

The authors express sincere thanks to the Head of the Department of Agricultural Extension, Federal University of Technology, Owerri, Imo State, Nigeria for the facilities provided to carry out this research work.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

ETHICS APPROVAL

Not applicable

FUNDING

This study received no specific funding from public, commercial, or not-for-profit funding agencies.

AI TOOL DECLARATION

The authors declares that no AI and related tools are used to write the scientific content of this manuscript.

DATA AVAILABILITY

Data will be available on request

REFERENCES

- Abu-Hassanein, Z. S., Benson, C. H., & Blotz, L. R. (1996). Electrical resistivity of compacted clays. *Journal of Geotechnical Engineering*, 122(5), 397–406.
- Akinlo, A. E. (2012). How important is oil in Nigeria's economic growth? *Journal of Sustainable Development*, 5, 165–179.
- Akinpelu, R. (2021). An analysis of oil spill data in Nigeria from 2011–2021. *Nigerian Journal of Environmental Studies*, 19(2), 56–72.
- Al-Hamadani, Y. A., Salal, A. M., Mohammed, Z. B., Assi, L. N., & Resheq, A. S. (2022). Water quality assessment of water treatment plants using environmental and statistical indicators within Baghdad City. *Periodicals of Engineering and Natural Sciences*, 10, 427–436.
- APHA. (1992). *Standard methods for the examination of water and wastewater* (18th ed.). Washington, DC: American Public Health Association.
- Babalola, O., Bebaji, G. A., & Mustapha, S. (1998). Soil management for sustainable agriculture and environmental harmony. In *Proceedings of the 24th Annual Conference of the Soil Science Society of Nigeria* (pp. xx–xx). Bauchi, Nigeria: Abubakar Tafawa Balewa University.
- Bremner, J. M. (1996). Nitrogen-total. In D. L. Sparks (Ed.), *Methods of soil analysis: Part 3—Chemical methods* (pp. 1085–1121). Madison, WI: Soil Science Society of America.
- Chijioke, A., Nnaji, C., & Ugwu, F. (2018). Agricultural challenges in oil-polluted areas of the Niger Delta. *Nigerian Agricultural Journal*, 14(1), 34–49.
- Choo, G., Ekpe, O. D., Park, W. K., Chung, D., Lee, J., & Oh, J. (2022). Temporal and spatial trends of chlorinated paraffins and organophosphate flame retardants in black-tailed gull (*Larus crassirostris*) eggs. *Science of the Total Environment*, 803, 150137.
- Denneman, A. B., & Datsi, A. A. (2009). Ecotoxicological risk assessment as a base for development of soil quality criteria. In *Contaminated soil '90* (pp. 157–164). Dordrecht, Netherlands: Springer.
- Denneman, C. A., & Robberse, J. G. (1990). Ecotoxicological risk assessment as a base for development of soil quality criteria. In *Contaminated soil '90* (pp. 157–164). Dordrecht, Netherlands: Springer.
- EGASPIN. (2018). *Environmental guidelines and standards for the petroleum industry in Nigeria* (3rd ed.). Abuja, Nigeria: Nigerian Upstream Petroleum Regulatory Commission.
- Ejechi, B. (2022). Oil production and its effects on the Niger Delta ecology. *Nigerian Journal of Ecology*, 18(2), 21–38.
- Ekeghe, N. (2022). Replacing oil as the mainstay of the Nigerian economy. *Business News*, Nigeria.
- Eneje, R. C., & Ebomotei, E. (2011). Effect of crude oil pollution on soil physicochemical properties and germination percentage of *Amaranthus*. *Nigerian Journal of Soil and Environmental Research*, 9, 97–103.
- Esu, I. E. (1991). *Detailed soil survey of NIHORT farm at Bunkure, Kano State, Nigeria*. Zaria, Nigeria: Institute for Agricultural Research, Ahmadu Bello University.
- Federal Ministry of Environment (FMENV). (1991). *Guidelines and standards for environmental pollution control in Nigeria*. Lagos, Nigeria: FMENV.
- Gulshen, A. B., & Datsi, A. A. (2012). Role of soil texture and depth on the emergence of buried weed seeds. *ARPJ Journal of Agricultural and Biological Science*, 7, 223–228.
- Ihem, E. E., Osuji, G. E., Onweremadu, E. U., Uzoho, B. U., Nkwopara, U. N., Ahukemere, C. M., ... Okeli, N. H. (2015). Variability in selected properties of crude oil-polluted soils of Izombe, Northern Niger Delta, Nigeria. *International Journal of Applied Research*, 4(3), 29–33.
- Kujawinski, E. B., Reddy, C. M., & Rodgers, R. P. (2020). The first decade of scientific insights from the Deepwater Horizon oil release. *Nature Reviews Earth & Environment*, 1, 237–250.
- Li, Y., Zhao, Z., & Zhang, X. (2021). Nutrient availability and soil amendment effectiveness of various organic materials: A laboratory study. *Geoderma*, 380, 114703.

- Li, W., Yang, Y., Liu, Z. H., *et al.* (2020). Progression of mental health services during the COVID-19 outbreak in China. *International Journal of Biological Sciences*, 16, 1732–1738.
- Ministry of Housing, Netherlands. (1994). *Dutch intervention values for soil remediation* (Report HQ 94-021). The Hague, Netherlands: Ministry of Housing.
- Moneke, U., Ayo, O., & Chukwu, B. (2022). Health effects of oil pollution in the Niger Delta. *Journal of Environmental Health*, xx(x), xx–xx.
- Moral, R., Pérez-Murcia, M. D., Pérez-Espinosa, A., Moreno-Caselles, J., Paredes, C., & Rufete, B. (2008). Salinity, organic matter, and micronutrients in soils amended with organic wastes. *Bioresource Technology*, 99, 225–231.
- Nelson, D. W., & Sommers, L. E. (1982). Total carbon, organic carbon, and organic matter. In A. L. Page, R. H. Miller, & D. R. Keeney (Eds.), *Methods of soil analysis: Part 2* (pp. 539–579). Madison, WI: American Society of Agronomy.
- Onweremadu, E. U., & Peter, K. D. (2016). Pedogenesis of soils of two tropical microclimates in Owerri area, southeastern Nigeria. *International Journal of Soil Science*, 11, 14–18.
- Osuji, I. C., & Onakpoge, C. M. (2004). The Ebocha oil spillage II: Fate of associated heavy metals six months after. *Agricultural Journal and Environmental Assessment Management*, 9, 78–87.
- Owugah, L. (2006). *Oil transnationals, state and development in the oil-producing communities of the Niger Delta*. Accra, Ghana: Third World Network Africa.
- Wang, M., Jiang, J., Tan, S., Sun, Z., & Han, X. (2009). In situ ruminal crude protein and starch degradation of three classes of feedstuffs in goats. *Journal of Applied Animal Research*, 36, 23–28.
- Wang, X., Feng, J., & Wang, J. (2009). Petroleum hydrocarbon contamination and its impact on soil characteristics in an oil field of the Momoge wetland. *Environmental Science*, 30(8), 2394–2401. (In Chinese)
- Wokocha, G. A., Emeodu, D., & Ihenko, S. (2011). Impact of crude oil spillage on soil and food production in Rivers State, Nigeria. *Journal of Money, Investment and Banking*, 19, 29–34.
- World Health Organization. (1996). *Trace elements in human nutrition and health*. Geneva, Switzerland: WHO.