



Research Article

CHARACTERIZATION OF ZINC TOLERANT BACTERIAL STRAINS FROM THE ELECTROPLATING EFFLUENT CONTAMINATED SOIL

^{1*}Ranjithkumar M, ²Sathya P, and ²Mahalingam PU

¹Department of Microbiology, Syed Ammal Arts and Science College, Ramanathapuram- 623 513, Tamil Nadu.

²Department of Biology, Gandhigram Rural Institute-Deemed to be University, Dindigul-624 302, Tamil Nadu.

Article History: Received 27th December 2021; Accepted 07th January 2022; Published 14th January 2022

ABSTRACT

Toxic heavy metal pollution is expanding throughout the world as a result of industrial progress. This work focuses on the characterization of zinc tolerant bacterial strains from an electroplating effluent polluted soil sample in order to minimize/control metal pollution. pH, Temperature, Electrical Conductivity, Total Solids, Total Dissolved Solids, Total Suspended Solids, Chloride, Sodium, Calcium, Potassium, Biological Oxygen Demand, and Chemical Oxygen Demand were all measured and evaluated in the zinc-containing electroplating effluent sample. The sixteen bacterial strains were isolated from a polluted soil sample with electroplating effluent and identified using morphological and biochemical features. Using nutritional agar medium enriched with zinc metal, all of the chosen strains were evaluated for metal tolerance. Only six bacterial strains were chosen as potential metal tolerant strains based on the screening study, and these strains were characterized under various environmental conditions such as different pH (pH 5, pH 7, and pH 9), different temperatures (5°C, 28°C, 37°C, and 45°C), and different metal concentrations (100ppm, 200ppm, 300ppm and 400ppm). *Pseudomonas* sp strain 1 was shown to be a better zinc resistant organism, according to the findings.

Keywords: Electroplating effluents, heavy metals, Zinc, Biosorption, *Pseudomonas* sp strain 1.

INTRODUCTION

With the advancement of technology, pollution from toxic heavy metals is spreading around the world. The most common heavy metals released from industries such as electroplating, metal finishing, tanning, and so on are copper, cadmium, chromium, and nickel (Aksu, 1998; APHA, 1998; Avery & Tobin, 1993; Brady *et al.*, 1994). Toxic metals (Hg, Cr, Pb, Zn, Cu, Ni, Cd, As, Co, Sn), precious metals (Pd, Ag, Au, Ru), and radionuclides (Wang & Chen, 2006) are the three types of heavy metals that are of importance to the environment, according to (Wang & Chen, 2006) (U, Th, Ra, Am). Because of their toxicity and threat to human life and the environment, electroplating industries are a key contributor to the large influx of chromium into the biosphere (Dönmez & Aksu, 1999). As a result, effluent processes are designed to ensure that when wastewater is discharged into natural water sources, it meets varying effluent discharge criteria depending on where it is deposited, as well as reducing or preventing detrimental effects (Holt *et al.*, 1994).

Microbial biomass can operate as effective metal scavengers in both the bioaccumulation and biosorption processes. Bioaccumulation relies on metals being incorporated into living biomass, whereas biosorption is a metabolism-independent process in which metallic ions remain at the cellular surface. Complex formation, ion exchange, coordination, adsorption, and chelation are some of the mechanisms that cause biosorption (Hu *et al.*, 1996). In general, the mechanism has been relied on ion interaction at the biomass/aqueous medium interface (Huang *et al.*, 1990). Heavy metal traces are required as co-factors in enzyme reactions, but large quantities can be extremely harmful to living organisms due to metabolic reaction inhibition. Transport across the cell membrane, biosorption to the cell walls and trapping in extracellular capsules, precipitation, complexation, and oxidation-reduction reactions are some of the ways microorganisms respond to heavy metals (Dermentzis *et al.*, 2011; Jamaluddin *et al.*, 2012; Pandian *et al.*, 2014).

*Corresponding Author: Dr. M. Ranjithkumar, Assistant Professor and Head, Department of Microbiology, Syed Ammal Arts and Science College, Ramanathapuram- 623 513, Tamil Nadu, India, Email: ranjithmk85@gmail.com, Mobile: +91 96593 48393.

Zinc is a trace element that has a role in the formation of complexes (such as zinc fingers in DNA) and as a component of biological enzymes. Bacterial organisms collect zinc by an unspecific, quick absorption process, and it is typically found in larger amounts (but is less hazardous) than other heavy metals (Mahalingam *et al.*, 2014). Zinc ion uptake is often linked to magnesium ion uptake, and the two ions may be transported by comparable methods in bacteria (Mergey *et al.*, 1985). Some bacterial species have high tolerance and resistance to heavy metals in high concentrations. Because of the advent of resistant species (Nies, 1999), the reasons for microorganisms' resistance and tolerance behaviour require further investigation. As a result, in this study, an attempt was made to investigate zinc-tolerant bacterial strains.

MATERIALS AND METHODS

Collection of sample

The zinc electroplating effluent was collected from the direct outlet of Meena Electroplating Industry, Madurai, Tamil Nadu, India and the sample was immediately transported to the laboratory, Department of Biology, GRI, Gandhigram for further analysis.

Physicochemical characteristics of the electroplating effluent

The physicochemical parameters of electroplating effluents, such as pH, Temperature, Electrical Conductivity, Total Solids, Total Dissolved Solids, Chloride, Sodium, Calcium, Potassium, Biological Oxygen Demand, and Chemical Oxygen Demand, were measured using the method recommended by APHA, 1998 (Nies & Silver, 1995), and the results were compared to the maximum permissible limit prescribed by BIS for effluent standard (Pandit *et al.*, 2013).

Isolation and Identification of metal tolerant Bacterial strains

The soil sample contaminated by electroplating effluent was collected in pre-cleaned polythene bags and brought to the laboratory for bacterial strain isolation. The sample was diluted up to 10⁻⁹ dilutions with normal saline (0.85 percent NaCl). 10⁻⁵ and 10⁻⁶ dilution samples (0.1ml) were obtained and distributed on nutrient agar plates, which were then incubated at 37°C for 24 hours. The sixteen most common colonies were chosen and kept as pure cultures on nutrient agar medium. Gram's staining, Motility test, Indole production, Methyl Red reaction, Voges Proskauer reaction, Citrate utilisation, Urease reaction, Starch hydrolysis, and Catalase reaction (Demirbaş, 2003) were used to identify the sixteen bacterial strains.

Screening of selected bacterial isolate for heavy metal zinc resistance

All sixteen bacterial isolates were then tested for their ability to withstand heavy metal. A nutrient agar medium

with a pH of 7 was created and zinc metal was added (100ppm). The medium was sterilised at 121°C for 20 minutes with 15 lbs (Lyons & Genc, 2016; Rai *et al.*, 1981). All sixteen bacterial isolates were streaked on zinc-containing agar medium and cultured for five days at 37°C. All of the bacterial isolates' growth performance was observed and documented.

Characterization of zinc tolerant bacterial strains

Six bacterial species that may be metal (Zn) tolerant *Pseudomonas* spp., *Escherichia coli*, *Proteus* spp., *Staphylococcus* spp., *Salmonella* spp. 2 and *Shigella* spp. 2 were characterised by cultivating them for 5 days in a metal-based nutrient agar medium with varied environmental conditions such as pH (pH 5, pH 7 & pH 9), temperature (5°C, 28°C, 37°C & 45°)

RESULT AND DISCUSSION

The electroplating industry is a major user of water, it contains high amounts of heavy metal ions will be released. This heavy metal in industrial wastewaters is a serious concern because they are highly toxic, non-biodegradable, a carcinogen and continuous deposition into receiving water lakes, streams and other water sources within vicinity cause bioaccumulation in the living organisms (Sivasubramanian, 2006). Therefore, in the present work an attempt was made up to study on isolation and characterization of zinc tolerant bacterial isolates. In this present study, zinc containing electroplating effluent sample was collected from the direct outlet and analyzed for various physicochemical parameters and found to containing higher concentrations physicochemical components. The zinc containing electroplating effluent posses all the physicochemical components in higher concentration than the permissible level prescribed by BIS. The total bacterial population in the soil contaminated by electroplating effluent was counted. The sixteen most common bacterial strains were isolated and identified based on their morphological and biochemical properties. Bergey's Manual of Determinative Bacteriology was used to verify all of the strains. Tables 1 and 2 show the characteristics of gramme positive (5 strains) and gramme negative (11 strains) bacterial isolates. Metal-tolerant bacterial species such as *Staphylococcus* sp, *Bacillus* sp, and *Pseudomonas aeruginosa* have been found in electroplating effluent contaminated soil samples by several authors (Hansford & Vargas, 1999; Sugumaran *et al.*, 2014; Veglio & Beolchini, 1997).

By growing all sixteen bacterial isolates in nutritional agar medium containing zinc metal, they were tested for their ability to withstand heavy metal, and the results were documented in Table 3. Only six bacterial strains, *Pseudomonas* spp. 1, *Escherichia coli*, *Proteus* spp. 2, *Staphylococcus* spp. 1, *Salmonella* spp. 2, and *Shigella* spp. 2, were identified as prospective metal tolerant strains based on their growth performance in the screening medium (Table 3). According to (Madigan & Pidcoo, 2003), the majority of metal-tolerant bacterial strains

belong to the Gram negative bacterial group rather than the Gram positive bacterial group. This is due to the gramme negative cell wall's composition and interaction with certain metal ions. Some Gram-positive strains, on the other hand, have been demonstrated to be extremely resistant to high concentrations of heavy metals.

Table 1. Morphological and biochemical characteristics of Gram negative bacterial isolates.


| Bacterial isolates No | Cultural characteristics | Biochemical characteristics | | | | | | | | | | | Name of the bacterial strain | |
|-----------------------|--------------------------|-----------------------------|---------------|-----------------|-------------------|---------------------|--------------------------|-------------------|-------------------|--------------------------|-----------------|----------------------|------------------------------|-----------------------------|
| | | Simple staining | Motility test | Gram's reaction | Indole production | Methyl red reaction | Voges Proskauer reaction | Catalase activity | Starch hydrolysis | Citrate utilization test | Urease reaction | Glucose fermentation | | Gelatin liquefaction |
| BIS-1 | WC | C | M | + | + | - | - | + | + | + | - | - | - | <i>Staphylococcus</i> spp 1 |
| BIS-2 | GBC | C | M | + | + | - | - | + | + | + | - | - | - | <i>Staphylococcus</i> spp 2 |
| BIS-3 | LYMC | R | M | + | - | - | + | - | - | - | - | + | - | <i>Bacillus</i> spp 1 |
| BIS-4 | WWC | R | M | + | - | - | + | - | - | - | - | + | - | <i>Bacillus</i> spp 2 |
| BIS-5 | BMC | C | M | + | + | + | - | - | - | + | - | + | + | <i>Micrococcus</i> sp |

BIS - Bacterial Isolate, WC - Whitish Colony, GBC - Golden Brown Colony, LYMC - Light Yellow Mucoïd Colony, WWC - White Waxy Colony, BMC - Brown color Mucoïd Colony, C - Cocci, R - Rod, M - Motile, + - Positive, - - Negative.

Table 2. Morphological and biochemical characteristics of Gram positive bacterial isolates.

| Bacterial isolates No | Cultural characteristics | Biochemical characteristics | | | | | | | | | | | Name of the bacterial strain | |
|-----------------------|--------------------------|-----------------------------|---------------|-----------------|-------------------|---------------------|--------------------------|-------------------|-------------------|--------------------------|-----------------|----------------------|------------------------------|--------------------------|
| | | Simple staining | Motility test | Gram's reaction | Indole production | Methyl red reaction | Voges Proskauer reaction | Catalase activity | Starch hydrolysis | Citrate utilization test | Urease reaction | Glucose fermentation | | Gelatin liquefaction |
| BIS-6 | OSC | R | M | - | - | - | - | + | - | + | - | + | + | <i>Pseudomonas</i> spp 1 |
| BIS-7 | GMC | R | M | - | - | - | - | + | - | + | - | + | + | <i>Pseudomonas</i> spp 2 |
| BIS-8 | DWC | R | M | - | - | - | - | + | - | + | - | + | + | <i>Pseudomonas</i> spp 3 |
| BIS-9 | OCC | R | M | - | - | - | - | + | - | + | - | + | + | <i>Pseudomonas</i> spp 4 |
| BIS-10 | CMC | R | M | - | + | + | - | + | - | - | - | + | - | <i>Escherichia coli</i> |
| BIS-11 | TGC | R | M | - | - | - | + | + | + | - | - | + | - | <i>Proteus</i> spp 1 |
| BIS-12 | WYC | R | M | - | - | - | + | + | + | - | - | + | - | <i>Proteus</i> spp 2 |
| BIS-13 | WPC | R | M | - | - | + | - | + | - | - | - | + | + | <i>Salmonella</i> spp 1 |
| BIS-14 | WOC | R | M | - | - | + | - | + | - | - | - | + | + | <i>Salmonella</i> spp 2 |
| BIS-15 | YC | R | M | - | - | - | - | - | + | - | - | - | - | <i>Shigella</i> spp 1 |
| BIS-16 | WMC | R | M | - | + | - | - | - | + | - | - | - | - | <i>Shigella</i> spp 2 |

BIS - Bacterial Isolate, OSC - Orange, Small Colony, GMC - Green color Mucoïd Colony, DWC - Dusty White Colony, OCC - Orange color Creamy Colony, CMC - Colorless Mucoïd Colony, TGC - Thin, Grey Colony, WYC - Whitish Yellow Colony, WPC - Whitish Pink Colony, WOC - Whitish Orange Colony, YC - Yellow Colony, WMC - Whitish Mucoïd Colony, C - Cocci, R - Rod, M - Motile, + - Positive, - - Negative

Figure 1. Growth performance of 16 bacterial isolates in zinc containing nutrient agar medium on 5 days.

- | | | | |
|--------------------------------|-----------------------------|-----------------------------|-----------------------------|
| 1. <i>Staphylococcus</i> spp 1 | 5. <i>Micrococcus</i> spp | 9. <i>Pseudomonas</i> spp 4 | 13. <i>Salmonella</i> spp 1 |
| 2. <i>Staphylococcus</i> spp 2 | 6. <i>Pseudomonas</i> spp 1 | 10. <i>Escherichia coli</i> | 14. <i>Salmonella</i> spp 2 |
| 3. <i>Bacillus</i> spp 1 | 7. <i>Pseudomonas</i> spp 2 | 11. <i>Proteus</i> spp 1 | 15. <i>Shigella</i> spp 1 |
| 4. <i>Bacillus</i> spp 2 | 8. <i>Pseudomonas</i> spp 3 | 12. <i>Proteus</i> spp 2 | 16. <i>Shigella</i> spp 2 |

Table 3. Growth performance of 16 bacterial isolates in zinc containing nutrient agar medium on 5 days.

| Bacterial Isolate No. | Strain Type | Name of the Isolates | Growth Characterization |
|-----------------------|-------------|-----------------------------|-------------------------|
| BIS-1 | + ve | <i>Staphylococcus</i> spp 1 | GG |
| BIS-2 | + ve | <i>Staphylococcus</i> spp 2 | MG |
| BIS-3 | + ve | <i>Bacillus</i> spp 1 | PG |
| BIS-4 | + ve | <i>Bacillus</i> spp 2 | GG |
| BIS-5 | + ve | <i>Micrococcus</i> sp | MG |
| BIS-6 | - ve | <i>Pseudomonas</i> spp 1 | EG |
| BIS-7 | - ve | <i>Pseudomonas</i> spp 2 | MG |
| BIS-8 | - ve | <i>Pseudomonas</i> spp 3 | PG |
| BIS-9 | - ve | <i>Pseudomonas</i> spp 4 | MG |
| BIS-10 | - ve | <i>Escherichia coli</i> | GG |
| BIS-11 | - ve | <i>Proteus</i> spp 1 | PG |
| BIS-12 | - ve | <i>Proteus</i> spp 2 | GG |
| BIS-13 | - ve | <i>Salmonella</i> spp 1 | PG |
| BIS-14 | - ve | <i>Salmonella</i> spp 2 | GG |
| BIS-15 | - ve | <i>Shigella</i> spp 1 | MG |
| BIS-16 | - ve | <i>Shigella</i> spp 2 | GG |

BIS - Bacterial Isolates, EG - Excellent Growth, GG - Good Growth , MG - Moderate Growth ,PG - Poor Growth.

Six potential bacterial isolates, *Pseudomonas* spp. 1, *Escherichia coli*, *Proteus* spp. 2, *Staphylococcus* spp. 1, *Salmonella* spp. 2 and *Shigella* spp. 2, were characterised by growing them for 5 days in zinc-containing nutrient agar medium with various environmental conditions such as pH (pH 5, pH 7 & pH 9), temperature (5°C, 28°C, 37°)Table 4 contains all of the results.

Table 4. Growth performance of six potential bacterial isolates in zinc enriched nutrient agar medium with various environmental conditions and metal concentration.

| pH | Bacterial Isolates | Metal concentration (ppm) | | | | | | | | | | | | | | | |
|----|-----------------------------|---------------------------|----|----|----|-----|----|----|----|-----|----|----|----|-----|----|----|----|
| | | 100 | | | | 200 | | | | 300 | | | | 400 | | | |
| | | Temperature (°C) | | | | | | | | | | | | | | | |
| | | 5 | 28 | 37 | 45 | 5 | 28 | 37 | 45 | 5 | 28 | 37 | 45 | 5 | 28 | 37 | 45 |
| 5 | <i>Pseudomonas</i> spp 1 | NG | PG | GG | NG | NG | MG | GG | NG | NG | MG | GG | PG | NG | PG | GG | MG |
| | <i>Staphylococcus</i> spp 1 | NG | MG | MG | NG | NG | PG | MG | PG | NG | MG | GG | MG | NG | PG | PG | NG |
| | <i>Escherichia coli</i> | NG | PG | GG | NG | NG | MG | GG | MG | NG | PG | MG | PG | NG | PG | MG | PG |
| | <i>Proteus</i> spp 2 | NG | PG | MG | PG | NG | MG | MG | MG | NG | PG | MG | PG | NG | PG | PG | MG |
| | <i>Salmonella</i> spp 1 | NG | PG | MG | MG | NG | MG | EG | MG | NG | MG | PG | NG | NG | MG | MG | MG |
| | <i>Shigella</i> spp 2 | NG | MG | MG | NG | NG | PG | MG | PG | NG | MG | MG | PG | NG | MG | PG | PG |
| 7 | <i>Pseudomonas</i> spp 1 | NG | MG | GG | PG | NG | MG | GG | PG | NG | GG | EG | PG | NG | MG | GG | NG |
| | <i>Staphylococcus</i> spp 1 | NG | PG | MG | MG | NG | MG | PG | NG | NG | PG | MG | PG | NG | NG | MG | PG |
| | <i>Escherichia coli</i> | NG | MG | GG | PG | NG | MG | MG | MG | NG | MG | GG | PG | NG | PG | MG | PG |
| | <i>Proteus</i> spp 2 | NG | MG | MG | NG | NG | MG | MG | PG | NG | MG | MG | MG | NG | PG | PG | MG |
| | <i>Salmonella</i> spp 1 | NG | PG | GG | PG | NG | PG | PG | NG | NG | PG | PG | PG | NG | PG | PG | PG |
| | <i>Shigella</i> spp 2 | NG | MG | MG | NG | NG | PG | MG | NG | NG | PG | MG | PG | NG | MG | PG | PG |
| 9 | <i>Pseudomonas</i> spp 1 | NG | MG | MG | PG | NG | MG | MG | PG | NG | MG | PG | PG | NG | NG | PG | NG |
| | <i>Staphylococcus</i> spp 1 | NG | PG | MG | NG | NG | PG | MG | NG | NG | NG | PG | NG | NG | PG | PG | NG |
| | <i>Escherichia coli</i> | NG | PG | PG | NG | NG | PG | PG | PG | NG | NG | PG | PG | NG | PG | NG | NG |
| | <i>Proteus</i> spp 2 | NG | PG | MG | NG | NG | PG | MG | NG | NG | PG | NG | NG | NG | NG | NG | NG |
| | <i>Salmonella</i> spp 1 | NG | PG | PG | PG | NG | MG | PG | PG | NG | PG | NG | NG | NG | NG | NG | NG |
| | <i>Shigella</i> spp 2 | NG | MG | MG | NG | NG | PG | MG | PG | NG | PG | PG | NG | NG | NG | NG | NG |

EG - Excellent Growth, GG - Good Growth, MG - Moderate Growth, PG - Poor Growth, NG - No Growth.

Pseudomonas spp strain 1 was identified as a potential zinc tolerant organism by displaying superior growth in zinc containing medium with pH 7, temperature 37°C, and 300 ppm concentration on the fifth day (Table 4). pH, temperature, and moisture all have an impact on microbial growth and activity. Although microorganisms have been isolated from extreme environments, most of them grow optimally across a restricted range, therefore achieving optimal circumstances is critical. Various environmental conditions were used to cultivate *Staphylococcus* spp. in chrome electroplating effluent medium. *Staphylococcus* spp grew best in a 20% electroplating effluent sample with pH 7 at 30°C, according to the findings.

CONCLUSION

The investigation on the isolation and characterisation of zinc-tolerant bacterial strains indicated that *Pseudomonas* sp 1 is a potential metal-tolerant strain that would be a better alternative for removing zinc metals from electroplating effluents before they are released onto agricultural land/water bodies. As a result, it reduces water and soil contamination.

ACKNOWLEDGMENT

The authors wish to thank the Meena Electroplating Industry, Madurai, Tamil Nadu, India and also Syed Ammal Arts and Science College, Ramanathapuram for supporting this study.

REFERENCES

- Aksu, Z. (1998). Biosorption of heavy metals by microalgae in batch and continuous systems *Wastewater Treatment with Algae*. 37-53.
- APHA (1998). Standard methods for the examination of water and wastewater (Vol. 2): American Public Health Association.
- Avery, S. V., & Tobin, J. M. (1993). Mechanism of adsorption of hard and soft metal ions to *Saccharomyces cerevisiae* and influence of hard and soft anions. *Applied and Environmental Microbiology*, 59(9), 2851-2856.
- Brady, D., Stoll, A., Starke, L., & Duncan, J. (1994). Chemical and enzymatic extraction of heavy metal binding polymers from isolated cell walls of *Saccharomyces cerevisiae*. *Biotechnology and Bioengineering*, 44(3), 297-302.
- Demirbaş, E. (2003). Adsorption of cobalt (II) ions from aqueous solution onto activated carbon prepared from hazelnut shells. *Adsorption Science & Technology*, 21(10), 951-963.
- Dermentzis, K., Christoforidis, A., & Valsamidou, E. (2011). Removal of nickel, copper, zinc and chromium from synthetic and industrial wastewater by electrocoagulation. *International Journal of Environmental Sciences*, 1(5), 697-710.

- Dönmez, G., & Aksu, Z. (1999). The effect of copper (II) ions on the growth and bioaccumulation properties of some yeasts. *Process Biochemistry*, 35(1-2), 135-142.
- Hansford, G., & Vargas, T. (1999). Chemical and electrochemical basis of bioleaching processes *Process Metallurgy*, (9)13-26.
- Holt, J. G., Krieg, N. R., Sneath, P. H., Staley, J. T., & Williams, S. T. (1994). *Bergey's manual of determinative bacteriology*. 9th. *Baltimore: William & Wilkins*.786-788.
- Hu, M. Z. C., Norman, J. M., Faison, B. D., & Reeves, M. E. (1996). Biosorption of uranium by *Pseudomonas aeruginosa* strain CSU: characterization and comparison studies. *Biotechnology and Bioengineering*, 51(2), 237-247.
- Huang, C.-p., Huang, C.-p., & Morehart, A. L. (1990). The removal of Cu (II) from dilute aqueous solutions by *Saccharomyces cerevisiae*. *Water Research*, 24(4), 433-439.
- Jamaluddin, H., Zaki, D. M., & Ibrahim, Z. (2012). Isolation of metal tolerant bacteria from polluted wastewater. *Pertanika Journal of Tropical Agricultural Science*, 35(3).
- Lyons, G., & Genc, Y. (2016). Commercial humates in agriculture: real substance or smoke and mirrors? *Agronomy*, 6(4), 50.
- Madigan, M. L., & Pidcoe, P. E. (2003). Changes in landing biomechanics during a fatiguing landing activity. *Journal of Electromyography and Kinesiology*, 13(5), 491-498.
- Mahalingam, P., Ranjithkumar, M., & Ramalakshmi, P. (2014). Partial characterization of metal resistant *Bacillus* spp. isolated from electroplating effluent sediments. *Journal of Microbiology and Biotechnology Research*, 4(5), 43-45.
- Mergeay, M., Nies, D., Schlegel, H., Gerits, J., Charles, P., & Van Gijsegem, F. (1985). *Alcaligenes eutrophus* CH34 is a facultative chemolithotroph with plasmid-bound resistance to heavy metals. *Journal of Bacteriology*, 162(1), 328-334.
- Nies, D. H. (1999). Microbial heavy-metal resistance. *Applied Microbiology and Biotechnology*, 51(6), 730-750.
- Nies, D. H., & Silver, S. (1995). Ion efflux systems involved in bacterial metal resistances. *Journal of Industrial Microbiology*, 14(2), 186-199.
- Pandian, K., Thatheyus, A., & Ramya, D. (2014). Bioremoval of chromium, nickel and zinc in electroplating effluent by *Pseudomonas aeruginosa*. *Open Journal of Water Pollution and Treatment*, 2014(2), 75-82.
- Pandit, R., Patel, B., Kunjadia, P., & Nagee, A. (2013). Isolation, characterization and molecular identification of heavy metal resistant bacteria from industrial effluents, Amala-khadi-Ankleshwar, Gujarat. *International Journal of Environmental Sciences*, 3(5), 1689-1699.
- Rai, L., Gaur, J., & Kumar, H. (1981). Phycology and heavy-metal pollution. *Biological Reviews*, 56(2), 99-151.
- Sivasubramanian, V. (2006). Phycoremediation-Issues and challenges. *Indian Hydrobiology*, 9(1), 13-22.
- Sugumaran, T., Ramu, A., & Kannan, N. (2014). Journal homepage: <http://www.journalijar.com> *International Journal of Advanced Research Research Article*.2(4), 610-624.
- Veglio, F., & Beolchini, F. (1997). Removal of metals by biosorption: a review. *Hydrometallurgy*, 44(3), 301-316.
- Wang, J., & Chen, C. (2006). Biosorption of heavy metals by *Saccharomyces cerevisiae*: a review. *Biotechnology Advances*, 24(5), 427-451.