

CHARACTERIZATION OF WATERMELON (*CITRULLUS LANATUS*) INSECT PESTS AND THE DIFFERENT CONTROL METHODS USED IN SOUTHERN BENIN

^{*1,3,7} Akotegnon Azonwakin Rodrigue, ^{1,2}Faton Manhognon Oscar Euloge, ^{1,2,3}Bonou-Gbo Zaki,
^{1,2}Nouhouégnon Rock, ¹Kpetehoto Hinnoutondji Wilfrid, ^{1,5}Lucie Fanou, ^{1,2}Gnancadja-Andre
Léopold Simplicie

¹Laboratory for Research in Biological Resources/UATM GASA-Formation/Benin

²Laboratory of Plant Physiology and Study of Environmental Stresses: Research Unit in Phytopathology and Plant
Protection/ UAC/ FAST/ Benin

³ Nutritional Sciences Laboratory, Department of Food Sciences and Nutrition, Guinean Higher School of Tourism and
Hospitality, Conakry, Guinea

⁴ Laboratory of pharmacology and Improved Traditional Medicines, Department of Animal Physiology, Faculty of Science
and Technology, University of Abomey-Calavi, BP 526 Cotonou, Republic of Benin

⁵Department of Animal Sciences, Higher Institute of Agronomy and Veterinary of Faranah/Guinea

⁶Laboratory of Biotechnology, Genetic Resources and Plant and Animal Breeding / UAC / FAST /Benin

⁷Laboratory of Food Technology and Quality Control/UATM GASA-Formation/Benin

⁸Higher Institute of Sustainable Development, University of Yembila Abdoulaye Toguyeni/Burkina Faso

Article History: Received 22nd March 2025; Accepted 29th April 2025; Published 31st May 2025

ABSTRACT

The majority of watermelon pests have long been controlled by the often-excessive use of pesticides, whose harmful effects on ecosystems are well known. These impacts are all the more worrying when they affect crops. In light of this, integrated pest management (IPM) appears to be a sustainable alternative aimed at limiting the use of chemicals. This study aims to improve the productivity of *Citrullus lanatus* in southern Benin through integrated pest management techniques. It focuses on the classification of pests and diseases encountered, as well as the evaluation of the control methods used. The survey was conducted in several municipalities, including Missérété, Covè, Za-kpota, Comè, Dangbo, Dogbo, Lokossa, Tori-Bossito, Allada, Adjohoun, Kétou, Grand-Popo, and Ouidah. The results reveal the presence of flies, beetles, root-knot nematodes, downy mildew, leaf spot, anthracnose, fusarium wilt, and mosaic virus. Growers mainly use chemicals such as Pacha 25EC, Acarius 018EC, Emacot019EC, Kaobtimal, and Lambda. However, their overuse leads to the destruction of beneficials and pollution. Few studies mention the use of biopesticides, although various cultural practices can effectively control these pests.

Keywords: Insect pests, Biological control, Chemical control, Integrated control, Watermelon.

INTRODUCTION

Agriculture constitutes an essential foundation of the Beninese economy, representing 27% of the national GDP and employing over 70% of the working population, primarily in rural areas (MAEP, 2019). Market gardening, driven by growing urban demand, offers regular and rapid income, particularly attractive to young people and women

(Yarou *et al.*, 2015). Among these, watermelon (*Citrullus lanatus*) stands out for its nutritional qualities and strong economic potential. Native to Africa, this cucurbit is widely cultivated in southern Benin, particularly in Sèmè-Kpodji, Grand-Popo, and Ouidah, where the agroclimatic conditions are favorable (Agbangla *et al.*, 2017). However, its production is hampered by significant phytosanitary

challenges, linked to a wide variety of insect pests that affect leaves, stems and fruits, reducing yields and crop quality (Adandonon *et al.*, 2009; FAO, 2020). The main pests include aphids (*Aphis gossypii*), thrips (*Thrips tabaci*), whiteflies (*Bemisia tabaci*), leafminers (*Liriomyza* spp.) and fruit moths (*Diaphania indica*), often vectors of serious viral diseases such as watermelon mosaic (Sikirou *et al.*, 2013; Atachi *et al.*, 2016). Their proliferation is facilitated by inappropriate agricultural practices, such as monoculture, lack of crop rotation and irrational use of pesticides (Houunkpèvi *et al.*, 2020). Producers' response to these threats relies largely on synthetic insecticides. Although effective in the short term, their overuse generates resistance among pests, destroys natural beneficial organisms, and disrupts ecological balances (Ryckewaert, 1998; Tamo *et al.*, 2014). Furthermore, soil and water contamination, as well as the presence of chemical residues in fruit, pose a major risk to human health (Yarou *et al.*, 2015). In response, research institutions and NGOs recommend alternative methods such as integrated pest management, which combines sustainable cultural, biological, and chemical techniques, or biological control through the use of natural enemies (Goudegnon *et al.*, 2012).

In this context, this study aims to identify the main insect pests of watermelon in southern Benin and to evaluate producers' phytosanitary practices. It focuses on the municipalities of Sèmè-Kpodji, Abomey-Calavi, Ouidah and Cotonou. The specific objectives are: to identify the dominant pest species, analyze their mode of attack, determine the critical periods of their appearance, and evaluate the control strategies in use. The study is based on a combined entomological, agronomic and sociotechnical approach, in order to propose ecologically sustainable, economically viable and technically adapted solutions for local producers (Ryckewaert, 1998; Tamo *et al.*, 2014; Goudegnon *et al.*, 2012).

STUDY MATERIALS AND METHODS

Study Area

The Republic of Benin is located in West Africa between latitudes 6°10' N and 12°25' N and longitudes 0°45' E and 3°55' E (Akoègninou *et al.* 2006). It covers a total area of 112,622 km². The study was conducted in the southern part of the country, characterized by a relatively humid agro-ecological zone with two rainy seasons and annual rainfall ranging from 1100 to 1400 mm/year. Average annual temperatures range from 26°C to 28°C, and the vegetation types are semi-deciduous forests or forests or forests and savannah forests (Gbaguidi *et al.* 2013; Yabi and Afouda 2012). The surveys were conducted in several municipalities, including Missérété, Covè, Za-kpota, Comè, Dangbo, Dogbo, Lokossa, Tori-Bossito, Allada, Adjohoun, Kétou, Grand-Popo, and Ouidah.

Study Materials

Watermelon (*Citrullus lanatus*) is an annual, monoecious herbaceous plant with a creeping or climbing habit, reaching 4 to 10 meters in length. It clings to the ground using simple tendrils (Photo 1) and develops a superficial root system composed of a taproot accompanied by numerous lateral roots. Its ribbed stem is covered with long, soft hairs (E.O. Kichah, 2016). Its leaves are simple, alternate, without stipules, and borne on a 2 to 14 cm petiole that is also hairy. They have lobes with weakly toothed edges and long, hairy veins. The flowering, axillary and solitary, produces yellow, unisexual flowers, measuring 2 to 3.5 cm in diameter, with a campanulate calyx and petals fused at the base. Male flowers have three free stamens, while female flowers have an inferior ovary (E.O. Kichah, 2016). The fruit, a large globose to ellipsoidal berry, measures 5 to 70 cm long and can weigh between 1.5 and 30 kg. Its epidermis varies from white to green, gray or yellow, plain, spotted or striated. The flesh, juicy and sweet, can be white, pale green, yellow or red. The seeds, numerous, are elliptical, flat, smooth, 0.5 to 1.5 cm, yellow, brown, black or rarely white. Germination is epigeal, with rounded to oblong leaf-like cotyledons (Y.I. Pawendkisgou, 2021).

Study Methods

To achieve the objectives set out in this study, the methodology employed is based on three stages: the division and categorization of the study area, a document review, and data collection and processing.

Division and Categorization of the Study Area

The selection of the study areas was based on an in-depth document review and data from the Territorial Agricultural Development Agencies (ATDA). On this basis, watermelon-producing municipalities were divided into three categories: low-production municipalities ("tailing municipalities"), medium-production municipalities, and high-production municipalities ("leading municipalities"). To ensure balanced representation of the different production dynamics, a random sample was conducted, including the following municipalities:

- **Tailing municipalities:** Bonou, Covè, Za-Kpota, Zogbodomey ;
- **Medium-production municipalities:** Aguégoués, Athiémè, Comè, Dangbo, Djidja, Dogbo, Lokossa, Tori-Bossito, Zè ;
- **Leading municipalities:** Abomey-Calavi, Adja-Ouèrè, Adjohoun.

The main selection criterion was the proven presence of active watermelon farms in these localities (ATDA, 2023).

Literature review

This step allowed for the collection, analysis, and synthesis of existing data on watermelon production in Benin, particularly on insect pests and diseases affecting the crop.

It provided an essential theoretical and empirical framework to guide the rest of the study (Yarou *et al.*, 2017; FAO, 2020).

Data collection and processing

With the help of ATDA, a targeted survey was conducted in the thirteen identified municipalities to identify producers actually active in the watermelon sector. The data collected was structured by department, municipality, number of villages visited, and number of producers

surveyed. This phase provided a reliable basis for subsequent analyses.

RESULTS AND DISCUSSION

The results reflecting the watermelon production targets and customer types are shown in Table 1. The primary production objective is sales, and the potential customers are consumers and wholesalers (Table 1). The graphical representation of the data in Table 1 is shown in Figure 1 opposite.

Table 1. Production Target and Customer Types.

Designation	Terms and conditions	Rate
Production target	Consumption	4.17
	Sales	95,83
Types of customers	Consumers	100,00
	Wholesalers	95,83
	Transformers	2,78

Source : Field surveys, March 2023.




Figure 1. Production and Customer Graph.

Source. Field Surveys, March 2023.

The graph in Figure 1 from Table 1 presents the production objectives and customer types received by watermelon producers in southern Benin. It should be noted that 95.93% of producers have a production objective for sale, compared to 4.17% for consumption. Regarding customer types, 100% of watermelon producers deliver part of their production to consumers, 95.83% to wholesalers, and 2.78% to processors. These results fully justify the availability of watermelon customers in the study area. Watermelon is susceptible to various fungal diseases such as downy mildew, leaf spot, and powdery mildew, as well as mosaic virus. The insect pests are thrips (*Thrips* spp.), mites (*Tretranychus* spp.), aphids (*Aphisgossypii*), fruit

flies (*Dacusciliatus*), cucumber beetles (*Diabrotica* spp.), pumpkin leaf beetles (*Aulacophara* spp.), looper caterpillars, ladybugs (*Epilachna*), and leaf miners (*Liriomyza* spp.). The results relating to the distribution of watermelon pests and diseases in the thirteen municipalities receiving the surveys in southern Benin are presented in Table 2 below. Table 3 shows the presence of watermelon pests and/or diseases in the thirteen municipalities in southern Benin considered without exception. This represents a widespread, almost uniform distribution of watermelon pests and/or diseases. The survey conducted on the distribution of watermelon pests and diseases in southern Benin reveals that the pests that attack watermelon

crops are flies, beetles, and root-knot nematodes, and the diseases that commonly affect them are downy mildew, leaf spot, anthracnose, fusarium wilt, and mosaic virus. Figure 2

opposite presents the schematic distribution of watermelon pests and diseases in the municipalities receiving the surveys.

Tableau 2. Distribution of watermelon pests and diseases in the thirteen affected communes of southern Benin.

Municipalite s	Fl y	Milde w	Coleopter a	Mosai c	Anthracnos e	Nematode s	Fusarium wilt	Cladoporosi s	Effectiv e
Adjohoun	1	1	1	0	0	0	0	0	3
Allada	0	0	0	1	1	1	0	0	3
Comé	0	1	0	0	0	1	1	0	3
Covè	0	1	0	1	0	0	0	0	2
Dangbo	1	0	1	1	0	0	0	0	3
Dogbo	1	1	0	0	0	1	0	1	4
Grand-Popo	0	1	0	1	1	1	0	0	4
Kétou	1	1	1	1	0	0	0	0	4
Lokossa	1	1	0	1	0	1	0	0	4
Missrété	0	1	0	1	0	1	0	0	3
Ouidah	0	1	0	1	0	1	0	0	3
Tori-Bossito	1	1	0	0	1	1	0	0	4
Zakpota	1	1	0	1	0	0	0	0	3

Source : Field surveys, March 2023.

Note: The numbers 0 and 1 indicate the absence and presence of the watermelon pest or disease in the municipality, respectively.




Figure 2. Distribution of watermelon pests and diseases across municipalities. **Source:** Field surveys, March 2023.

Classification, description of affected organs, and types of losses caused by watermelon pests and diseases. Tables 3 to 4 present the classification, description of affected organs, and types of losses caused by watermelon pests and diseases.

Table 3. Classification and description of defoliating beetles and some watermelon flies.

INSECTS					
Affected organs		Types of losses			
Leaves	Fruits	Number of plants	Number of fruits per plant	Fruit size	Quality of ripe fruit
Defoliating beetles: <i>Chrysomelidae</i> (<i>Aulacophora africana</i>, <i>Monolepta spp.</i>, <i>Acalymma vittata</i>, <i>Diabrotica undecimpunctata</i>, <i>Asbecesta cyanipennis</i>, <i>Asbecesta transversa</i>, <i>Podagricra spp.</i>)					
Leaf perforation by adults. Larvae can also attack the stem at the collar and the roots.	Perforation of the skin due to feeding by adult beetles.	Death of young plants in the event of a strong attack			Decrease in market value, export impossible.
Fruit flies: <i>Dacus vertebratus</i>; <i>Dacus ciliatus</i>; <i>Bactrocera cucurbitae</i>; <i>Bactrocera invadens</i>; <i>Bactrocera latifrons</i>; <i>Bactrocera zonata</i>					
	Maggots thrive in fruitv		Sharp decrease if fruits are attacked at an early stage.	Fruit rot. Infested fruits lose all value..	

Source: Field surveys, March 2024

Table 4. Classification and description of whiteflies and some fungi.

INSECTS (continued)					
Affected organs		Types of losses			
Leaves	Fruits	Number of plants	Number of fruits per plant	Fruit size	Quality of ripe fruit
Whiteflies: <i>Bemisia tabaci</i>					
Adults and larvae feed on leaves.			Decrease, due to a reduction in photosynthesis following the development of sooty mold on the honeydew secreted by whiteflies.		Honeydew decreases the market value of fruits, due to the reduction in aesthetic value.
MUSHROOMS					
Fusarium wilt: <i>Fusarium oxysporum</i>, <i>Fusarium sp.</i> ; <i>Fusarium nivium</i>					
The mycelium grows in the stem		Death of young plants due to rot or death of older plants due to successive wilting.			If the fruit is already formed, it stops growing and loses all commercial value.

Source: Field surveys, March 2024

Tableau 5. Classification and description of some watermelon fungi.

MUSHROOMS (continued)								
Affected organs		Number of plants	Number of fruits per plant	Types of losses	Quality of ripe fruit			
Leaves	Fruits			Fruit size				
Anthracnose: <i>Collectotrichum orbiculare</i>; <i>Collectotrichum lagenarium</i>								
Mycelium development on the stem, leaves and fruits.				Even if the infection does not reach the flesh, secondary rot can develop and cause significant damage. Infected fruits lose all commercial value.				
Downy mildew: <i>Pseudoperonospora cubensis</i>								
Development on the leaves.		Premature death of plants in case of heavy infection.		Decrease due to leaf loss.	The fruits are rarely directly affected, but may remain small and tasteless. Such fruits cannot be exported.			
Cladosporiose : <i>Cladosporium cucumerinum</i>								
Development on leaves, leaf petioles, stems and fruits.				Les fruits comportant des cavités perdent toute valeur commerciale.				
Source: Field surveys, March 2024.								

Tableau 6. Classification and description of watermelon viruses.

VIRUS				
Affected organs		Types of losses		
Plante entière	Number of plants	Number of fruits per plant	Fruit size	Quality of ripe fruit
Watermelon mosaic virus: (Potyviridae: Potyvus)				
Viruses are transmitted from plant to plant by aphids (e.g. Aphis gossypii; Myzus persicae etc.). Aphids are infected with the virus present in an infected plant and then transmit it to a healthy plant through their bites. They can transmit the virus after feeding for 9 seconds. These viruses are not carried by seeds.				
Once transmitted, the virus spreads throughout the plant		Significantly reduced growth if plants are attacked during the early stages before flowering.	Diseased fruits cannot be exported or traded upstream in local urban markets	
NEMATODES				
Root-knot nematodes: Meloidogyne spp.				
The presence of Meloidogyne promotes and aggravates infection by soil-borne fungi such as Fusarium spp. Infected plants are very sensitive to drought and irregular irrigation.				
Infested roots develop galls and rot if conditions are humid.	If plants are heavily attacked during the early stages, they may die.	Significant decrease if plants are heavily attacked during early stages.		
Source : Enquêtes de terrains, Mars 2024.				

The analysis of Tables 3 to 6 shows, on the one hand, that among the plant organs of watermelon plants, the most vulnerable and frequently attacked are the leaves, followed by the fruits, the floral organs, and sometimes the entire plant. The consequences caused by watermelon pests and/or diseases include yield reduction, post-harvest losses, a decrease in the market value of harvested watermelons, a deterioration in the nutritional and medicinal quality of harvested products, disruptions to the growth cycles of production, and the inability to sell on local or export markets. The areas planted and yields obtained over the

past five (05) years by the responding watermelon producers are shown in Table 7. The fight against watermelon pests and diseases in southern Benin is based primarily on the use of chemicals. These pesticides are mostly purchased on the market. Commonly used chemicals are presented in Table 9. Regarding the fertilizers used by watermelon producers in southern Benin, most use chemical fertilizers. It should be noted that poultry droppings are specifically used by some producers in Fonsa, in the commune of Ouidah, in addition to other fertilizers.



Photo 1. Some watermelon pests and diseases.

Source: Field surveys, March 2024

Table 7. Areas and Yields Obtained Over the Past Five Years.

Year	Area (m ²)	Yield (fruits)
2018	83270	35123
2019	107170	45322
2020	127220	53477
2021	159220	68066
2022	155500	65117

Source : Field surveys, March 2024

The analysis of Table 7 shows an increase in sown areas from 83,270 to 159,220 m2 between 2018 and 2021, reflecting a continued increase in yields, rising from 35,123 to 68,066 m2 over the same period before falling slightly in 2022. Table 8 presents the average, minimum, maximum and standard deviation of sown areas and yields obtained over the last five years.

Table 8. Average sown area and average yield obtained over the last five years.

Designation	Average	Minimum	Maximum	Standard deviation
Sup18	1205,846	0,0000	4362,50	1328,004
Sup19	1522,702	0,0000	5975,00	1672,662

Sup20	1711,263	50,0000	6125,00	1412,301
Sup21	2226,187	666,6667	10000,00	2027,325
Sup22	2121,869	700,0000	5000,00	1240,645
Rend18	503,784	0,0000	1999,00	569,245
Rend19	634,518	0,0000	2683,75	709,176
Rend20	714,191	21,0000	2762,50	621,534
Rend21	935,652	273,3333	4000,00	828,110
Rend22	1008,405	209,0000	4600,00	961,642

Source : Field surveys, March 2024

Table 8 shows an increasing trend from 2018 to 2022 in all the parameters indicated, namely means, minimums, maximums, and standard deviations. The average trend in watermelon production in southern Benin over the past five years is shown in Figure 3.




Figure 3. Average evolution of production.

Source : Field surveys, March 2024

Table 9. Commonly used chemicals against watermelon pests and diseases in southern Benin.

Designation	Chemical fertilizer	Organic fertilizer	Chemical control	Biological control	Integrated control
Proportion(%)	99.986	0.014	100	00	00
Elements used	NPK, UREE, K ₂ SO ₄ , SUPER GROS	Les fientes	Pacha 25Ec, Acarius 018Ec, Emacot 019Ec, Kaobtimal, Lambda	-	-

Source : Field surveys, March 2024

Chemical fertilizers (NPK; UREA: K₂SO₄; SUPER GROS) and chemical control (Pacha 25Ec; Acarius 018Ec; Emacot 019Ec; Kaobtimal; Lambda) are widely used in proportions of 99.99% and 100%, respectively, while organic fertilizers made from droppings are used very sparingly, at 0.014%.

The survey allowed us to gather watermelon producers' impressions of the impacts of chemicals used to control watermelon pests and diseases on the environment and the fruit produced. The responses obtained are presented in Table 10 and the graph below.

Table 10. Effects of the use of plant protection products on the environment.

Consequence	Rate (%)
Depletion of production land	1,389
Groundwater pollution	2,778
None	95,833

Source : Field surveys, March 2024.

The consequences of the use of phytosanitary products on the environment are extremely widely declared without major negative impacts at 95.833% (Table XV). The graph in Figure 4 is the schematic representation of Table 10.




Figure 4. Effect of Chemicals Used on the Environment.

Source. Field Surveys, March 2024

Graph 5 above describes the effect of chemicals according to watermelon producers. It shows that 1.389% of watermelon producers acknowledged that the pesticides used contribute to the depletion of production land, 2.778% stated that these chemicals pollute the water table and according to the remaining 95.833%, these chemicals used have no impact on the environment and watermelon fruits. The control of watermelon pests and diseases in southern Benin is based primarily on the use of chemicals. However, limiting pest populations can be achieved through natural biological control using beneficial insects: this is the case for thrips, whiteflies, leaf miners, aphids, etc. No mass release of beneficial insects is carried out for technical and economic reasons. Only introductory releases of new species can be considered. Sometimes, for other pests, biological control is still insufficient (few or no beneficial insects reach them), and it is necessary to treat them specifically and preventively according to a schedule: this is the case for cucurbit moth caterpillars. It is necessary to support this biological control with specific chemical treatments when beneficial insects are insufficient in number, for example, against leaf miners or aphids. In this case, curative treatments are used with products that preserve beneficial insects. Finally, it is important to note that new pests and diseases appear regularly. Therefore, the use of an effective control method must be considered. In a comparative approach, these results show that chemical control remains the primary method used by watermelon producers in southern Benin to deal with pests and diseases. This observation is largely confirmed by James *et al.* (2010), who emphasize that in tropical market gardening systems, the intensive use of insecticides remains a direct

response to the economic losses caused by pests, despite the deleterious effects on biodiversity and health. However, more sustainable alternatives exist. In this regard, Van Lenteren (2000) points out that biological control, particularly through the conservation or increase of natural beneficial insects, constitutes an ecologically viable solution. Several beneficial insects are known to effectively control populations of thrips, whiteflies, leafminers, and aphids (Hassan, 1993; Bale *et al.*, 2008). In Benin, the lack of mass releases of beneficials is often due to technical and economic constraints, which limits the effectiveness of this method on a large scale (Kéita *et al.*, 2017).

Furthermore, some pests, such as cucurbit moth caterpillars (*Diaphania* spp.), remain relatively unresponsive to locally available beneficials, necessitating the use of specific chemical treatments. In this vein, the work of Kogan (1998) and Ehler (2006) argues that, in an integrated pest management approach, chemical treatments can be compatible with the protection of natural enemies, provided they are selective and applied in a targeted manner. It is also crucial to note the regular emergence of new pests linked to factors of mutation and climate change, and the globalization of trade, without ignoring the evolution of agricultural practices. In their research, Altieri and Nicholls (2003) emphasize that the resilience of agroecosystems depends in part on their ability to integrate different control methods (biological, cultural, mechanical, and rational chemical). The watermelon pest control strategy in southern Benin must evolve toward an integrated approach, combining technical effectiveness, reasonable costs, and environmental protection.

Integrated pest management offers an alternative to traditional chemical control by keeping pest populations low, preventing them from causing economic damage. (Ryckewaert, 1998). It also applies to disease control. Thus, controlling fungal diseases in watermelon is based on the application of preventive and cultural measures such as: Use of healthy seeds and the use of disease-free plant material; Proper management of factors influencing greenhouse climatic conditions, namely irrigation; Avoiding the presence of free water on the crop; Rationalizing fertilization and limiting excessive nitrogen inputs; The plot must be well-drained to avoid water stagnation and excess humidity; 6- Implementing crop rotations. Removing and destroying plant debris and anything left over from pruning the crop. Furthermore, it is essential to follow certain rules regarding chemical treatments: Wetting, i.e., the amount of water used per unit area, must be sufficient: from approximately 400 liters per hectare for a young plantation to 1,200 liters per hectare for a high-yield crop (e.g., tomatoes); for a sprayer, the wetting rate must be halved. Comply with the prescribed doses (i.e., the amount of product per 10 liters of water, for example). Remember to double the dose when using a sprayer because the wetting rate is halved (the amount of product per area, however, always remains the same). Too high a dose risks causing phytotoxicity. • Products should be alternated whenever possible to avoid habituation or resistance among insects and diseases. Products from different chemical families should be chosen. Respect the pre-harvest application intervals for each crop, otherwise there is a risk of residues in the parts consumed (Ryckewaert, 1998). It should be noted that when using integrated pest management, other benefits are also obtained, such as improved air, water, soil, and plant and animal quality in cultivated and non-cultivated environments. The comparative study of watermelon pest management approaches in southern Benin highlights the growing importance of integrated pest management as a sustainable alternative to conventional chemical control. The latter, although still largely dominant among producers, has limitations from both environmental and health perspectives. According to Ryckewaert (1998), integrated pest management aims to maintain pest populations below the threshold of economic harm through a combination of preventive, cultural, biological and, as a last resort, chemical practices.

One of the main advantages of integrated pest management lies in its ability to reduce dependence on pesticides, the excessive use of which is often observed in horticultural production systems in West Africa (James *et al.*, 2010). This systematic use, without strict adherence to doses, safety intervals, or active ingredient rotation principles, creates risks of phytotoxicity, pest resistance, and harmful residues on fruits, as Ryckewaert also points out. In terms of fungal disease management, the integrated approach is essentially based on prevention, through the adoption of good agricultural practices: use of healthy seeds, humidity control (through rational irrigation and drainage), crop rotation, and removal of plant debris, among others. These measures are widely recommended in

the work of Elad and Pertot (2014), who demonstrate that prophylaxis is often more effective and less costly than curative treatments, particularly in tropical environments where conditions favor the development of pathogenic fungi. Furthermore, the rational application of plant protection products as part of integrated pest management involves strict adherence to application procedures, particularly with regard to wetting, dosage, and alternating chemical families, in order to limit the resistance of pests and pathogens. This principle is supported by Kogan's (1998) recommendations on the importance of preserving beneficials and reducing environmental impacts. Another fundamental aspect of integrated pest management is its positive impact on the environment. By reducing chemical inputs, it promotes better soil, water, and air quality, while preserving biodiversity, particularly the natural enemies of pests. This agroecological approach is in line with the principles defended by Altieri and Nicholls (2003) who emphasize the increased resilience of diversified and ecologically managed crop systems.

CONCLUSION

This research is part of the characterization of watermelon (*Citrullus lanatus*) insect pests and diseases and the various control methods used in southern Benin. The study covered the municipalities of Missirété, Covè, Za-kpota, Comè, Dangbo, Dogbo, Lokossa, Tori-Bossito, Allada, Adjohoun, Kétou, Granp-Popo, and Ouidah. It made it possible to identify watermelon pests and diseases in southern Benin. These include: flies, beetles, root-knot nematodes, downy mildew, leaf spot, anthracnose, fusarium wilt, and mosaic viruses. Indeed, the fight against watermelon (*Citrullus lanatus*) pests and diseases in southern Benin is based primarily on the use of chemical products such as Pacha 25EC, Acarius 018EC, Emacot019EC, Kaobtimal, Lambda, etc. However, some particularly harmful pests or diseases, such as root-knot nematodes (*Meloidogyne* sp.), fusarium wilt (*Oxysporum* sp., *Oxysporum niveum*), and bacterial fruit spot (*Acidovorax avenaesub*, *Acidovorax* sp., *Acidovorax citrulli*), can be avoided through long crop rotation, good drainage, and the use of tolerant or resistant cultivars. Similarly, polyethylene mulch, especially when coated with a layer of reflective aluminum paint, repels thrips and aphids. We suggest that officials from the Ministry of Agriculture, Livestock, and Fisheries (MAEP), particularly those from the Territorial Agency for Agricultural Development (ATDA), educate watermelon producers through Field Schools (FS) on integrated pest management, which is an alternative to conventional chemical control and also helps keep pest populations low, preventing them from causing economic damage.

ACKNOWLEDGMENT

The authors would like to thank all those who contributed to the completion of this work.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

ETHICS APPROVAL

Not applicable

AI TOOL DECLARATION

The authors declares that no AI and related tools are used to write the scientific content of this manuscript.

DATA AVAILABILITY

Data will be available on request

REFERENCES

- Abdoul Habou Z., Haubruge E., Adam T., François J. Verheggen (2013). Focus on: Insect pests and biocidal properties of *Jatropha curcas* L. (Euphorbiaceae): bibliographic summary. *Biotechnologie, Agronomie, Société et Environnement*, 17(4), 604 – 612.
- Adandonon, A., Tamò, M., & Aveling, T. (2009). Integrated pest and disease management in West Africa: an alternative to chemical pesticides. IITA, Cotonou.
- Agbangla, C., Savi, M. K., & Houndete, T. A. (2017). Contribution of watermelon to food security and producers' incomes in southern Benin. *African Journal of Agriculture and Sustainable Development*, 12(3), 45-53.
- Aglinglo A.L., Azon F.C., Lègba C.E., Djido U., Agossou O.C., Francisco R., Fassinou Hotegni V.N., Achigan-Dako G.E. (2018). Synthetic technical sheet for the production of watermelon (*Citrullus lanatus* (Thunb.) Matsum. Et Nakai.).
- Aglinglo E. G., Lys Amavi, Azon, F. Christel, Legba, C. Eric, Djido, Ulrich, Agossou, Aboègnonhou Chaldia Odette, Francisco, A. Rachidi, Hotegni, Vodjo Nicodeme Fassinou, Achigan-Dako, "Synthetic technical data sheet for the production of watermelon (*Citrullus lanatus* (Thunb.) Matsum. Et Nakai.)," Lab.
- Ahouangninou C.C.A. (2013). Sustainability of market gardening production in southern Benin: an attempt at the ecosystem approach. Doctoral thesis from the University of Abomey, 333pp.
- Altieri, M.A., & Nicholls, C.I. (2003). Agroecology: Rescuing a science, revitalizing a practice. *Journal of Sustainable Agriculture*, 22 (3), 93-120.
- Atachi, P., Tossou, G., & Zannou, S. (2016). Agroecological management of market garden pests in Benin. Proceedings of scientific conferences in Benin.
- Bale, J.S., Van Lenteren J.C., Bigler F. (2008). Biological control and sustainable food production. *Philosophical Transactions of the Royal Society B*, 363 (1492), 761-776.
- Bassirou S. et Guero M.A. (2024). Regional Chamber of Agriculture of Maradi / Niger. Watermelon production in the Maradi Region. 7pp.
- Braun U. (1982). Morphological studies in the genus *Oidium* III. *Zentralblatt für Mikrobiologie*. 137 (4), 314-324.
- CABI (2005). Collection on crop protection, edition 2005
- CACED, (2018). The Market for Vegetable Products in Southern Benin: Dynamics and Perspectives. Action Center for the Environment and Sustainable Development (ACED), Benin. 33 pp.
- Cendres A. (2010). Innovative process for extracting fruit juice using microwaves: manufacturing viability and nutritional quality of juices. Food and Nutrition. *University of Avignon*, 2010. 289pp.
- Chomicki S., Renner S. (2015). Watermelon origin solved with molecular phylogenetics including Linnaean material: another example of museomics. *New Phytology*, 205-526 – 32.
- Dansou C.S. (2019). Structuring and spatial dynamics of the cities of Ouidah and Grand-Popo on the Beninese coast: Environmental issues and perspectives. Geography. University of Abomey-Calavi (Benin), 2019, 376pp.
- DSA (2016). Final results of the agricultural campaign: 2015-2016. Directorate of Agricultural Statistics. 100pp.
- Dupin T. (2017). Observation of pests and their natural enemies in citrus orchards managed with agroecological practices in Martinique. Master 2 thesis, Sciences Technology Health, Mention Biology and Plant Technology, Specialization Production and Plant Technology. University of Angers and University of Rennes, 90pp.
- Ehler, L.E. (2006). Integrated pest management (IPM): Definition, historical development and implementation, and the other IPM. *Pest Management Science*, 62(9), 787–789.
- Elad, Y., & Pertot, I. (2014). Climate change impacts on plant pathogens and plant diseases. *Journal of Crop Protection*, 72, 86 -98.
- Emmanuelle Osbert Kichah (2016). Root architecture of herbaceous species: diversity of establishment and plasticity. Agricultural sciences. Doctoral thesis at the University of Avignon, 2016. French. 174pp.
- Esquinas-Alcazar J. T, Gulick PJ (1983), Genetic resources of Cucurbitaceae – a global report. International Board for Plant Genetic Resources.
- FAO (Organisation des Nations Unies pour l'alimentation et l'agriculture). (2017). Integrated pest management : principles and practices. Rome: FAO.
- FAO. (2020). Good practices for pest control of vegetable crops in West Africa. Rome: Food and Agriculture Organization of the United Nations.
- Goudegnon, A. B., Sikirou, R., & Kpéra, G. N. (2012). Biological control of agricultural pests in Benin: current

- status and prospects. *Benin Agricultural Review*, 10(3), 21-29.
- Hamza N. (2014). Application of arbuscular mycorrhizae in market gardening: case of watermelon (*Citrullus lanatus*). Master's thesis in Plant Biology and Physiology, Ferhat Abbas University, Sétif 1.
- Hassan, S.A. (1993). The current status of biological control of pests in greenhouses. *IOBC/WPRS Bulletin*, 16(2), 1–8.
- Houhou Mohamed Labib (2022). Study of the phytosanitary situation of cucurbits in the wilaya of Biskra, Zeb el Gharbi region. Master's thesis at the University of Biskra, *Natural and Life Sciences*, 74pp.
- Houkpèvi, A., Zossou, E., & Gandonou, C. (2020). Vulnerability factors of market gardening farms to pests in southern Benin. *Journal of Agronomic Sciences*, 25(1), 30-42.
- James B., Atcha-Ahowé, C. Godonou, I., Baimey H., Goergen G., Sikirou, R., Toko M. (2010). Integrated Pest Management in Vegetable Production: A Guide for Extension Workers in West Africa. IITA-Cotonou.
- Jean-Philippe Deguine (2023). Biological control and biocontrol: a need for clarification, *Cah. Agriculture*. 2023, 32, 11
- Jeroen Boland, Irene Koomen, Joep van Lidth de Jeude, Jan Oudejans. (2004). Pesticides: Composition, Use, and Risks. 124 pp.
- Kéita B., Zannou A., Glèlè Kakai R. (2017). Constraints of implementing biological control in market gardening systems in southern Benin. *African Journal of Agronomy and Sustainable Development*, 12 (1), 45 - 58.
- Kogan, M. (1998). Integrated Pest Management: Historical perspectives and contemporary developments. *Annual Review of Entomology*, 43, 243 - 270.
- Koudjil Mohamed (2021). Plant Protection: Biological Control. Master II Course, Hassiba Benbouali University of Chlef, 59pp.
- Kyriacou M., Leskovar D., Colla G., Roupheal, Y. (2018). Watermelon and melon fruit quality: Genotypic and agro-environmental factors involved. *Scientia Horticulturae*, 234, 393-408.
- Lecoq H., Desbiez C. (2012). Chapitre 3 – Cucurbit crop viruses in the Mediterranean region: a constantly evolving picture. Editeur(s): Gad Loebenstein, Hervé Lecoq, *Advances in Virus Research*, Academic Press, 84, 67-126.
- Morita M., Hayashi T., Ochiai M., Tsukaguchi K., Tsuboi Y., Matsumoto Y., Hara Y. (2014). Oral supplementation with watermelon extract increases plasma L-arginine concentrations and reduces blood pressure in prehypertensive individuals: A randomized, double-blind, placebo-controlled trial. *American Journal of Hypertension*, 27 (1), 75-80.
- Moutaïrou, K., Aho, N., & Afouda, F. (2018). Identification and dynamics of watermelon pests in southern Benin. *Beninese Agricultural Review*, 9(2), 15-23.
- Myriam Siegwart (2021). Combinations, characteristics and origins of resistance mechanisms to (bio)insecticides in crop pest insects. Agricultural Sciences. *University of Avignon*, 2021. 193pp.
- N’Ggo S., Diome T., Diop B. et Sembene M. (2020). Effect of aqueous extracts of *Calotropis procera* on the main pests of cabbage grown in Senegal. *International Journal of Biological and Chemical Sciences*. 14 (5), 1600 -1610.
- Niane K., Seid Ali M., Ayessou N.C., Cissé M. et Faye G. (2021). Nectar production and stabilization of two varieties of watermelon (*Citrullus lanatus*). *Africa Science*, 19 (5), 1-11.
- Oyédé M. Inès (2023). Department of Climatology and Agrometeorology. Study and Training Department. State of the climate in Benin for the year 2023. 23pp.
- PARIS H. S. (2015) Origin and emergence of the sweet dessert watermelon, *Citrullus lanatus*. *Journal Annals of Botany*, 116 -133 - 48, doi : 10.1093/aob/mcv077.
- PNACCB (2022). Benin's National Climate Change Adaptation Plan. Ministry of Living Environment and Sustainable Development, General Directorate of *Environment and Climate* (DGEC). 175pp.
- Pretty, J., & Bharucha, Z. P. (2015). Integrated pest management for sustainable intensification of agriculture in Asia and *African Insects*, 6(1), 152-182.
- Ryckewaert, P. (1998). Integrated crop protection: concepts, methods and issues. CIRAD, Montpellier.
- Ryckewaert, P. (1998). Integrated protection of market garden crops under cover in humid tropical zones: methods and means. CIRAD.
- Sherf A.F. and A.A. Macnab (1986). Plant Diseases and Their Control. 2nd Edition. ISBN: 0-471-05860-2
- Sikirou, R., & al. (2013). Virulent pests and diseases associated with watermelon: observations in Beninese market gardening areas. Technical report INRAB.
- Tamò, M., Arodokoun, D. Y., & Adandonon, A. (2014). Ecological approaches for sustainable pest management in West Africa. IITA-Benin, Scientific Report.
- USDA (2023). U.S. Department of Agriculture), national nutrient database for standard reference legacy release, basic report: 09326, Watermelon, raw. <https://ndb.nal.usda.gov/ndb/foods/show/09326> consulté le 04 Février 2023.
- Van Lenteren, J. C. (2000). A greenhouse without pesticides: Fact or fantasy?. *Crop Protection*, 19 (6), 375-384.
- Xavier M. (2018). Natural regulation of insect pests in vegetable crops and its consequences on production: quantification of the service provided and search for levers for its intensification. *Animal Biology, Agrocampus* Ouest,

2018. Doctoral thesis in agronomy at the University of Brittany Loire, Agro Campus, 279pp.
- Yanogo Pawendkisgou I.Y., Sanogo S., Yaméogo J. (2021). Watermelon (*Citrullus lanatus*), an opportunity for marketing stakeholders in the city of Koudougou (Burkina Faso). *Ivorian Review of Savannah Geography*, (10), pp.177-195.
- Yarou B.B., Pierre S., Komlan F.A., Mensah A., Alabi T., Verheggen F., Francis F. (2017). Pesticidal plants and protection of market garden crops in West Africa (bibliographic synthesis), *Biotechnologie, Agronomie, Société et Environnement*, 21 (4), 288 – 304.
- Yarou, B. B., Dossa, G., & Zannou, S. (2015). Pesticide Use in Agriculture in Benin: Perceptions and Risks. *Journal of Environmental Sciences*, 13(2), 85-97.