



Review Article

## RED FLOUR BEETLE, *TRIBOLIUM CASTANEUM* (HERBST): BIOLOGY AND MANAGEMENT

Mukesh Kumar Chaubey

Department of Zoology, National Post Graduate College, Barahalganj-273402, Gorakhpur, U.P

**Article History:** Received 22<sup>nd</sup> February 2023; Accepted 3<sup>rd</sup> March 2023; Published 23<sup>rd</sup> March 2023

### ABSTRACT

According to an estimate, the world's population will reach upto 9.1 billion till 2050 and food production is projected to increase up to 70%. Developing countries already facing food scarcity contribute most in population growth. In these countries, one in every six children suffers from hunger and malnutrition. In these critical situation, protection of the grains and food products from insect infestation during pre-harvest and post-harvest is urgently necessary it causes mass losses of grains. In several countries, about 15% of grains are lost harvesting or post-harvesting stages. The Food and Agricultural Organization (FAO) has estimated this loss up to 40% in India. Among the culprits of these losses, coleopteran insect species play major roles. Among coleopterans, red flour beetle, *Tribolium castaneum* causes loss of grains and food products to larger extent in post-harvest stage. *T. castaneum* is a secondary pest with a wide host range infesting among, barley, beans, ground nuts, lentils, maize, oat, peas, rice, rye, sorghum, wheat and other processed grain products. Adults and larvae attack grains already damaged by primary pest and feed mainly on the germ. For the management of *T. castaneum* population, several approaches including synthetic insecticides have been applied with lots of success but leaving several environmental and human health issues too. To minimize these adverse effects, plant derived volatile chemicals i.e. essential oils and its constituents alone or in combination have been used and got promising results. Since these are volatile, non-persistent and biodegradable, essential oil based formulations can be used for the management of *T. castaneum* at large as green eco-friendly insecticide.

**Keywords:** Biopesticides, Essential oil, Insecticide resistance, *Tribolium castaneum*.

### INTRODUCTION

About 20,000 species of pests including 40 rodents, 150 fungi, 355 mites, 70 moths and more than 600 species of beetles have been reported to damage of agricultural products both qualitative and quantitative at pre-harvest as well as post-harvest stages (Rajendran, 2005). This is a serious problem especially in developing countries where stored-product insect pests cause severe pre- and post-harvest losses estimated to be more than 20% in developing countries and up to 9% in developed countries (Pimentel, 2009). Insects deteriorate grain quality in storage making more susceptible to fungal attack, unsuitable for consumption, planting and trade. Among the stored-product insect pests, members of order Coleoptera and Lepidoptera are common (Boyer *et al.*, 2012). Of these, coleopterans are widely distributed and the most serious pests (Stejskal *et al.*, 2015). Both larvae and adults feed on the grain/grain

products causing serious economic losses. The red flour beetle, *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae) is one of the most damaging insect species that attack stored grain and food products throughout the world (Andrić *et al.*, 2010; Edde *et al.*, 2012). It is a secondary pest of all grains and primary pests of flour and other processed grain products. It infests usually those grains that had already been damaged by other pests. The affected grains and products become contaminated prompting fungal growth. Economic losses include unpleasant smell, reduced weight, reduced trade and nutritional value. Presence of this insect pest may also cause allergic responses.

### TAXONOMIC POSITION AND DISTRIBUTION

The red flour beetle, *T. castaneum* is a secondary pest of stored grain and products throughout the world. It is

\*Corresponding Author: Mukesh Kumar Chaubey, Department of Zoology, National Post Graduate College, Barahalganj -273402, Gorakhpur, U.P, Email: mgpgc9839427296@gmail.com

regarded as one of the most damaging polyphagous insect pest of stored grains and products (Goodman *et al.*, 2012; Neenah, 2014; Devi and Devi, 2015). The adults are long-lived and sometimes living more than three years. Several names of red flour beetle viz. *Tribolium navale* (Fabricius, 1775), *Colydium castaneum* (Herbst, 1787), *Tenebrio castaneus* (Schönherr, 1806), *Phaleria castanca* (Gyllenhal, 1810), *Uloma ferruginea* (Dejean, 1821), *Margus castaneus* (Dejean, 1833), *Stene ferruginea* (Westwood, 1839) and *Tribolium ferrugineum* (Wollaston, 1854) are being used as synonyms. *T. castaneum* is considered to have Indian or Indo-Australian origin with a high dispersion rate (Hill, 2002; Ridley *et al.*, 2011). It is cosmopolitan, but, more common in warmer regions than to temperate regions of the world (Hill, 2002).

### MORPHOLOGY

Adults are small, flat, 2.3-4 mm long and red brown coloured (Figure 1)(Mahroof and Hagstrum, 2012). The antennae of *T. castaneum* are composed of eleven segments, with last three abruptly enlarged segments, in contrast with *T. confusum* in which antennal segment enlarge gradually (Figure 2) (Baldwin and Fasulo, 2014). Thorax is slightly darker than the elytra. Elytra are punctuated containing parallel-sided striations. The males have a setaceous lesion on the ventral surface of front femur (Haines, 1991). Adult bears a functional hind wings often used in flight (Ridley *et al.*, 2011). Larvae are initially whitish, but, later become brown and the size reaches up to 5 mm. Its terminal abdominal segment bears a pair of dark and upturned projections.

### HOST

*T. castaneum*, one of the most cosmopolitan pests of stored products, is polyphagous damaging grains and products both quantity and quality (Bachrouch *et al.*, 2010; Neenah, 2014). It attacks a widespread variety of stored grains and its products including beans, biscuits, cornflakes, dried fruits, lentils, maize, rice, sorghum, wheat and wheat flour (Ranga Rao *et al.*, 2010, Devi and Devi, 2015). The damaged grains and products contain frass, carcasses and exuviae. It also turns grey in colour with a pungent unacceptable odour due to benzoquinones, a defensive chemical secreted from their prothoracic and abdominal glands. All these make the grains and products unsuitable for human consumption decreasing trade value as well (Devi and Devi, 2015). Both larvae and adults feed externally on grains, flours and other processed food products. Its larvae consume mostly the germ of the wheat grains (White and Lambkin, 1988).

### LIFE CYCLE AND BIOLOGY

*T. castaneum* is a holometabolous insect showing eggs, larval, pupal and adult phases in the life cycle. The females mate several times and lay about 300-600 eggs. Eggs are white, microscopic and covered with sticky material to which flour particles adhere. About 5-12 days of

incubation, eggs hatch into larvae. The hatched larvae feed on any available food for about 3-4 weeks and undergo pupation. Pupal phase lasts for 5-9 days. It takes 7 to 12 weeks to complete its life cycle, however, life span depends on temperature and relative humidity. A temperature of 27°C has been found to be the most suitable for development. There are four to five generations per year.

### Eggs

Eggs are whitish, microscopic (0.61mm long and 0.3mm width) and cylindrical with remnants of flour particles adhering to the surface with the help of sticky substance present over the eggs shell (Li and Arbogast, 1991; Devi and Devi, 2015). This makes their detection in flour very different. The egg period lasts in about 5-9 days (Devi and Devi, 2015).

### Larvae

The larvae are minute, 4-5 mm in length, cylindrical, slender, brownish-white, active feeders and well sclerotised when fully grown. Larval phase includes six to seven instars (Devi and Devi, 2015). Duration of larval phase varies depending upon on availability, type and quality of food, temperature and relative humidity (Haines, 1991). It requires 12-13 days with an average of 12.9 days to complete (Haines, 1991; Devi and Devi, 2015).

### Pupae

Pupa is brown coloured, quiescent, non-feeding and without cocoon. Pupal length varies between the sexes. The average body length of male pupa is about 3.18±0.3mm with a body width of 1.07±0.03mm. The average body length of female pupa is 4.12±0.01mm with a body width of 1.15±0.01mm (Devi and Devi, 2015). Pupal duration also varies according to sex, with male pupal period of 6-7days, while female period lasts in 7-9 days (Devi and Devi, 2015).

### Mating behavior

Females of *T. castaneum* show polyandry ( a single female can copulate with several males in the same copulation period). Polyandry increases progeny production as during copulation with several males, female receives large amount of sperms and choose cryptically which sperm is used for fertilization (Fedina and Lewis, 2004). It is also important as many sexually active males are non-virgins and may be sperm-depleted (Pai *et al.*, 2005). Polyandry also increases genetic variability in the population increases genetic variability in the population. Females of different geographic regions often show variations in mating behaviour suggesting that polyandry can be advantageous in some populations but not in others (Pai *et al.*, 2005). Males also select females to mate. Males with large number of odour receptors prefer and select mature virgin females by recognizing non-virgin females through scent produced by the reproductive glands. Virgin females lack such scent glands (Arnaud, 1999).

## ECONOMIC IMPORTANCE

*T. castaneum* damage stored grains and food products causing huge economic loss especially in warmer areas of the world. It infests grains already damaged by other pests or during harvest and storage. The affected product becomes contaminated with faeces, has unpleasant smell and loses nutritional and trade value. Quinones secreted from thoracic and abdominal glands deteriorate grain and product quality to worst. Temperature and moisture content affect feeding behaviour. Increase in temperature by 4°C from 28 to 32°C increases weight loss of whole grain two times (Majeed *et al.*, 2016). Increase in moisture content of whole grain from 12.2 to 14.2% increases damage to 1.6 times during the first 60 days, 1.3 times during next 60 days and 1.2 times during the last 60 days of infestation (Daniels, 1956). During development, a single *T. castaneum* larva consumes 13 mg of wheat flour and adult during their lifetime consumes 315 mg of wheat flour (Hagstrum and Subramanyam, 2000). After consuming the kernel of wheat, a larva feeds on the endosperm (Daniels, 1956). During life, adult consumes the entire germ of 7-12 wheat (Karunakaran *et al.*, 2004). *T. castaneum* can spread infection of *Aspergillus flavus*, *Aspergillus fumigatus*, *Cladosporium herbarum*, *Enterococcus faecalis*, *Penicillium citinum* and *P. purpurogenum* (Channaiah *et al.*, 2010; Yun *et al.*, 2018).

## MANAGEMENT

### Detection methods

*T. castaneum* can be difficult to detect at low densities. Simple traps baited with food such as carob beans, ground nuts and cereals can help in detecting individuals when their density is low. At higher densities, adults and larvae can be seen in holes and tunnels in grains and dust produced during feeding. The eggs stuck to storage containers can also be detected. Presence of a pungent odour indicates high density infestation.

### Physical control

Proper sanitation of storage area, removal, destruction of all the sources of infestation and sieving of grains can reduce chances of flour beetle infestation. Addition of inert dusts to the grain causes death of insect by desiccation. Suspected grain can be stored in containers with tight fitting lids or in a freezer for 4-5 days or at temperatures of 50°C. This reduces oxygen levels and increases carbon dioxide level causing death of insect. The application of diatomaceous earth powder to adult *T. castaneum* infesting wheat causes cent percent mortality (Aldryhim, 1990).

### Molecular control

The RNAi technique is a new eco-friendly method of insect pest management. In this technique, synthetic RNA are used to target and inhibit specific genes within the insect species. This approach has been applied and found

effective against *T. castaneum* (Tang *et al.*, 2016; Gao *et al.*, 2017). However, this method has not yet been marketed due high cost of production of the RNA.

### Chemical Control

At present, insecticides like dichlorvos, pyrethrins, pyrethroids and insect growth regulators like methoprene and pyriproxyfen are being used frequently (Sutton *et al.*, 2011). Pyrethrin has found wide applications because of its low toxicity against mammals and high efficacy against a broad range of stored-product insect pests. Commercial formulations used often contain synergist like piperonyl butoxide to increase efficacy of pyrethrins (Brooke, 1958). Insect growth regulators cause malformation or reduces adult emergence. Applications of pyrethrins/pyrethroids alone or with insect growth regulators give better results insect management (Mullen, 1999). However, the disadvantage associated with these synthetic chemicals is the residual effects and persistence nature.

### Biological Control

Different strains of entomopathogenic fungus have shown some hope in the management of stored grain insect pest. Two entomopathogenic fungi, *Metarhizium anisopliae* and *Beauveria bassiana* have been reported to show positive results against many stored grain insect pests including *T. castaneum* (Golshan *et al.*, 2013; Zamani *et al.*, 2013; Sewify *et al.*, 2014; Abdu-Allah *et al.*, 2015; Rumbos and Athanassiou, 2017; Batta, 2018). Various studies have also conducted to explore the possibility of entomopathogenic fungi either alone or in combination with other strategies and received hopeful outcomes (Wakil *et al.*, 2012; Batta and Kavallieratos, 2018). *T. castaneum* beetles produce and secrete a mixture of quinone containing methyl-1,4-parabenzoquinone, ethyl-1,4-parabenzoquinones and 1-pentadecene (Villaverde *et al.*, 2007). These quinones act as an external immune defence to protect the insect against pathogens.

## INSECTICIDE RESISTANCE

A wide variety of insecticides has been applied as the main approach for the control and management of *T. castaneum*. These insecticides target the insect's nervous system including voltage-gated ion channels and acetylcholine esterase causing irreversible disruption of neurological function resulting in death of insects. But the continuous application of these insecticides has developed resistance in most of the the target insects. In Indian, the first case of insecticide resistance was reported in *T. castaneum* against malathion (Bhatia and Pradha, 1971). Since then, high cases of insecticide resistance have been reported in *T. castaneum* (Dhaliwal and Chawla, 1995; Padhee *et al.*, 2002; Arshad *et al.*, 2019). The development of insecticide resistance is a slow and gradual evolutionary process. After the initial exposure to the insecticide, there is a latent period during which resistance genes are segregated and recombine with other genes that make the conditions favourable for resistance development.

Now, the target insect species show an increased level of tolerance to the insecticide. Resistance develops gradually followed by a period of rapid development of resistance resulting in explosive growth of the insect's population. There are a variety of mechanisms that cause insecticide resistance in *T. castaneum*. These are:

- i. Target site insensitivity: In this, changes in sensitivity of target site inhibit insecticide binding.
- ii. Metabolic resistance: In this, production of increased quantity of enzymes leads to increased metabolic detoxification of insecticide.
- iii. Lack of penetration: In this, cuticle thickens or undergoes remodelling which prevents penetration of insecticides.
- iv. Behavioural resistance: In this, insects adapt some new behaviour to avoid exposure to insecticides.

#### Target site insensitive resistance

DDT and pyrethroids target voltage-gated sodium ion channels in insects (Narahasi *et al.*, 1992). Cyclodienes and fipronil bind to the gamma aminobutyric acid receptor (Casida, 1993). Organophosphates, carbamates and pyrethroids produce neurotoxicity by inhibiting activity of acetylcholine esterase (Russell *et al.*, 2004; Pang, 2014). These insecticides also affect other target sites such as voltage-gated sodium ion channels and gamma aminobutyric acid receptors. Resistant insect shows normal neurological functions irrespective of the presence of insecticide as they have evolved insensitive acetylcholine receptors which provide resistance to organophosphate and carbamate insecticides. There are mainly four types of target site insensitivity mechanisms viz. altered Acetylcholine esterase resistance mechanism, knockdown resistance mechanism, reduced gamma amino butyric acid receptor sensitivity mechanism and altered acetylcholine receptors.

Altered acetylcholine esterase resistance mechanism: Acetylcholine esterase is required to breakdown acetylcholine into acetate and choline to terminate synaptic transmission (Pitman, 1991). This enzyme is a target site of organophosphates and carbamates insecticides, which are bound to a serine residue on the active site of acetylcholine esterase making the enzyme inactive. The inhibition of acetylcholine esterase results in accumulation of acetylcholine at the synaptic cleft leading to prolonged binding of acetylcholine to its postsynaptic receptor. This causes neuroexcitation and produces symptoms like tremors, convulsions and death (Siegfried, 2001). Several organophosphates have been used to protect stored grains from insect infestation. But most of them have developed resistance due to insensitive acetylcholine esterase. Insensitivity of acetylcholine esterase occurs due to point mutations in the active site of Ace genes encoding acetylcholine esterase (Kono and Tomita, 2006).

Knockdown resistance mechanism: Pyrethrin, pyrethroids and DDT target voltage gated sodium ion

channels in insects. These bind to the voltage-gated ion channels in axonal membranes resulting in opening of ion channels for a long time and prolonged sodium influx, thereby, depolarizing the axonal membrane and stimulating the neurons to produce repetitive discharges causing paralysis and death in insects (Soderlund and Bloomquist, 1990). In insects, sodium ion channel is encoded by a single gene known as para (Dong, 2010). Changes in the structure of voltage-gated sodium ion channel occur by point mutation or substitution mutation. It reduces the binding affinity of the insecticides to ion channel protein causing insensitivity.

#### Metabolic resistance

This resistance cause over production of enzymes like cytochrome *P450* monooxygenases, carboxylesterases and glutathione transferases (Gilbert *et al.*, 2005; Feyerisen, 2006). These enzymes detoxify insecticides before exerting their toxic effect. *T. castaneum* has evolved metabolic resistance for organophosphates, carbamates and pyrethroids (Liang *et al.*, 2015; Huang *et al.*, 2019; Wei *et al.*, 2019).

#### Cuticle resistance mechanism

Generally, contact insecticide penetrates insect's cuticle and reaches the target site to exert its toxic action (Liu *et al.*, 2006). Cuticle resistance is the result of thickening or remodelling of cuticle which reduces the penetration of insecticide into the insect's body (Koganemaru *et al.*, 2013; Lilly *et al.*, 2016). The over expression of laccases and ABC transporters has been reported to be involved in this mechanism.

#### Behavioural resistance

In this, behaviour of insect is so modified that it reduces exposure period to insecticides. Guedes *et al.* (2009) reported higher rates of flight take-off in some insects which increases the insect's survival chances by reducing exposure time.

#### ESSENTIAL OILS

The continuous and indiscriminate applications of synthetic insecticides at large cause bioaccumulation and biomagnification in different environmental components. This adversely and severely affects non-target organisms including human and ecosystems. Therefore, plant based insecticides have been attracted as alternative of these synthetic insecticides. In this continuation, various essential oils and its constituents have been explored as repellent, insecticidal, antifeedant and insect growth regulator activities against a variety of stored grain insects. The essential oils are complex mixture of different chemical compounds with multiple modes of action that enhances their activity due to the synergistic action among constituents (Regnault-Roger, 2012). Since essential oils are volatile, biodegradable and non-persistent in nature, these can be a better candidate for insect pest management.

About 17500 aromatic plant species and more than 3000 essential oil constituents have been identified (Maddocks-Jennings *et al.*, 2005). These are produced as secondary metabolites in the members of Apiaceae, Asteraceae, Combretaceae, Geraniaceae, Gramineae, Lamiaceae, Myrtaceae, Meliaceae, Piperaceae, Rutaceae, Verbenaceae and Zingiberaceae (Bhattarai and Karki, 2004). These are extracted from plant parts like leaves, flowers, roots, buds, rhizomes etc through steam or water distillation. Generally,

essential oils have densities lower than those of water (Perricone *et al.*, 2015). These are lipophilic and, thus, can enter into insect and cause biochemical dysfunction and mortality (Lee *et al.*, 2004). The toxicity of essential oils does not depend only on the chemical compounds but also on many other factors like route of entry, molecular weights and the mechanisms of action. Essential oils with insecticidal activities can be inhaled, ingested or skin absorbed by insects.

Class: Insecta  
Order: Coleoptera  
Superfamily: Tenebrionoidea  
Family: Tenebrionidae  
Subfamily: Tenebrioninae  
Genus: *Tribolium*  
Species: *castaneum*



**Figure 1.** *Tribolium castaneum*

*T. castaneum*:  
Last three segments enlarge abruptly

*T. confusum*:  
Segments enlarge gradually

**Figure 2.** Difference between antenna of *T. castaneum* and *T. confusum*

### Repellent activity

The repellent activity of several essential oils has been reported against *T. castaneum*. Repellent behaviour of *Baccharis salicifolia* essential oil have been observed against *T. castaneum* adults (Garcia *et al.*, 2005). *Tagetes terniflora* and *Artemisia sieberi* oil have been reported effective as repellent against adults and fifth instar of *T. castaneum* (Stefanazzi *et al.*, 2006; Negahban and Moharamipour, 2007). *Laurelia sempervirens* and *Drimys winteri* essential oils repelled *T. castaneum* adults (Zapata and Smagghe, 2010). *Schinus molle* and *Salvia mirzayanii* essential oils have been reported for strong repellent activity against *T. confuse* adults (Abdel-Sattar *et al.*, 2010; Nikooei and Moharramipour, 2011). Repellent activity of *Cuminum cyminum*, *Piper nigrum*, *Illicium verum*, *Myristica fragrans*, *Foeniculum vulgare*, *Trachyspermum ammi*, *Anethum graveolens*, *Nigella sativa*, *Zingiber officinale*, *Piper cubeba*, *Allium sativum* essential oils has been determined against *T. castaneum* adults (Chaubey, 2007a,b; Shukla *et al.*, 2008; Chaubey, 2011; 2013; 2016). Besides complete essential oils, its pure constituents have also been tested for repellent activities (Kim *et al.*, 2010; Chaubey, 2011).

### Insecticidal activity

Since essential oils easily vaporise at room without any biochemical decomposition, these can be used as fumigants. The insecticidal activity of several essential oils has been reported against adults and larvae of *T. castaneum*. Essential oils of *Anethum sowa*, *Artemisia annua*, *Lippia alba* and *Elletaria cardomum* have been reported for fumigant toxicity (Tripathi *et al.*, 2000;

Aggrawal *et al.*, 2000; Verma *et al.*, 2000; Huang *et al.*, 2000). Fumigant toxicity of *C. cyminum*, *P. nigrum*, *I. verum*, *M. fragrans*, *F. vulgare*, *T. ammi*, *A. graveolens*, *N. sativa*, *Z. officinale*, *P. cubeba*, *A. sativum* essential oils has been determined against *T. castaneum* adults (Chaubey, 2007a, b; Shukla *et al.*, 2008; Chaubey, 2011; 2013; 2016). Fumigation of essential oils of *Carum copticum* and *C. carvi* causes mortality in *T. castaneum* (Sahaf *et al.*, 2008; Fang *et al.*, 2010). Besides complete essential oils, pure compounds have also been established for fumigant toxicity. Trans-anethole has been reported to show toxicity against *T. castaneum* (Koul *et al.*, 2007). The toxic effects of *Citrus reticulata* and *C. sinensis* oils compounds have been studied against larvae and adult of *T. castaneum* (Saleem *et al.*, 2013). Fumigation of  $\beta$ -caryophyllene and  $\alpha$ -pinene kills the larvae and adult of *T. castaneum* (Chaubey, 2012).

### Antifeedant activity

Antifeedants are compounds or substances that reduce insect feeding. These substances deter feeding by modification the insect behaviour affecting peripheral sensilla of insects (Isman, 2000). Tripathi *et al.* (2002) have reported feeding deterrence activity of *C. longa* leaf essential oil against adult and larvae of *T. castaneum*. Antifeedant activity of *Eucalyptus globulus* and *Lavandula stoechas* essential oils have been observed against *T. castaneum* (Ebdadollahi, 2011). Huang and Ho (1998) established antifeedant activity of cinnamaldehyde against *T. castaneum*. Antifeedant activity of 1,8-cineole has been demonstrated in *T. castaneum* (Tripathi *et al.*, 2001). Tripathi *et al.* (2002) have reported antifeedant activity of

*Curcuma* leaf oil and d-limonene in *T. castaneum*. Chaubey (2012) has evaluated antifeedant activities of *Zingiber officinale* and *Piper cubeba* essential oils as well as pure compounds,  $\alpha$ -pinene and  $\beta$ -caryophyllene against *T. castaneum*.

### Insect growth regulators activity

Several essential oils and its constituents have been established for their oviposition, growth inhibitory and progeny production inhibitory activities against different insect pests including *T. castaneum*. These disrupt mating behaviour, growth and metabolism of insects. *Z. officinale* and *P. cubeba* oils have been evaluated for developmental inhibitory activities against *T. castaneum*. These two oils inhibit development of larvae to pupae and the pupae to adults (Chaubey, 2011). Essential oil of *A. sativum* reduces oviposition potential of adults (Chaubey, 2013). *C. cyminum*, *P. nigrum*, *F. vulgare*, *T. ammi*, *A. graveolens*, *I. verum*, *M. fragrans* and *N. sativa* essential oils reduce oviposition potential and increase developmental period of *T. castaneum*. Fumigation inhibits development of larvae to pupae and the pupae to adults and also result in the deformities in different developmental stages of insect (Chaubey, 2007a, b; Shukla *et al.*, 2008). Ajayi and Olonisakin (2011) have been evaluated ovicidal activity of *Syzygium aromaticum*, *Piper guineense* and *Xylopi aethiopica* essential oils against *T. castaneum*. The three essential oils are able to reduce progeny emergence of *T. castaneum*. Besides complete oils, constituents have also shown insect growth regulatory activities. Developmental inhibitory activities of  $\alpha$ -pinene and  $\beta$ -caryophyllene alone or in binary combination have been determined against 4th instars larvae of *T. castaneum*. These compounds reduce transformation of larvae into pupae and pupae into adult (Chaubey, 2012). Carvone suppresses egg hatching in *T. castaneum* (Tripathi *et al.*, 2003).

### Mechanism of action

Essential oils and its constituents disrupt insect physiology in different ways. Some of these inhibit acetylcholine esterase enzyme activity or block octopamine receptors (Enan, 2001; Chaubey, 2013). Ryan and Byrne (1975) have reported that essential oils and monoterpenoids cause insect mortality by reversibly competitive inhibition of acetylcholine esterase by binding to active site of the enzyme (Ryan and Byrne (1975). Some essential oils block gamma aminobutyric acid and induce toxicity in insects (Priestley *et al.*, 2003). Chaubey (2013) has reported that *A. sativum* essential oil inhibits acetylcholine esterase activity in *T. castaneum* adults (Chaubey, 2013). Some monoterpenes such as thujone can induce the neurotoxic effects by acting gamma aminobutyric acid receptors in insects (Hold 2000; Ratra and Casida, 2001). Essential oils can also disrupt communication in mating behaviour of insect by blocking the function of antennal sensilla. This lowers fecundity and ultimately the population of insect pest (Ahmad *et al.*, 2001). Caryophyllene oxide inhibits

electron transport system by disturbing respiration rate of insects (Emeki *et al.*, 2004). Thus, in conclusion essential oils and its constituents exert its toxicity in insects by interfering with the nervous co-ordination and respiratory system.

### Structure activity relationship

The bioactivities of essential oils and their constituents depend on the functional group and molecular weights (Kumbhar and Dewang, 2001). L-Menthol and its acyl derivatives have been evaluated against stored grain insect pests. It has been reported that menthyl propionate and L-menthol have high insecticidal activity. High activity of menthyl propionate as compared to L-menthol can be due to increasing number of methyl groups in side chain (Aggarwal *et al.*, 2001). Due to nucleophilic properties of methyl group, increase in number of methyl groups on side chain cause increase in negative charge on carbon atom. Therefore, high activity of menthyl propionate may be due to increasing negative charge on carbon atom because of the presence of two methyl groups. Low activity of menthyl benzoate has been observed because they do not have methyl group and the benzene ring is attached directly to menthyl carbon. Activity of menthyl cinnamate has been found moderate as it has a double bond and benzene ring is not directly attached to carbon atom (Aggarwal *et al.*, 2001; Tripathi *et al.*, 2009). Benzene derivatives (eugenol, isoeugenol, methyl eugenol, safrole and isosafrole) and terpenes (cineole, limonene, p-cymene and  $\alpha$ -pinene) have been evaluated for insecticidal activity (Ngoh *et al.*, 1998). It has been observed that derivatives of benzene have higher insecticidal activity than that of terpenes. This toxicity may be due to the active groups in benzene derivatives. The presence of double bond in the side chain of aromatic ring and the substitution of methoxy group play an important role in the toxicity of these analogues. The knockdown and contact activity has been found to increase in methyl eugenol due to further methoxy group. In contrast, when double bond in side chain is nearer to aromatic ring, fumigant toxicity is decreased. However, benzene derivatives have more insecticidal activity than monoterpenes (Ngoh *et al.*, 1998). Further researches involving structure activity are required to enhance insecticidal potency of essential oils and its constituents. Further studies should be carried out to investigate whether the alteration in structure of essential oil constituents can modify mode of action.

### Synergistic action of essential oils

Regnault-Roger (1997) reported that the high toxic effect of essential oil may be due to the synergistic effect of essential oil active compounds. Various studies have been carried out to investigate the synergistic effect of essential oils, active components and their combinations against insects. Khalfi *et al.* (2006) studied the insecticidal activity of spearmint essential oils and two active compounds 1,8 cineole and carvone. They found that the activity is due to the synergistic action of 1,8-cineole and carvone.

Abbassy *et al.* (2009) reported that terpinen-4-ol and  $\alpha$ -terpinene increase the insecticidal activity of profenofos and methomyl. The essential oils are a complex of chemical compounds with multiple modes of action that enhances their activity due to the synergistic action.

## CONCLUSION

Essential oils are complex mixtures of volatile compounds produced as secondary metabolite in aromatic plants. These oils and its constituents have repellent, insecticidal, antifeedant and growth regulatory activities in insects. These interfere with the nervous and respiratory system of the insect. Due to their volatile and non-persistent nature, essential oils can be used as an alternative against insect pests of stored grain and food products. Most of these oils are selective in their role with little or no harmful effect on the environment and the non-target organisms including human. Thus, essential oils can be used as alternatives in the management of red flour beetle, *T. castaneum*.

## ACKNOWLEDGMENT

The authors express sincere thanks to the head of the Department of Zoology, National Post Graduate College, Barahalganj, Gorakhpur, U.P for the facilities provided to carry out this research work.

## REFERENCES

- Abbassy, M. A., Abdelgaleil, S. A. M., & Rabie, R. Y. A. (2009). Insecticidal and synergistic effects of *Majorana hortensis* essential oil and some of its major constituents. *Entomology Experiment Applicata*, 131, 225-232.
- Abdel-Sattar, A. A., Zaitoun, E., Farag, M. A., El Gayed, S. H. & Harraz, F. M. H. (2010). Chemical composition, insecticidal and insect repellent activity of *Schinus molle* L. leaf and fruit essential oils against *Trogoderma granarium* and *Tribolium castaneum*. *Natural Products Research*, 24, 226-235.
- Abdu-Allah, G., Abou-Ghadir, N.M., Nasser, M., & Metwaly, M.R. (2015). Comparative efficiency of the fungi, *Beauveria bassiana*, *Metarhizium anisopliae* and the natural product spinosad, using three economic coleopterous stored grain insects. *Egypt Journal of Biology and Pest Control*, 25(3), 715.
- Agarwal, M., Walia, S., & Dhingra, S. (2000). Insect growth inhibition, antifeedant and antifungal activity of compounds isolated/derived from *Zingiber officinale* rhizomes. *Pest Management Science*, 37, 289-300.
- Aggarwal, K. K., Tripathi, A. K., Ahmad, A., Prajapati, V., Verma, N., & Kumar, S. (2001). Toxicity of L-menthol and its derivatives against four storage insects. *Insect Science and Application*, 21, 229-235.
- Ahmed, K. S., Yosui, Y., & Lachikawa, T. (2001). Effect of neem oil on mating and oviposition behavior of azuki bean weevil, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). *Pakistan Journal of Biological Sciences*, 4(11), 1371-1373.
- Ajayi, F. A., & Olonisakin, A. (2011). Bioactivity of three essential oils extracted from edible seeds on the rust-red flour beetle, *Tribolium castaneum* (Herbst) infesting stored pearl millet. *Trakia Journal of Science*, 9(1), 28-36.
- Andrić, G., Kljajić, P., Perić, I., & Golić, M.P. (2010). Susceptibility of red flour beetle *Tribolium castaneum* (Herbst) populations from Serbia to contact insecticides. *Julius-Kühn Archives*, 425, 869-873.
- Arshad, A., Munawar, A., Mastoi, M. I., Sohail, S., Liang, C. (2019). Evaluation of Phosphine and Cypermethrin resistance in field collected strains of *Rhyzopertha dominica* (Coleoptera: Bostrichidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae) under laboratory conditions. *Evaluation*, 4(5).
- Bachrouh, O., Ben Jemba, J. M., Wissem, A. W., Talou, T., Marzouk, B., & Abderrab, M. (2010). Composition and insecticidal activity of essential oil from *Pistacia lentiscus* L. against *Ectomyelois ceratoniae* Zeller and *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae). *Journal of Stored Products Research*, 46, 242-247.
- Baldwin, R., & Fasulo, T. R. (2014). Featured creatures. University of Florida. ENY-289.
- Batta, Y.A. (2018). Efficacy of two species of entomopathogenic fungi against the stored grain pest, *Sitophilus granarius* L. (Curculionidae: Coleoptera), via oral ingestion. *Egypt Journal of Biology Pest Control*, 28(1), 44.
- Batta, Y.A., & Kavallieratos, N.G. (2018). The use of entomopathogenic fungi for the control of stored-grain insects. *International Journal of Pest Management*, 64(1), 77-87.
- Bhatia, S. K., & Pradhan, S. (1971). Studies on resistance to insecticides in *Tribolium castaneum* (Herbst)-III. Selection of a strain resistant to lindane and its biological characteristics. *Journal Stored Products Research*, 6(4), 331-337.
- Bhattarai, N., & Karki, M. (2004). Medicinal and Aromatic Plants-Ethnobotany and Conservation Status. In: Encyclopedia of Forest Sciences, Burley, J., Evans, J., & Youngquist, J. (Eds.). Academic Press, London, UK, 523-532.
- Boake, & Christine, R. B. (1985). Genetic Consequences of Mate Choice: A Quantitative Genetic Method for Testing Sexual Selection Theory. *Science*, 227(4690), 1061-1063.
- Boyer, S., Zhang, H., & Lempérière, G. (2012). A review of control methods and resistance mechanisms in

- stored-product insects. *Bulletin Entomology Research*, 102(2), 213-229.
- Brooke, J.P. (1958). The stabilization of pyrethrins by piperonyl butoxide in water, with special reference to their use as mosquito larvicides. *Annals of Applied Biology*, 46(1), 254-259.
- Casida, J.E. (1993). Insecticide action at the GABA-gated chloride channel: Recognition, progress, and prospects. *Archives Insect Biochemistry Physiology*, 22(1-2), 13-23.
- Channaiah, L. H., Subramanyam, B., & Zurek, L. (2010). Survival of *Enterococcus faecalis* OG1RF: pCF10 in poultry and cattle feed: vector competence of the red flour beetle, *Tribolium castaneum* (Herbst). *Journal of Food Protect*, 73, 568-73.
- Chaubey, M. K. (2007a). Insecticidal activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella sativa* (Ranunculaceae) essential oils against stored-product beetle *Tribolium castaneum* herbst (Coleoptera: Tenebrionidae). *African Journal of Agricultural Research*, 2(11), 596-600.
- Chaubey, M. K. (2007b). Toxicity of essential oils from *Cuminum cyminum* (Umbelliferae), *Piper nigrum* (Piperaceae) and *Foeniculum vulgare* (Umbelliferae) against stored product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Electronic Journal of Environmental Agriculture Food Chemistry*, 6(1), 1719-1727.
- Chaubey, M.K. (2011). Toxicity of Essential Oil Combinations against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *World Journal of Zoology*, 6(1), 22-25.
- Chaubey, M. K. (2012). Responses of *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae) against essential oils and pure compounds. *Herbal Pollution*. 58(3), 33-35.
- Chaubey, M. K. (2013). Insecticidal effect of *Allium sativum* (Alliaceae) essential oil against *Tribolium castaneum* (Coleoptera: Tenebrionidae). *TBAP*, 3(4), 248-258.
- Chaubey, M. K. (2016). Fumigant and contact toxicity of *Allium sativum* (Alliaceae) essential oil against *Sitophilus oryzae* L. (Coleoptera: Dryophthoridae). *Entomology Applied Science Letter*, 3(2), 43-48.
- Daniels, N. E. (1956). Damage and reproduction by the flour beetles *Tribolium confusum* and *T. castaneum* in wheat at three moisture contents. *Journal of Economics Entomology*, 49, 244-7.
- Devi, M. B., & Devi, N. V. (2015). Biology of rust-red flour beetle, *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Biological Forum- An International Journal*, 7, 12-15.
- Dhaliwal, B. K., & Chawla, R. F. (1995). Evaluation of current status of Malathion resistance in *Tribolium castaneum* (Herbst)(Coleoptera: Tenebrionidae) in Punjab, India. *Pesticide Research Journal*, 7(1), 54-7.
- Dong, K. (2010). Progress in insect sodium channel research. In: Gilbert LI, Gill SS, editors. *Insect pharmacology: Channels, receptors, toxins and enzymes*. Academic Press. 25-27.
- Ebadollahi, A. (2011). Antifeedant activity of essential oils from *Eucalyptus globules* Labill and *Lavandula stoechas* L. on *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Biharean Biologist*. 5(1). 8-10.
- Edde, P.A., Eaton, M., Kells, S.A., & Phillips, T.W. (2012). Biology, behavior, and ecology of pests in other durable commodities. In: Hagstrum, D.W., Phillips, T.W. and Cuperus, G. (eds.) *Stored product protection*. Manhattan, KS: Kansas State University, 45-62.
- Emekci, M., Navarro, S., Donahaye, E., Rindner, M., & Azrieli, A. (2004). Respiration of *Rhyzopertha dominica* (F.) at reduced oxygen concentration. *Journal of Stored Products Research*, 40, 27-38
- Enan, E., (2001). Insecticidal activity of essential oils: Octopaminergic sites of action. *Comparative Biochemistry Physiology Part C: Toxicology Pharmacology*, 130, 325-337.
- Fang, R., Jiang, C. H., Wang, X. Y., Zhang, H. M., Liu, Z. L., Zhou, L., Du, S. S., & Deng, Z. W. (2010). Insecticidal activity of essential oil of *Carum carvi* fruits from China and its main components against two grain storage insects. *Molecules*, 15(12), 9391-402.
- Fedina, T. Y. & Lewis, S. M. (2004). Female influence over offspring paternity in red flour beetle *Tribolium castaneum*. *Proceedings of the Royal Society B: Biological Sciences*, 271 (1546), 1393-1399.
- Feyereisen, R. (2006). Evolution of insect P450. *Biochemical Society Transactions*, 34(6), 1252-1255.
- Gao, S., Liu, X., Liu, J., Xiong, W., Song, X., Wu, W., Wei, L., & Li, B. (2017). Identification and evolution of latrophilin receptor gene involved in *Tribolium castaneum* development and female fecundity. *Genesis*, 55(12), e23081.
- Garcia, M., Donadel, O. J., Ardanaz, C. E., Tonn, C. E., Sosa, M. E. (2005). Toxic and repellent effects of *Baccharis salicifolia* essential oil on *Tribolium castaneum*. *Pest Management Science*, 61, 612-618.
- Gilbert, L. I., Iatrou, K., & Gill, S. S. (2005). *Comprehensive molecular insect science*. Oxford: Elsevier, 1-77.
- Golshan, H., Saber, M., Majidi-Shilsar, F., & Karimi, F. (2013). Efficacy of nine isolates of *Beauveria bassiana*



- against adults of *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Acta Entomologica Sinica*, 56(2), 201-206.
- Goodman, C. L., Stanley, D., Ringbauer Jr, J. A., Beeman, R. W., Silver, K., & Park, Y. (2012). A cell line derived from the red flour beetle *Tribolium castaneum* (Coleoptera: Tenebrionidae). *In Vitro Cellular & Developmental Biology-Animal*, 48(7), 426-433.
- Guedes, N. M. P., Guedes, R. N. C., Ferreira, G. H., & Silva, L. B. (2009). Flight take-off and walking behaviour of insecticide-susceptible and-resistant strains of *Sitophilus zeamais* exposed to deltamethrin. *Bulletin Entomology Research*, 99, 393-400.
- Hagstrum, D.W., & Subramanyam, B. H. 2000. Monitoring and decision tool. In: Subramanyam, B.H., & Hagstrum, D. W. (Ed.), *Alternatives to pesticides in stored-product IPM*. 1-28. Kluwer Academic Publishers.
- Haines, C. P. (1991). *Insects and arachnids of tropical stored products: their biology and identification (a training manual)*. 2.
- Hold, K.M., Sirisoma, N.S., Ikeda, T., Narahashi, T., & Casida, J. E. (2000).  $\alpha$ -Thujone (the active component of absinthe): $\gamma$ -Aminobutyric acid type A receptor modulation and metabolic detoxification. *Proceeding Natural Academic Science USA*.
- to phosphine in China and functions of cytochrome P450s in phosphine resistance. *Journal of Pest Science*. 92(3), 1239-1248.
- Isman, M.B., (2000). Plant essential oils for pest and disease management. *Crop Protection*. 19, 603-608.
- Karunakaran, C., Jayas, D. S., & White, N. D. G. (2004). Identification of wheat kernels damaged by the red flour beetle using x-ray images. *Biosystem Engineering*. 87,267-74.
- Khalfi, O., Benyoussef, E.H., & Yahiaoui, N. (2006). Extraction, analysis and insecticidal activity of spearmint essential oil from Algeria against *Rhyzopertha dominica* (F.). *Journal of Essential Oil-Bearing Plants*, 9, 17-21.
- Kim, S. I., Yoon, J. S., Jung, J. W., Hong, K. B., Ahn, Y. J., & Kwon, H. W. (2010). Toxicity and repellency of origanum essential oil and its components against *Tribolium castaneum* (Coleoptera: Tenebrionidae) adults. *Journal of Asia-Pacific Entomology*, 13, 369-373.
- Koganemaru, R., Miller, D. M., & Adelman, Z. N. (2013). Robust cuticular penetration resistance in the common bed bug (*Cimex lectularius* L.) correlates with increased steady-state transcript levels of CPR-type cuticle protein genes. *Pesticide Biochemistry and Physiology*, 106(3), 190-197.
- Kono, Y., & Tomita, T. (2006). Amino acid substitutions conferring insecticide insensitivity in Ace-paralogous acetylcholinesterase. *Pesticide Biochemistry and Physiology*, 85(3), 123-132.
- Koul, O., Singh, G., Singh, R., & Singh, J. (2007). Mortality and reproductive performance of *Tribolium castaneum* exposed to Anethol vapours at high temperature. *Biopesticides International*, 3, 126-137.
- Kumbhar, P.P. & Dewang, P.M. (2001). Monoterpenoids: The natural pest management agents. *Fragrance Flavor Assoc. India*, 3, 49-56.
- Lee, B.H., Annis, P.C., Tumaalii, F. & Choic, W.S. (2004). Fumigant toxicity of essential oils from the Myrtaceae family and 1,8-cineole against 3 major stored-grain insects. *Journal of Stored Products Research*, 40, 553-564.
- Li, L., & Arbogast, R. T. (1991). The effect of grain breakage on fecundity, development, survival and population increase in maize of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Stored Products Research*, 27, 87-94.
- Liang, X., Xiao, D., He, Y., Yao, J., Zhu, G., & Zhu, K. Y. (2015). CYP-mediated up-regulation of cytochrome P450 genes in the red flour beetle (*Tribolium castaneum*). *International Journal of Molecular Sciences*, 16(1), 2078-2098.
- Lilly, D. G., Latham, S. L., Webb, C. E., & Doggett, S. L. (2016). Cuticle thickening in a pyrethroid-resistant strain of the common bed bug, *Cimex lectularius* L. (Hemiptera: Cimicidae). *Plos One*. 11(4):e0153302.
- Liu, N., Zhu, F., Xu, Q., Pridgeon, J. W., & Gao, X. (2006). Behavioral change, physiological modification, and metabolic detoxification: mechanisms of insecticide resistance. *Acta Entomologica Sinica*. 49, 671-679.
- Maddocks-Jennings, W., Wilkson, J. M., & Shillington, D. (2005). Novel approaches to radiotherapy-induced skin reactions: A literature review. *Complementary Therapies in Clinical Practice*, 11, 224-231.
- Mahroof, R. M., & Hagstrum, D. W. (2012). Biology, behavior, and ecology of insect in processed commodity. In: *Stored Product Protection*. Hagstrum, D. W., Phillips, T. W. & Cuperus, G. (Eds). Kansas State University.
- Majeed, M. Z., Javed, M., Khaliq, A., & Afzal, M. (2016). Estimation of losses in some advanced sorghum genotypes incurred by red flour beetle, *Tribolium castaneum* L. (Herbst.) (Tenebrionidae: Coleoptera). *Pakistan Journal of Zoology*, 48, 1133-1139.
- Mullen, M. A. (1999). Special use pheromone-baited trap for the red flour beetle (Coleoptera: Tenebrionidae). *Journal of Entomology Science*, 34, 497-500.

- Narahashi, T., Frey, J. M., Ginsburg, K. S., & Roy ML. (1992). Sodium and GABA-activated channels as the targets of pyrethroids and cyclodienes. *Toxicology Letters*, 64, 429-36.
- Negahban, M., & Moharamipour, S. (2007). Repellent activity and persistence of essential oil from *Artemisia sieberi* besser on three stored-product insect species. *Iranian Journal of Medicinal and Aromatic Plants*, 22, 293-302.
- Nenaah, G. E. (2014). Chemical composition, toxicity and growth inhibitory activities of essential oils of three *Achillea* species and their nano-emulsions against *Tribolium castaneum* (Herbst). *Industrial Crops and Products*, 53, 252-260
- Ngoh, S. P., Choo, L. E. W., Pang, F. Y., Huang, Y., Kini, M. R., & Ho, S. H. (1998). Insecticidal and repellent properties of nine volatile constituents of essential oils against the American cockroach, *Periplaneta americana* (L.). *Pest Science*, 54, 261-268.
- Nikooei, M., & Moharramipour, S. (2011). Fumigant toxicity and repellency effects of essential oil of *Salvia mirzayani* on *Callosobruchus maculatus* and *Tribolium confusum* (Coleoptera: Tenebrionidae). *Journal of Entomological Society of Ira*, 30, 17-30.
- Padhee, A. K., Saxena, J. D., Sinha, S. R., & Srivastava, C. (2002). Selection for Resistance to Deltamethrin in Red-Flour Beetle, *Tribolium castaneum* (Herbst). *Annals Plant Protection Science*, 10(2), 220-4.
- Pai, A., Bennett, L., & Yan, G. (2005). Female multiple mating for fertility assurance in red flour beetles (*Tribolium castaneum*). *Canadian Journal of Zoology*, 83(7), 913-919.
- Pang, Y. P. (2014). Insect acetylcholinesterase as a target for effective and environmentally safe insecticides. *Advance Insect Physiology*, 46, 435-94.
- Perricone, M., Arace, E., Corbo, M. R., Sinigaglia, M., & Bevilacqua, A. (2015). Bioactivity of essential oils: A review on their interaction with food components. *Frontiers Microbiology*, 6, 76
- Pimentel, D. (2009). Pesticides and pest control. In: Peshin, R. & Dhawan, A.K. (Eds.) *Integ.*
- Pitman, R. M. (1971). Transmitter substances in insects: a review. *Comp. General Pharmacology*, 2(7), 347-371.
- Priestley, C. M., Williamson, E. M., Wafford, K.A., Sattelle, D.B. (2003). Thymol, a constituent of thyme essential oil, is a positive allosteric modulator of human GABA<sub>A</sub> receptors and a homo-oligomeric GABA receptor from *Drosophila melanogaster*. *British Journal of Pharmacology*, 140, 1363-1372.
- Rajendran, S. (2005). Detection of insect infestation in stored foods. *Advances in Food and Nutrition Research*, 49, 163-232.
- Ranga Rao, G. V., Rameshwar Rao, V. & Nigam, S. N. (2010). Post-harvest insect pests of groundnut and their management. Information Bulletin No. 84. Patancheru 502324, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. ISBN 978-92-9066-528-1. Order code IBE 084. 20.
- Ratra, G. S. & Casida, J.E. (2001). GABA receptor subunit composition relative to insecticide potency and selectivity. *Toxicology Letters*, 122, 215-222.
- Regnault-Roger, C. (1997). The potential of botanical essential oils for insect pest control. *Integrated Pest Management. Reviews*, 2, 25-34.
- Regnault-Roger, C., Vincent, C. & Arnason, J.T. (2012). Essential oils in insect control: Low-risk products in a high-stakes world. *Annual Review of Entomology*, 57, 405-424.
- Ridley, A., Hereward, J. P., Daghli, A., Raghu, G. J., Collins, P. J. & Walter, G.H. (2011). The spatiotemporal dynamics of *Tribolium castaneum* (Herbst): adult flight and gene flow. *Molecular Ecology*, 20, 1635-1646.
- Rumbos, C. I., & Athanassiou, C. G. (2017). Use of entomopathogenic fungi for the control of stored-product insects: can fungi protect durable commodities? *Journal of Pest Control Science*, 90(3), 839-854.
- Russell, R. J., Claudianos, C., Campbell, P. M., Horne, I., Sutherland, T. D., & Oakeshott, J. G. (2004). Two major classes of target site insensitivity mutations confer resistance to organophosphate and carbamate insecticides. *Pesticide Biochemistry Physiology*, 79(3), 84-93.
- Ryan, M. E., Byrne, O. (1988). Plant insect coevolution and inhibition of acetylcholinesterase. *Journal of Chemical Ecology*, 14, 1965-1975.
- Sahaf, B. Z., Moharramipour, S., Meshkatsadat, M. H. & Filekesh, E. (2008). Repellent activity and persistence of the essential oils from *Carum copticum* and *Vitex pseudo-negundo* on *Tribolium castaneum*. *Integrated Protect Stored Prod. IOBC Bull.* 40, 205-210.
- Saleem, M., Hussain, D., Rashid, R. H., Saleem, H. M., Ghouse, G., & Abbas, M. (2013). Insecticidal activities of two citrus oils against *Tribolium castaneum* (Herbst). *American Journal of Research Community*, 1(6), 67-74.
- Sewify, G., El Shabrawy, H., Eweis, M. & Naroz, M.H. (2014). Efficacy of entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae* for controlling certain stored product insects. *Egypt Journal of Biology Pest Control*. 24(1), 191.
- Shukla, J., Tripathi, S. P., & Chaubey, M. K. (2008). Toxicity of Myristica fragrance and Illicium verum essential oils against flour beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Elective Journal of Environmental. Agricultural Food Chemistry*, 7(7), 3059-3064.
- Siegfried, B. D., & Scharf, M. E. (2001). Mechanisms of organophosphate resistance in insects. In: Biochemical sites of insecticide action and resistance. 269-291. Springer, Berlin, Heidelberg.
- Soderlund, D. M., & Bloomquist, J. R. (1990). Molecular mechanisms of insecticide resistance. In: Roush, R. T., Tabashnik, B.E. Editors. Pesticide resistance in arthropods. Boston, MA: Springer. 58-96.
- Stefanazzi N, Gutierrez MM, Stadler T, Bonini NA, Ferrero AA. Biological activity of essential oil of *Tagetes*

- terniflora* Kunth (Astereaceae) against *Tribolium castaneum* Herbst (Insecta, Coleoptera, Tenebrionidae). *Biology Sanidad Veg Plagas*. 2006; 32(3), 439-447.
- Stejskal, V., Hubert, J., Aulicky, R. & Kucerova, Z. (2015). Overview of present and past and pest-associated risks in stored food and feed products: European perspective. *Journal of Stored Products Research*, 64, 122-132.
- Sutton, A. E., Arthur, F. H., Zhu, K. Y., Campbell, J. F., & Murray, L. W. (2011). Residual efficacy of synergized pyrethrin and methoprene aerosol against larvae of *Tribolium castaneum* and *Tribolium confusum* (Coleoptera: Tenebrionidae). *Journal of Stored Products Research*, 47, 399-406.
- Tang, B., Wei, P., & Zhao, L. (2016). Knockdown of five trehalase genes using RNA interference regulates the gene expression of the chitin biosynthesis pathway in *Tribolium castaneum*. *BMC Biotechnology*, 16, 67.
- Tripathi, A. K., Prajapati, V., Aggarwal, K., & Kumar, S. (2001). Toxicity, feeding deterrence, and effect of activity of 1,8-cineole from *Artemisia annua* on progeny production of *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Journal of Economic Entomology*, 94, 979-983.
- Tripathi, A. K., Prajapati, V., Aggrawal, K.K., Khanuja, S. P. S., & Kumar, S. (2000). Toxicity towards *Tribolium castaneum* in the fraction of essential oil of *Anethum sowa* seeds. *Journal of Medicine Aromatic Plant Science*, 22, 40.
- Tripathi, A. K., Prajapati, V., Khanuja, S. P. S., & Kumar, S. (2003). Effect of d-limonene on three stored-product beetles. *Journal of Economic Entomology*, 96(3), 990-995.
- Tripathi, A. K., Prajapati, V., Verma, N., Bahl, J. R., Bansal, R. P., Khanuja, S. P. S., & Kumar, S. (2002). Bioactivities of the leaf essential oil of *Curcuma longa* (var. *ch-66*) on three species of stored-product beetles (Coleoptera). *Journal of Economic Entomology*, 95(1), 183-189.
- Tripathi, A. K., Upadhyay, S., Bhuiyan, M., Bhattacharya, P. R. (2009). A review on prospects of essential oils as biopesticides in insect pest management. *Journal of Pharmacognostic and Phototherapy*, 1, 52-63.
- Verma, N., Tripathi, A. K., Prajapati, V., Bahl, J. R., Khanuja, S. P. S., Kumar S. (2000). Toxicity of essential oil from *Lippia alba* towards stored grain insects. *Journal of Medicine Aromatic Plant Science*, 22, 50.
- Villaverde, M. L., Juárez, M. P. & Mijailovsky, S. (2007). Detection of *Tribolium castaneum* (Herbst) volatile defensive secretions by solid phase microextraction-capillary gas chromatography (SPME-CGC). *Journal of Stored Products Research*, 43(4), 540-545.
- Wakil, W., Kavallieratos, N. G., Usman, M., Gulzar, S., El-Shafie, H. A. (2021). Detection of phosphine resistance in field populations of four key stored-grain insect pests in Pakistan. *Insects*, 12(4), 288.
- Wei, L., Gao, S., Xiong, W., Liu, J., Mao, J., Lu, Y., Song, X., & Li, B. (2019). Latrophilin mediates insecticides susceptibility and fecundity through two carboxylesterases, esterase4 and esterase6, in *Tribolium castaneum*. *Bulletin. Entomology Research*, 109(4), 534-543.
- Yousif, N. A. (1990). Efficacy of the amorphous silica dust, Dryacide, against *Tribolium confusum* Duv. and *Sitophilus granarius* (L.) (Coleoptera: Tenebrionidae and Curculionidae), *Journal of Stored Products Research*, 26(4), 207-210.
- Yun, T. S., Park, S. Y., Yu, J. Y., Hwang, Y. J., & Hong, K. J. (2018). Isolation and identification of fungal species from the insect pest *Tribolium castaneum* in rice processing complexes in Korea. *Plant Pathology Journal*, 34, 356-66.
- Zamani, Z., Aminaee, M. M., & Khaniki, G.B. (2013). Introduction of *Beauveria bassiana* as a biological control agent for *Tribolium castaneum* in Kerman province. *Archives Phytopathology Plant Protection*, 46, 2235-2243.
- Zapata, N., & Smagghe, G. (2010). Repellency and toxicity of essential oils from the leaves and bark of *Laurelia sempervirens* and *Drimys winteri* against *Tribolium castaneum*. *Indian Journal of Crops Production*, 32, 405-410.