



Research Article

POPULATION DYNAMICS OF MAJOR INSECT PESTS INFESTING CHILLY, BOTTLE GOURD AND CUCUMBER

¹Arti Yadav, ¹Parveen Kumar and ^{2*}Kirpa Ram

¹Department of Zoology, Baba Mastnath University, Asthal Bohar, Rohtak-124024

²Department of Botany, Baba Mastnath University, Asthal Bohar, Rohtak-124024

Article History: Received 19th March 2023; Accepted 22nd April 2023; Published 08th May 2023

ABSTRACT

A set of experiments was carried out to investigate the population dynamics of important insect pests that infest capsicum, bottle gourd and cucumber. Results revealed that aphid (*Aphis gossypii* Glover), whitefly (*Bemisia tabaci* Genn.) and leafhopper (*Amrasca biguttula*), and commenced the activity during the second and third week of October and November. The mean thrips peak population found in the second and third week of October in chilly (6.90) whereas third week for bottle gourd as well as cucumber (8.88 and 9.67 respectively). The average white fly density maximum is seen between the second of October in chilly, bottle gourd and cucumber (6.19, 8.88 and 9.67 respectively). Leaf hopper population also high in second week of October (5.02, 5.76 and 5.65) for chilly, bottle gourd and cucumber. Second and third week of October had maximum incidence of thrips and white fly population in bottle gourd as compare to chilly and cucumber.

Keywords: Aphid, Whitefly, Leafhopper, Chilly, Bottle gourd and Cucumber.

INTRODUCTION

Pepper (*Capsicum annum* L.), Cucumber (*Cucumis sativus* L.) and Bottle Guard (*Bactrocera cucurbitae* L.) are the one of the important economic, most popular and highly remunerative vegetable crops grown in most parts of the world; that for the nutritional values of fruits, mainly due to the fact that they are an excellent source of natural colors antioxidant and bioactive nutrients (Ellaithy *et al.*, 2015). Sucking insect pests like aphids, whitefly, thrips and leafhoppers attack the crop throughout the growth period resulting in the reduction of yields (Haque *et al.*, 2011). Unlike many of the field problems, insects and animal pests problems are peculiar to greenhouse cultivation. Aphids, two spotted spider mite, thrips, whitefly, leafhoppers, caterpillars, leaf miner, gall midge, nematodes and snails are serious problems on vegetable crops under protected condition and present the major factor limiting pepper production (Giraddi *et al.*, 2012). These insect pests caused economic damage either directly by sucking the juice of plants or indirectly by transmitted the plant viral diseases (Refaei *et al.*, 2016). Like other cucurbits, bottle gourd is also being subjected to damage by wide array of insect and non-insect pests, major being, fruit

fly, red pumpkin beetle, leaf miner, aphids, whitefly and mites (Tiwari *et al.*, 2012) right from the initial stages of the crop to harvest of the products in India. Such pests attacking pointed gourd in other countries have been reported so far by (Hasan *et al.*, 2015). The cucurbits are subjected to damage by a wide array of insect-pests. major being, fruit fly (Rai *et al.*, 2008, Sapkota *et al.*, 2010) Fruit fly, *B. cucurbitae* is one of the most destructive pests often rendering cultivation of bottle gourd unprofitable (Meena *et al.*, 2019). Thrips (*Scirtothrips dorsalis* Hood), whiteflies (*Bemisia tabaci* Genn), aphids (*Aphis gossypii* Glover) and yellow mites (*Polyphagotarsonemus latus* Banks) are the important sucking pests which contributed to reduction in the crop yield (Montasser *et al.*, 2011).

The piercing-sucking pests are very serious problem on pepper where both nymphs and adults suck the cell sap from leaves and tender parts thereby inducing premature senescence; also, excretes honeydew on which sooty mold grows and inhibits the photosynthesis in addition to indirect damage of spread viral diseases (Dogan *et al.*, 2016). So management interventions are required to save the yield loss. Coccinellids are used as an effective eater for sucking insect pest management (Elliott and Kieckhefer, 1990). The

*Corresponding Author: Dr. Kirpa Ram, Department of Botany, Baba Mastnath University, Asthal Bohar, Rohtak-124024, Email: dr.kirparamjangra@gmail.com.

beetles prey on a number of species of aphids on different host plants (Chowdhury *et al.*, 2008). The ladybeetles are predacious both at larval and adult stages and feed on pests such as aphids, brown plant hopper and thrips (Sapkota *et al.*, 2010). This paper deals with the seasonal incidence of sucking insect pests on Pepper, Cucumber and Bottle Guard and determines the role of predatory coccinellid beetles in suppressing sucking pest populations.

MATERIALS AND METHODS

The study was conducted during the 2019-20 crop season in the study area of Baba Mastnath University, Rohtak, utilizing the varieties Hisar Shakti (HC-44; Chilly), Kashi Ganga (Bottle Gourd), and Pusa Barkha (Cucumber). All of the agronomic techniques that were advised were implemented. Major insect pests were recorded at weekly intervals from five randomly chosen tagged plants per plot during the cropping period. The abundance of insect pests was counted to assess the frequency of various insect pests. The aphid, whitefly, and leaf Hoffer populations were observed by numbering both the nymphs and adults from three leaves (top, middle, and bottom) each plant. The

population of leafhoppers was collected using the sweep method and a hand net. Three sweeps per plot were considered and data were recorded in weekly intervals and analyzed by the OP-Stat by CCS Haryana Agriculture, University.

RESULT AND DISCUSSION

Significant different was observed on thrips population in every weeks of each month. The results presented in Table-1 revealed that maximum number of thrips incidence was found in second week of October for chilly (9.19), second week of November for bottle gourd and cucumber (11.54 and 12.00) respectively. The population of thrips was found significant at 5% CD level with incidence days, crop and weekly interval as well as interaction effect was also significant in all tested environment. Mean weekly maximum population of thrips was observed in third week for chilly, bottle guard and cucumber. The current findings are consistent with previous observations by Rajput *et al.*, 2017, who noted a greater occurrence of leafhoppers between the months of August and September.

Table 1. Weekly incidence of thrips on chilly, bottle gourd and cucumber.

Incidence Days	August	September	October	November	Mean (W)	
Chilly	Week-1	0.00	1.76	2.89	3.16	1.95
	Week-2	4.47	6.06	9.19	7.89	6.90
	Week-3	4.97	6.14	6.36	5.38	5.71
	Week-4	4.44	2.80	2.18	1.82	2.81
Bottle Gourd	Week-1	0.48	1.86	3.54	3.95	2.46
	Week-2	4.96	7.40	10.44	11.54	8.58
	Week-3	8.50	9.37	9.97	7.67	8.88
	Week-4	5.77	4.30	2.86	2.31	3.81
Cucumber	Week-1	0.62	2.14	3.94	4.68	2.85
	Week-2	5.12	8.02	10.81	12.00	8.99
	Week-3	9.15	10.47	10.94	8.13	9.67
	Week-4	5.78	5.62	3.41	3.04	4.46
Mean (C)	4.52	5.50	6.38	5.96		
Factors	C.D.		SE(d)		SE(m)	
Crop (C)	0.078		0.039		0.028	
Week (W)	0.090		0.045		0.032	
Interaction (C×W)	0.156		0.079		0.056	
Month (M)	0.090		0.045		0.032	
Interaction (C×M)	0.156		0.079		0.056	
Interaction (W×M)	0.180		0.091		0.064	
Interaction (C×W×M)	0.312		0.157		0.111	

Table 2. Weekly incidence of whitefly on chilly, bottle gourd and cucumber.

Incidence Days		August	September	October	November	Mean (W)
Chilly	Week-1	1.52	2.73	3.02	3.63	2.72
	Week-2	4.17	7.07	7.19	6.32	6.19
	Week-3	6.12	5.76	3.53	2.81	4.56
	Week-4	2.16	2.27	2.14	2.01	2.15
Bottle Gourd	Week-1	2.44	4.29	4.76	5.69	4.30
	Week-2	6.56	11.61	11.59	9.59	9.84
	Week-3	9.82	9.09	5.70	4.39	7.25
	Week-4	3.45	3.54	3.43	3.07	3.37
Cucumber	Week-1	2.72	5.46	6.54	7.14	5.47
	Week-2	9.14	11.49	12.11	10.78	10.88
	Week-3	10.49	9.87	7.25	5.58	8.30
	Week-4	4.41	3.76	3.54	3.10	3.70
Mean (C)		5.25	6.41	5.90	5.34	
Factors		C.D.		SE(d)		SE(m)
Crop (C)		0.067		0.034		0.024
Week (W)		0.078		0.039		0.028
Interaction (C×W)		0.135		0.068		0.048
Month (M)		0.078		0.039		0.028
Interaction (C×M)		0.135		0.068		0.048
Interaction (W×M)		0.155		0.078		0.055
Interaction (C×W×M)		0.269		0.136		0.096

Table 3. Weekly incidence of leafhopper on chilly, bottle gourd and cucumber.

Incidence Days		August	September	October	November	Mean (W)
Chilly	Week-1	1.17	1.55	1.87	2.15	1.68
	Week-2	3.31	5.93	5.71	5.11	5.02
	Week-3	4.93	4.21	2.17	1.99	3.33
	Week-4	1.89	1.45	1.05	0.61	1.25
Bottle Gourd	Week-1	1.96	2.35	2.65	2.82	2.44
	Week-2	4.07	6.71	6.40	5.84	5.76
	Week-3	5.58	5.06	2.87	2.84	4.09
	Week-4	2.64	2.18	1.84	1.37	2.01
Cucumber	Week-1	1.85	2.24	2.54	2.71	2.34
	Week-2	3.96	6.60	6.29	5.73	5.65
	Week-3	5.47	4.95	2.76	2.73	3.98
	Week-4	2.53	2.07	1.74	1.26	1.90
Mean (C)		3.28	3.77	3.16	2.93	
Factors		C.D.		SE(d)		SE(m)
Crop (C)		0.038		0.019		0.014
Week (W)		0.044		0.022		0.016
Interaction (C×W)		N/A		0.039		0.027
Month (M)		0.044		0.022		0.016
Interaction (C×M)		N/A		0.039		0.027
Interaction (W×M)		0.088		0.045		0.031
Interaction (C×W×M)		N/A		0.077		0.055

The white fly population in chilly commenced the activity second weeks of August, September, October and

November whereas third week of August, September, October and November for bottle gourd and cucumber

(Table-2). The peak population (7.19: chilly and 11.59; bottle gourd) was observed during the second week October and 12.11 in cucumber at second week of November. The mean population was recorded minimum (6.19, 9.84, 10.88) in chilly, bottle gourd and cucumber respectively in second week of every month. The population of thrips was found significant at 5% CD level with incidence days, crop and weekly interval as well as interaction effect was also significant in all tested environment. The findings are consistent with those of Patnaik *et al.*, 2004; Alim *et al.*, 2012 and Meena *et al.*, 2019 who indicated that the highest occurrence of this pest occurred in September to November in chilly, bottle gourd and cucumber respectively.

The leafhopper population in Chilly started activity in the second, third, and fourth weeks of August, September, October, and November, respectively, for bottle gourd and cucumber (Table-3). The highest population was 5.71 for chilly, 6.40 and 6.29 for cucumber. In the second week of each month, the mean population was observed as low as (5.02, 5.76, and 5.65) in chilly, bottle gourd, and cucumber. The thrips population was shown to be non-significant at the 5% CD level with incidence days, crop, and weekly intervals, as well as an interaction effect in all examined environments. The findings of Solangi *et al.*, (2008) and Deepika *et al.*, (2013) that leafhopper population was substantially and positively connected with session in chilly, bottle gourd, and cucumber are identical.

CONCLUSION

Population density, crop and week incidence showed significant association with each other in all tested crops. The data revealed that maximum population of Aphid, whitefly, and leafhopper was recorded on bottle gourd followed by cucumber, respectively. It is also concluded that based on population density of different insect pest of chilly, bottle gourd and cucumber second and third week of October thrips, white fly and leaf hopper population destruction is maximum.

ACKNOWLEDGEMENTS

Authors are also thankful to all the staff of Department of Zoology and Botany for their helpful co-operation and valuable suggestion.

REFERENCES

- Alim, M. B., Hossain, M. A., Khan, M., Khan, S. A. M., Islam, S. and Khalequzzaman, M. (2012). Seasonal variations of melon fly, *Bactrocera cucurbitae* (Coq.) (Diptera: Tephritidae) in different agricultural habitats of Bangladesh. *Journal of Agricultural Biological Science*, 7(11), 905-911.
- Chowdhury, S. P., Ahad, M. A., Amin, M. R., Hasan, M. S., (2008). Biology of lady beetle, *Micraspis discolors* (Fabr.) (Coccinellidae: Coleoptera). *International of Sustainable Crop Production*, 3, 39-44.
- Deepika, K., Roshan, L., Dahiya, K. K. and Bharti, Y. P. (2013). Population dynamics of sucking pest and its correlation with abiotic factors. *Agriways*, 1(1), 23-29.
- Dogan, A., Erler, F., Erkan, M., Ozgur Ates, A., Sule Sabanci, H. and Polat, E. (2016). Microbial based Production System: A Novel Approach for Plant Growth and Pest and Disease Management in Greenhouse-grown Peppers (*Capsicum annuum* L.). *Journal of Agricultural Science and Technology*, 18, 371-386.
- Ellaithy, A.Y.M., Shafeek, M. R., Hoda, Hussein, E., and Abo-Elella, G. M. (2015). Piercing sucking pests, growth and yield of sweet pepper cultivars as affected by alternative covers under plastic tunnel conditions. *International Journal of ChemTech Research*. 8(9), 149-161.
- Elliott, N. C., and Kieckhefer, R. W. (1990). Dynamics of aphidophagous coccinellid assemblages in small grain fields in eastern South Dakota. *Environmental Entomology*, 19, 1320-1329.
- Giraddi, R.S, Patil, R.K. and Wasin, M.A. (2012). Organic pest management in byadagi chilli. National symposium on Pest Management Organisation 10-11 December, 2012 UAS, Dharwad and KSDA, Bangalore, 2012, 72-76.
- Haque, M., Jyoti, Singh., N.K., Rath, S.S. and Ghosh, S. (2011) Epidemiology and seasonal dynamics of ixodid ticks of dairy animals of Punjab state, India. *Indian Journal of Animal Science*, 81, 661-664.
- Hasan, R., Hossain, M.K., Alam, N., Bashar, A., Islam, S. and Tarafder, M.J.A. (2015). Genetic divergenc in commercial cucumber (*Cucumis sativus* L.) genotypes. *Bangladesh Journal of Botany*, 44(2), 201-207.
- Meena, D. S., V. S. Acharya and Dinesh Kumar. (2019). Seasonal Incidence of Fruit Fly, *Bactrocera cucurbitae* on Bottle Gourd and their Correlation with Abiotic Factors. *International Journal Current Microbiology Applied Science*. 8(12), 381-388.
- Montasser, A.A., Marzouk, A.S., Hanafy, A.R.I. and Hassan, G.M. 2011. Seasonal fluctuation of the broad mite *Polyphagotarsonemus latus* (Acari: Tarsonemidae) and its predatory mites on some pepper cultivars in Egypt. *International Journal of Environmental Science and Engineering*, 2, 9-20.
- Patnaik, HP, Sarangi, PK and Mahapatra, P. (2004). Studies on the incidence of fruitflies and jassids on summer bitter gourd and their control. *Orissa Journal of Horticulture*, 32(2), 87-90.
- Rai, M., Pandey, S. and Kumar, S. (2008). Cucurbit research in India: aretrospect. Proc. X EUCARPIA

- Meeting on Genetics and Breeding of Cucurbitaceae, Avignon (France), May 21-24. Pp.292-93.
- Rajput, V.S., Prajapati, B.G., Pareek, A. and Patel, P.S. (2017). Studies on Population Dynamics of Major Insect Pests Infesting Chilli (*Capsicum annum* L.), *International Journal of Pure and Applied Bioscience*, 5(6), 1465-1470.
- Refaei, E.A., El-Sarand, E.A. and Khalifa, A.A. (2016). Population fluctuation of certain sucking insects and their natural enemies associated with seed watermelon, *Citrullus lanatus* at rasheed region, Beheira governorate, *Journal of Plant Protection Research*, 4(4), 61-69.
- Sapkota, R., Dahal, K. C. and Thapa, R.B. (2010). Damage assessment and management of cucurbit fruit flies in spring-summer squash. *Journal of Entomology and Nematology*, 2, 7-12.
- Solangi, G.S., Mahar, G.M., and Oad, F.C. (2008). Presence and absence of different insect predators against sucking insect pest of cotton. *Journal of Entomology*, 5(1), 31-37.
- Tiwari AK, Snehi SK, Khan MS, Sharma PK, Raj SK, Rao G P (2012). Molecular detection and identification of Tomato leaf curl New Delhivirus associated with yellow mosaic and leaf curl disease of *Luffacylindrica* in India. *Indian Phytopathology*, 65(1), 80-84.